

7-1-2008

# In Situ Sims, Micro-Baddeleyite U-Pb Dating Method for Mafic Rocks

K. R. Chamberlain

T. M. Harrison

A. K. Schmitt

L. M. Heaman

Susan Swapp

University of Wyoming, [swapp@uwyo.edu](mailto:swapp@uwyo.edu)

*See next page for additional authors*

Follow this and additional works at: [http://repository.uwyo.edu/geology\\_facpub](http://repository.uwyo.edu/geology_facpub)

 Part of the [Geology Commons](#)

---

## Publication Information

Chamberlain, K. R.; Harrison, T. M.; Schmitt, A. K.; Heaman, L. M.; Swapp, Susan; and Khudoley, A. K. (2008). "In Situ Sims, Micro-Baddeleyite U-Pb Dating Method for Mafic Rocks." *Geochimica Et Cosmochimica Acta* 72.12, A147-A147.

This Conference Proceeding is brought to you for free and open access by the Geology and Geophysics at Wyoming Scholars Repository. It has been accepted for inclusion in Geology and Geophysics Faculty Publications by an authorized administrator of Wyoming Scholars Repository. For more information, please contact [scholcom@uwyo.edu](mailto:scholcom@uwyo.edu).

---

**Authors**

K. R. Chamberlain, T. M. Harrison, A. K. Schmitt, L. M. Heaman, Susan Swapp, and A. K. Khudoley

## Mg and Si isotopes in the solar system

R. CHAKRABARTI\* AND S.B. JACOBSEN

Dept. of Earth and Planetary Sciences, Harvard University,  
Cambridge, MA 02138 (rama@eps.harvard.edu)

Chemical heterogeneity in our solar system is well-documented. Silicate portions of the Earth, Moon and Mars are different from the chondritic meteorites in their volatile element budgets and in Mg/Si. Isotopic compositions also vary between Earth and other Solar System bodies. Chemical and isotopic differences between Earth and bulk meteorites can be broadly attributed to the following processes: (1) Terrestrial fractionation, (2) Fractionation and incomplete mixing in the solar nebula prior to accretion and (3) Late injection of pre-solar material and its inhomogeneous distribution.

It has been suggested [1] that the Earth and Moon have heavier Si-isotopic compositions compared to meteorites, all of which have indistinguishable Si-isotopes, resulting from core formation on Earth. Several efforts have been initiated to resolve the Mg isotopic compositions of the bulk Earth and meteorites [2, 3] but available results are contradictory.

We have initiated a combined Mg- and Si-isotopic study in terrestrial and planetary samples to address the following questions: (1) What are the BSE Mg- and Si-isotopic compositions? (2) Does the BSE and average chondrites have resolvable differences in Si and Mg-isotopic compositions and (3) if they are different, what is the most plausible explanation? So far our Mg-isotopic data suggest that terrestrial olivines from peridotites and whole-rock basalts show similar Mg-isotopic compositions which are slightly lighter than those of whole-rock chondrites. Olivines from several pallasites show heavier Mg isotopic compositions compared to both chondrites and terrestrial samples. This difference suggests that pallasites, likely derived from the core-mantle boundary of early differentiated planetesimals, sampled Mg-isotopic heterogeneity in the solar nebula.

- [1] Georg, R. B. *et al.* (2007) *Nature* **447**, 1102-1106.  
[2] Wiechert, U. & Halliday, A. N. (2007) *EPSL* **256**, 360-371. [3] Teng F-Z *et al.* (2007) *EPSL* **261**, 84-92.

## *In situ* SIMS micro-baddeleyite U-Pb dating method for mafic rocks

K.R. CHAMBERLAIN<sup>1</sup>\*, T.M. HARRISON<sup>2</sup>,  
A.K. SCHMITT<sup>2</sup>, L.M. HEAMAN<sup>3</sup>, S.M. SWAPP<sup>1</sup>,  
AND A.K. KHUDDOLEY<sup>4</sup>

<sup>1</sup>Dept. of Geology and Geophysics, Univ. of Wyoming,  
Laramie, WY, 82071 USA

(\*correspondence: kchamber@uwyo.edu)

<sup>2</sup>Department of Earth and Space Sciences, UCLA,  
Los Angeles, CA 90095-1567, USA

<sup>3</sup>Dept. of Earth and Atmospheric Sciences, University of  
Alberta, Edmonton, AB T6G 2E3, Canada

<sup>4</sup>Geological Department, St. Petersburg State University,  
University nab. 7/9, St. Petersburg, 199034, Russia

Baddeleyite (ZrO<sub>2</sub>) is a favorite target for U-Pb dating of mafic and ultramafic rocks. Although samples with grains that are large enough (100-200 micron long) for traditional separation and dating methods are scarce, micro-baddeleyite crystals (<20 microns long) are nearly ubiquitous. With refined methods, it is possible to separate 10 to 20 micron long grains for ID-TIMS analysis.

We are currently developing an *in situ* technique to date micro-baddeleyite crystals using the CAMECA *ims* 1270 ion microprobe, avoiding the separation and dissolution steps of ID-TIMS and permitting analysis of even smaller grains (2-5µm). The ion microscope optics of the *ims* 1270 can reduce the effective sampling diameter of a 20 µm primary ion beam to a few to 10 µm. An aperture blocks undesired secondary ions, in particular common Pb ions emitted from host phases surrounding micro-baddeleyite, and the target crystals can be dated in polished thin sections. CAMECA *ims* 1270 analytical conditions also involve oxygen-flooding the sample chamber (~10<sup>5</sup> Torr) to enhance Pb secondary ion yields for baddeleyite by approximately 10-fold, and a UO<sub>2</sub>/U vs. U/Pb relative sensitivity calibration. Three baddeleyite standards analyzed under these conditions yielded U/Pb reproducibilities that only marginally exceeded those determined for NIST610 glass, suggesting that crystal orientation related bias in U-Pb sensitivity is minor to absent. Proterozoic-aged standards yielded 2 sigma precisions on <sup>207</sup>Pb/<sup>206</sup>Pb dates from 1% to 0.2%. Preliminary results for micro-baddeleyite from Ordovician and Mesoproterozoic-aged mafic sills analyzed *in situ* yield U-Pb ages roughly equivalent in precision to those achieved on larger standard baddeleyite grains. Precisions on Phanerozoic <sup>206</sup>Pb/<sup>238</sup>U dates are ~6%; Precambrian <sup>207</sup>Pb/<sup>206</sup>Pb date precisions are 1% or less. The *in situ* technique is applicable to most unmetamorphosed to low-grade mafic rocks including dikes, sills, fine-grained volcanic rocks and extra-terrestrial rocks, greatly increasing the number of mafic samples that are dateable by U-Pb methods.