

2008

## A note on the largest eigenvalue of non-regular graphs

Bolian Liu  
liubl@scnu.edu.cn

Gang Li

Follow this and additional works at: <http://repository.uwyo.edu/ela>

---

### Recommended Citation

Liu, Bolian and Li, Gang. (2008), "A note on the largest eigenvalue of non-regular graphs", *Electronic Journal of Linear Algebra*, Volume 17.  
DOI: <https://doi.org/10.13001/1081-3810.1249>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact [scholcom@uwyo.edu](mailto:scholcom@uwyo.edu).

## A NOTE ON THE LARGEST EIGENVALUE OF NON-REGULAR GRAPHS\*

BOLIAN LIU<sup>†</sup> AND GANG LI<sup>†</sup>

**Abstract.** The spectral radius of connected non-regular graphs is considered. Let  $\lambda_1$  be the largest eigenvalue of the adjacency matrix of a graph  $G$  on  $n$  vertices with maximum degree  $\Delta$ . By studying the  $\lambda_1$ -extremal graphs, it is proved that if  $G$  is non-regular and connected, then  $\Delta - \lambda_1 > \frac{\Delta + 1}{n(3n + \Delta - 8)}$ . This improves the recent results by B.L. Liu et al.

**AMS subject classifications.** 05C50, 15A48.

**Key words.** Spectral radius, Non-regular graph,  $\lambda_1$ -extremal graph, Maximum degree.

**1. Introduction.** Let  $G = (V, E)$  be a simple graph on vertex set  $V$  and edge set  $E$ , where  $|V| = n$ . The eigenvalues of the adjacency matrix of  $G$  are called the eigenvalues of  $G$ . The largest eigenvalue of  $G$ , denoted by  $\lambda_1(G)$ , is called the spectral radius of  $G$ . Let  $D$  denote the diameter of  $G$ . We suppose throughout the paper that  $G$  is a simple graph. For any vertex  $u$ , let  $\Gamma(u)$  be the set of all neighbors of  $u$  and  $d(u) = |\Gamma(u)|$  be the degree of  $u$ . A nonincreasing sequence  $\pi = (d_1, d_2, \dots, d_n)$  of non-negative integers is called (*connected*) *graphic* if there exists a (connected) simple graph on  $n$  vertices, for which  $d_1, d_2, \dots, d_n$  are the degrees of its vertices. Let  $\Delta$  and  $\delta$  be the maximum and minimum degree of vertices of  $G$ , respectively. A graph is called regular if  $d(u) = \Delta$  for any  $u \in V$ . It is easy to see that the spectral radius of a regular graph is  $\Delta$  with  $(1, 1, \dots, 1)^T$  as a corresponding eigenvector. We will use  $G - e$  ( $G + e$ ) to denote the graph obtained from  $G$  by deleting (adding) the edge  $e$ . For other notations in graph theory, we follow from [2].

Stevanović [8] first found a lower bound of  $\Delta - \lambda_1$  for the connected non-regular graphs. Then the results from [8] were improved in [9, 4, 7, 3]. In [4, 7], the authors showed that

$$\Delta - \lambda_1 \geq \frac{1}{n(D + 1)} \quad ([4, 7]) \quad (1.1)$$

---

\*Received by the editors November 21, 2007. Accepted for publication February 15, 2008. Handling Editor: Stephen J. Kirkland.

<sup>†</sup>School of Mathematics Sciences, South China Normal University, Guangzhou, 510631, P.R. China (liubl@scnu.edu.cn, tepal.li@sohu.com). This work was supported by the National Natural Science Foundation of China (No.10771080) and by DF of the Ministry of Education of China (No.20070574006).

and

$$D \leq \frac{3n + \Delta - 5}{\Delta + 1} \quad ([7]). \quad (1.2)$$

B.L. Liu et al. obtained

$$\Delta - \lambda_1 \geq \frac{\Delta + 1}{n(3n + 2\Delta - 4)} \quad ([7]). \quad (1.3)$$

Recently, S.M. Cioabă [3] improved (1.1) as follows:

$$\Delta - \lambda_1 > \frac{1}{nD} \quad ([3]). \quad (1.4)$$

Thus combining (1.2) and (1.4), the inequality (1.3) can be improved as follows:

$$\Delta - \lambda_1 > \frac{\Delta + 1}{n(3n + \Delta - 5)}. \quad (1.5)$$

In this note we improve the inequality (1.2) on  $\lambda_1$ -extremal graphs. Furthermore, we obtain the following inequality which improves (1.5).

$$\Delta - \lambda_1 > \frac{\Delta + 1}{n(3n + \Delta - 8)}.$$

**2. Preparation.** Firstly, we state a well-known result which is just Frobenius's theorem applied to graphs.

**LEMMA 2.1.** *Let  $G$  be a connected graph and  $\lambda_1(G)$  be its spectral radius. Then  $\lambda_1(G + uv) > \lambda_1(G)$  for any  $uv \notin E$ .*

**DEFINITION 2.2.** [7] Let  $G$  be a connected non-regular graph. Then the graph  $G$  is called  $\lambda_1$ -extremal if  $\lambda_1(G) \geq \lambda_1(G')$  for any other connected non-regular graph  $G'$  with the same number of vertices and maximum degree as  $G$ .

**THEOREM 2.3.** *Let  $G$  be a  $\lambda_1$ -extremal graph on  $n$  vertices with maximum degree  $\Delta$ . Define*

$$V_{<\Delta} = \{u : u \in V \text{ and } d(u) < \Delta\}.$$

*Then  $G$  must have one of the following properties:*

- (1)  $|V_{<\Delta}| \geq 2$  and  $V_{<\Delta}$  induces a complete graph.
- (2)  $|V_{<\Delta}| = 1$ .
- (3)  $V_{<\Delta} = \{u, v\}$ ,  $uv \notin E(G)$  and  $d(u) = d(v) = \Delta - 1$ .

*Proof.* By contradiction, suppose that  $G$  is a  $\lambda_1$ -extremal graph without properties (1), (2) and (3). It follows that  $|V_{<\Delta}| \geq 2$ . Then there are two cases.

**Case 1:**  $V_{<\Delta} = \{u, v\}$ ,  $uv \notin E(G)$ ,  $d(u) < \Delta - 1$  and  $d(v) \leq \Delta - 1$ . Then the graph  $G + uv$  has the same maximum degree as  $G$ . By Lemma 2.1, we obtain  $\lambda_1(G + uv) > \lambda_1(G)$ , contradicting the choice of  $G$ .

**Case 2:**  $|V_{<\Delta}| > 2$  and  $V_{<\Delta}$  does not induce a complete graph. Then there exist two vertices  $u, v \in V_{<\Delta}$  and  $uv \notin E(G)$ . Similarly arguing to case 1, we obtain  $\lambda_1(G + uv) > \lambda_1(G)$ , contradicting the choice of  $G$ .

Combining the above two cases, the proof follows.  $\square$

Using the properties mentioned in Theorem 2.3, we give the following definition.

DEFINITION 2.4. Let  $G$  be a connected non-regular graph on  $n$  vertices with maximum degree  $\Delta$ . Then

- the graph  $G$  is called *type-I* if it has property (1),
- the graph  $G$  is called *type-II* if it has property (2),
- the graph  $G$  is called *type-III* if it has property (3).

LEMMA 2.5. [6] Let  $G$  be a simple connected graph with  $n$  vertices,  $m$  edges and spectral radius  $\lambda_1(G)$ . Then

$$\lambda_1(G) \leq \frac{\delta - 1 + \sqrt{(\delta + 1)^2 + 4(2m - \delta n)}}{2}$$

and equality holds if and only if  $G$  is either a regular graph or a graph in which each vertex has degree either  $\delta$  or  $n - 1$ .

We first consider the  $\lambda_1$ -extremal graphs with  $\Delta = 2$  or  $\Delta = n - 1$ . When  $\Delta = 2$ , the  $\lambda_1$ -extremal graph is the path with  $\lambda_1(P_n) = 2\cos(\frac{\pi}{n+1})$ . When  $\Delta = n - 1$ , similarly arguing to Theorem 2.3, we know that the  $\lambda_1$ -extremal graph is  $K_n - e$ . By Lemma 2.5, we obtain

$$\lambda_1(K_n - e) = \frac{n - 3 + \sqrt{(n + 1)^2 - 8}}{2}. \tag{2.1}$$

Theorem 2.3 shows that the  $\lambda_1$ -extremal graphs must be type-I, type-II or type-III, but in what follows, we will prove that when  $2 < \Delta < n - 1$ , any type-III graph is not  $\lambda_1$ -extremal.

LEMMA 2.6. [5] Let  $\pi = (d_1, d_2, \dots, d_n)$  be a nonincreasing sequence of non-negative integers. Then  $\pi$  is graphic if and only if

$$\sum_{i=1}^n d_i \text{ is even and } \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\}, \text{ for all } k = 1, 2, \dots, n-1. \tag{2.2}$$

LEMMA 2.7. Let  $\pi = (d_1, d_2, \dots, d_n)$  be a nonincreasing sequence of positive

integers and  $d_{n-1} \geq 2$ ,  $d_n \geq 1$ . Then  $\pi$  is graphic if and only if it is connected graphic.

*Proof.* If  $\pi$  is connected graphic, then it is obviously graphic. Conversely, suppose that  $\pi$  is graphic and  $G$  is a disconnected graph with the degree sequence  $\pi$ . Without loss of generality, suppose that  $G$  has two components  $G_1$  and  $G_2$ . Noticing  $d_{n-1} \geq 2$  and  $d_n \geq 1$ , we suppose that any vertex in  $G_1$  has degree at least two and any edge  $u_2v_2 \in E(G_2)$ . Then it follows that there exists one edge  $u_1v_1$  in  $G_1$  which is not the cut edge, i.e.  $G_1 - u_1v_1$  is still connected. Otherwise,  $G_1$  is a tree, a contradiction. Consider  $G' = G - u_1v_1 - u_2v_2 + u_1u_2 + v_1v_2$ . It is easy to see that  $G'$  is a connected graph with the degree sequence  $\pi$ .  $\square$

**LEMMA 2.8.** Let  $\pi = (d_1, d_2, \dots, d_n) = (\Delta, \Delta, \dots, \Delta, \Delta - 1, \Delta - 1)$  and  $\pi' = (d'_1, d'_2, \dots, d'_n) = (\Delta, \Delta, \dots, \Delta, \Delta - 2)$  with  $2 < \Delta < n - 1$ . If  $\pi$  is connected graphic, then  $\pi'$  is connected graphic.

*Proof.* Since  $2 < \Delta < n - 1$ , we obtain  $d'_{n-1} \geq 3$  and  $d'_n \geq 1$ . Then by Lemma 2.7, we need only to prove that  $\pi'$  is graphic. Let  $G$  be a connected graph with the degree sequence  $\pi$ . Since  $\pi$  is graphic, by Lemma 2.6, we obtain

$$\sum_{i=1}^n d_i \text{ is even and } \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\}, \text{ for all } k = 1, 2, \dots, n-1.$$

For  $\pi'$  we will prove that (2.2) is still true. Obviously,  $\sum_{i=1}^n d'_i = \sum_{i=1}^n d_i$  is even. Then we need only to prove that the inequality is true. We split our proof into four cases.

**Case 1:**  $1 \leq k \leq \Delta - 2$ , then  $k \leq n - 4$  and

$$\begin{aligned} \sum_{i=1}^k d'_i &= \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\} \\ &= k(k-1) + k(n-k) \\ &= k(k-1) + \sum_{i=k+1}^n \min\{k, d'_i\}. \end{aligned}$$

**Case 2:**  $1 < k = \Delta - 1$ , then  $k \leq n - 3$  and

$$k(k-1) + k(n-k) - \sum_{i=1}^k d_i = k(k-1) + k(n-k) - k\Delta = k(n-k-2) \geq k > 1.$$

Thus

$$\sum_{i=1}^k d'_i = \sum_{i=1}^k d_i < k(k-1) + k(n-k) - 1 = k(k-1) + \sum_{i=k+1}^n \min\{k, d'_i\}.$$

**Case 3:**  $\Delta \leq k \leq n - 2$ , then

$$\begin{aligned} \sum_{i=1}^k d'_i &= \sum_{i=1}^k d_i \leq k(k-1) + \sum_{i=k+1}^n \min\{k, d_i\} \\ &= k(k-1) + (n-k-2)\Delta + 2\Delta - 2 \\ &= k(k-1) + \sum_{i=k+1}^n \min\{k, d'_i\}. \end{aligned}$$

**Case 4:**  $k = n - 1$ , then

$$\begin{aligned} k(k-1) + \Delta - \sum_{i=1}^k d_i &= (n-1)(n-2) + \Delta - [(n-1)\Delta - 1] \\ &= (n-2)(n-1-\Delta) + 1 \geq 4, \end{aligned}$$

where the last inequality holds since  $2 < \Delta < n - 1$ . Hence

$$\begin{aligned} \sum_{i=1}^k d'_i &= \sum_{i=1}^k d_i + 1 < k(k+1) + \Delta - 2 \\ &= k(k-1) + \sum_{i=k+1}^n \min\{k, d'_i\}. \end{aligned}$$

Combining the above four cases, the inequality is true. Then by Lemma 2.6, the  $\pi'$  is graphic. This completes the proof.  $\square$

As we know, *majorization* on degree sequences is defined as follows: for two sequences  $\pi = (d_1, d_2, \dots, d_n)$ ,  $\pi' = (d'_1, d'_2, \dots, d'_n)$  we write  $\pi \preceq \pi'$  if and only if  $\sum_{i=1}^n d_i = \sum_{i=1}^n d'_i$  and  $\sum_{i=1}^j d_i \leq \sum_{i=1}^j d'_i$  for all  $j = 1, 2, \dots, n$ . We claim that  $G$  has the *greatest maximum* eigenvalue if  $\lambda_1(G) \geq \lambda_1(G')$  for any other graph  $G'$  in the class  $\mathcal{C}_\pi$ , where  $\mathcal{C}_\pi = \{G : G \text{ is a connected graph with the degree sequence } \pi\}$ .

**LEMMA 2.9.** [1] *Let  $\pi$  and  $\pi'$  be two distinct degree sequences with  $\pi \preceq \pi'$ . Let  $G$  and  $G'$  be graphs with the greatest maximum eigenvalues in classes  $\mathcal{C}_\pi$  and  $\mathcal{C}_{\pi'}$ , respectively. Then  $\lambda_1(G) < \lambda_1(G')$ .*

**THEOREM 2.10.** *Let  $G$  be a connected graph with degree sequence*

$$\pi = (\Delta, \Delta, \dots, \Delta, \Delta - 1, \Delta - 1)$$

*and  $2 < \Delta < n - 1$ . Then there exists a connected graph  $G'$  with degree sequence  $\pi' = (\Delta, \Delta, \dots, \Delta, \Delta - 2)$  such that  $\lambda_1(G) < \lambda_1(G')$ .*

*Proof.* By Lemma 2.8, there exists a connected graph  $G'$  with degree sequence  $\pi' = (\Delta, \Delta, \dots, \Delta, \Delta - 2)$ . We suppose that  $G$  ( $G'$ ) is the graph with the greatest

maximum eigenvalue in  $\mathcal{C}_\pi$  ( $\mathcal{C}_{\pi'}$ ). It is obvious that  $\pi \preceq \pi'$ . By Lemma 2.9, we obtain  $\lambda_1(G) < \lambda_1(G')$ .  $\square$

**THEOREM 2.11.** *Let  $G$  be a  $\lambda_1$ -extremal graph on  $n$  vertices with the maximum degree  $\Delta$  and  $2 < \Delta < n - 1$ . Then  $G$  must be either type-I or type-II.*

*Proof.* Suppose that  $G$  is a type-III graph with the greatest maximum eigenvalue in class  $\mathcal{C}_\pi$ , where  $\pi = (\Delta, \Delta, \dots, \Delta, \Delta - 1, \Delta - 1)$ . By Theorem 2.10, there exists a graph  $G'$  with degree sequence  $\pi' = (\Delta, \Delta, \dots, \Delta, \Delta - 2)$  and greatest maximum eigenvalue in class  $\mathcal{C}'_\pi$  such that  $\lambda_1(G') > \lambda_1(G)$ . It follows that  $G$  is not  $\lambda_1$ -extremal.  $\square$

**REMARK.** Although Theorem 2.11 shows that the  $\lambda_1$ -extremal graph with  $2 < \Delta < n - 1$  must be type-I or type-II, there exist some graphs with property (1) or (2) which are not  $\lambda_1$ -extremal. Let  $G_1, G_2$  be connected graphs with degree sequences  $(3, 3, 3, 3, 2, 2), (5, 5, 5, 5, 5, 5, 2)$ , respectively. Clearly  $G_1$  ( $G_2$ ) is a type-I (type-II) graph. However, by checking the Table 1 of [7], we know they are not the  $\lambda_1$ -extremal graphs. After some computer experiments, we give a conjecture about the  $\lambda_1$ -extremal graphs as follows:

**CONJECTURE 2.12.** *Let  $G$  be a connected non-regular graph on  $n$  vertices and  $2 < \Delta < n - 1$ . Then  $G$  is  $\lambda_1$ -extremal if and only if  $G$  is a graph with the greatest maximum eigenvalue in classes  $\mathcal{C}_\pi$  and  $\pi = (\Delta, \Delta, \dots, \Delta, \delta)$ , where*

$$\delta = \begin{cases} (\Delta - 1), & \text{when } n\Delta \text{ is odd,} \\ (\Delta - 2), & \text{when } n\Delta \text{ is even.} \end{cases}$$

### 3. Main Results.

**THEOREM 3.1.** *Let  $G$  be a type-I or type-II graph on  $n$  vertices with diameter  $D$ . Then*

$$D \leq \frac{3n + \Delta - 8}{\Delta + 1}. \quad (3.1)$$

*Proof.* Since  $G$  is a type-I or type-II graph, we have  $\Delta \geq 3$ . Let  $u, v$  be two vertices at distance  $D$  and  $P : u = u_0 \leftrightarrow u_1 \leftrightarrow \dots \leftrightarrow u_D = v$  be the shortest path connecting  $u$  and  $v$ . We first claim that  $|V_{<\Delta} \cap V(P)| \leq 2$ . Otherwise,  $G$  must be a type-I graph and suppose  $\{u_p, u_q, u_r\} \subseteq V(P) \cap V_{<\Delta}$  with  $p < q < r$ . Then by definition of type-I graph, we obtain that  $u_p u_q, u_q u_r$  and  $u_p u_r \in E(G)$ . Therefore,  $P$  is not the shortest path connecting  $u$  and  $v$ , a contradiction.

Then there are two cases.

**Case 1:**  $V_{<\Delta} \cap V(P) = \emptyset$ . Define  $T = \{i : i \equiv 0 \pmod{3} \text{ and } i \leq (D - 3)\} \cup \{D\}$ . Thus  $|T| = \lceil \frac{D+1}{3} \rceil$ . Let  $d(u_i, u_j)$  denote the distance between  $u_i$  and  $u_j$ . Since  $P$  is

the shortest path connecting  $u$  and  $v$ , we have  $d(u_i, u_j) \geq 3$  and  $\Gamma(u_i) \cap \Gamma(u_j) = \emptyset$  for any distinct  $i, j \in T$ . Notice that  $u_i \in V(P)$  for any  $i \in T$ . We obtain

$$|\Gamma(u_i) - V(P)| = \begin{cases} \Delta - 1, & \text{if } i \in \{0, D\}, \\ \Delta - 2, & \text{otherwise.} \end{cases}$$

Then

$$\begin{aligned} n &\geq |V(P)| + \sum_{i \in T} |\Gamma(u_i) - V(P)| \\ &\geq D + 1 + (|T| - 2)(\Delta - 2) + 2(\Delta - 1) \\ &\geq D + 1 + \left(\frac{D+1}{3} - 2\right)(\Delta - 2) + 2(\Delta - 1). \end{aligned}$$

Thus

$$D \leq \frac{3n - \Delta - 7}{\Delta + 1}.$$

**Case 2:** Either  $V_{<\Delta} \cap V(P) = \{u_p, u_q\}$  with  $q = p + 1$  or  $V_{<\Delta} \cap V(P) = \{u_p\}$ . The proof is similar to the proof of [7]. We obtain the same result

$$D \leq \frac{3n + \Delta - 8}{\Delta + 1}.$$

Combining the above two cases, the proof follows.  $\square$

**LEMMA 3.2.** [3] *Let  $G$  be a connected non-regular graph on  $n$  vertices with maximum degree  $\Delta$  and diameter  $D$ . Then*

$$\Delta - \lambda_1 > \frac{1}{nD}.$$

**THEOREM 3.3.** *Let  $G$  be a connected non-regular graph on  $n$  vertices with maximum degree  $\Delta$ . Then*

$$\Delta - \lambda_1 > \frac{\Delta + 1}{n(3n + \Delta - 8)}.$$

*Proof.* Without loss of generality, we suppose that  $G$  is a  $\lambda_1$ -extremal graph. Since  $G$  is connected and non-regular, then  $n \geq 3$  and  $\Delta \geq 2$ . When  $\Delta = n - 1$  and  $n \geq 5$ , the  $\lambda_1$ -extremal graph is  $K_n - e$  with  $D = 2$ . Then by Lemma 3.2, we obtain

$$\lambda_1(K_n - e) < \Delta - \frac{1}{2n} < \Delta - \frac{\Delta + 1}{n(3n + \Delta - 8)}.$$



When  $\Delta = n - 1$  and  $n = 3$ , the  $\lambda_1$ -extremal graph is  $P_3$  with  $\lambda_1(P_3)=1.4142$ . When  $\Delta = n - 1$  and  $n = 4$ , the  $\lambda_1$ -extremal graph is  $K_4 - e$  with  $\lambda_1(K_4 - e)=2.5616$ . By direct calculation, we know that the inequality is true. When  $2 < \Delta < n - 1$ , applying Theorem 3.1 and Lemma 3.2, we obtain the result. When  $\Delta = 2$  and  $n > 3$ , the  $\lambda_1$ -extremal graph is  $P_n$ . By adding some edges to  $P_n$ , we can attain  $K_n - e$ . Then following the Lemma 2.1, we obtain  $\lambda_1(P_n) < \lambda_1(K_n - e)$ . This completes the proof.  $\square$

**Acknowledgement.** The authors would like to thank an anonymous referee for valuable comments and suggestions that improved our presentation.

#### REFERENCES

- [1] T. Biyikoglu and J. Leydold. Largest eigenvalue of degree sequences. Available at <http://arxiv.org/abs/math.CO/0605294>.
- [2] J.A. Bondy and U.S.R. Mury. *Graph Theory with Application*. North-Holland, New York, 1976.
- [3] S.M. Cioabă. The spectral radius and the maximum degree of irregular graphs. *Electron. J. Combin.*, 14 (2007), R38.
- [4] S.M. Cioabă, D.A. Gregory, and V. Nikiforov. Note: extreme eigenvalues of nonregular graphs. *J. Combin. Theory Ser. B*, 97:483–486, 2007.
- [5] P. Erdős and T. Gallai. Graphs with prescribed degree of vertices (in Hungarian). *Mat. Lapok* 11, 1960.
- [6] Y. Hong, J.L. Shu, and K. Fang. A sharp upper bound of the spectral radius of graphs. *J. Combin. Theory Ser. B*, 81:177–183, 2001.
- [7] B.L. Liu, J. Shen, and X.M. Wang. On the largest eigenvalue of non-regular graphs. *J. Combin. Theory Ser. B*, 97:1010–1018, 2007.
- [8] D. Stevanović. The largest eigenvalue of nonregular graphs. *J. Combin. Theory Ser. B*, 91:143–146, 2004.
- [9] X.D. Zhang. Eigenvectors and eigenvalues of non-regular graphs. *Linear Algebra Appl.*, 409:79–86, 2005.