

2010

A. Horn's result on matrices with prescribed singular values and eigenvalues

Tin-Yau Tam

Follow this and additional works at: <http://repository.uwyo.edu/ela>

Recommended Citation

Tam, Tin-Yau. (2010), "A. Horn's result on matrices with prescribed singular values and eigenvalues", *Electronic Journal of Linear Algebra*, Volume 21.

DOI: <https://doi.org/10.13001/1081-3810.1411>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact scholcom@uwyo.edu.

A. HORN'S RESULT ON MATRICES WITH PRESCRIBED SINGULAR VALUES AND EIGENVALUES*

TIN-YAU TAM[†]

Abstract. We give a new proof of a classical result of A. Horn on the existence of a matrix with prescribed singular values and eigenvalues.

Key words. Eigenvalues, singular values.

AMS subject classifications. 15A45, 15A18.

Let $A \in \mathbb{C}_{n \times n}$ and let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A arranged in the order $|\lambda_1| \geq \dots \geq |\lambda_n|$. The singular values of A are the nonnegative square roots of the eigenvalues of the positive semi-definite matrix A^*A and are denoted by $s_1 \geq \dots \geq s_n$. Weyl's inequalities [7] provide a very nice relation between the eigenvalues and singular values of A :

$$(1.1) \quad \prod_{j=1}^k |\lambda_j| \leq \prod_{j=1}^k s_j, \quad k = 1, \dots, n-1,$$

$$(1.2) \quad \prod_{j=1}^n |\lambda_j| = \prod_{j=1}^n s_j.$$

The equality follows from two ways of expressing the absolute value of the determinant of A . A. Horn [2] established the converse of Weyl's result.

THEOREM 1.1. (A. Horn) If $|\lambda_1| \geq \dots \geq |\lambda_n|$ and $s_1 \geq \dots \geq s_n$ satisfy (1.1) and (1.2), then there exists $A \in \mathbb{C}_{n \times n}$ such that $\lambda_1, \dots, \lambda_n$ are the eigenvalues and s_1, \dots, s_n are the singular values of A .

Horn's original proof is divided into two cases: (i) $s_n \neq 0$ (the nonsingular case) and (ii) $s_n = 0$ (the singular case. There is a typo: $C_{m,m+1} = \gamma$ and $C_{i,i+1} = \alpha_i$ should be $C_{m+1,m} = \gamma$ and $C_{i+1,i} = \alpha_i$ on [2, p.6]). In this note we provide a new proof of Horn's result. Our proof differs from Horn's proof in two ways that (i) our proof is divided into two cases according to $\lambda_1 = 0$ and $\lambda_1 \neq 0$, and (ii) our induction technique is different. It is very much like Chan and Li's technique [1] (the same

*Received by the editors on June 10, 2009. Accepted for publication on July 31, 2010. Handling Editors: Roger A. Horn and Fuzhen Zhang.

[†]Department of Mathematics and Statistics, Auburn University, AL 36849-5310, USA (tamtiny@auburn.edu).

technique is also used in [8]) for proving another result of Horn [3] (on the diagonal entries and eigenvalues of a Hermitian matrix): ours is multiplicative and Chan and Li's is additive. See [4, Section 3.6] for a proof of Theorem 1.1 using the result of Horn [3]. Also see [5] for an extension of Weyl-Horn's result and a numerically stable construction of A . A technique similar to that of Chan and Li can be found in Thompson's earlier work [6] on the diagonal entries and singular values of a square matrix.

Proof. We divide the proof into two cases: nilpotent or not.

Case i: $\lambda_1 = 0$. Then $s_n = 0$ by (1.2) and we choose

$$A := \begin{pmatrix} 0 & s_1 & & \\ & 0 & \ddots & \\ & & & s_{n-1} \\ & & & 0 \end{pmatrix}$$

Case ii: $\lambda_1 \neq 0$. We will use induction on n . When $n = 2$, the matrix

$$A = \begin{pmatrix} \lambda_1 & \mu \\ 0 & \lambda_2 \end{pmatrix}$$

has singular values $s_1 \geq s_2$ if we set

$$\mu := (s_1^2 + s_2^2 - |\lambda_1|^2 - |\lambda_2|^2)^{1/2}.$$

Suppose that the statement of Theorem 1.1 is true for $\lambda_1 \neq 0$ when $n = m \geq 2$. Let $n = m + 1$ and let $j \geq 2$ be the largest index such that $s_{j-1} \geq |\lambda_1| \geq s_j$. Clearly $s_1 \geq \max\{|\lambda_1|, s_1 s_j / |\lambda_1|\} \geq \min\{|\lambda_1|, s_1 s_j / |\lambda_1|\}$. Then there exist 2×2 unitary matrices U_1 and V_1 such that

$$U_1 \begin{pmatrix} s_1 & \\ & s_j \end{pmatrix} V_1 = \begin{pmatrix} \lambda_1 & \mu' \\ 0 & s_1 s_j / |\lambda_1| \end{pmatrix},$$

where $\mu' = (s_1^2 + s_j^2 - |\lambda_1|^2 - s_1^2 s_j^2 / |\lambda_1|^2)^{1/2}$. Set $U_2 := U_1 \oplus I_{m-1}$, $V_2 := V_1 \oplus I_{m-1}$. Then

$$\begin{aligned} A_1 &:= U_2 \operatorname{diag}(s_1, s_j, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}) V_2 \\ &= \begin{pmatrix} \lambda_1 & \mu' \\ 0 & s_1 s_j / |\lambda_1| \end{pmatrix} \oplus \operatorname{diag}(s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}). \end{aligned}$$

It suffices to show that $(s_1 s_j / |\lambda_1|, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1})$ and $(\lambda_2, \dots, \lambda_{m+1})$ satisfy (1.1) and (1.2). Since $s_{j-1} \geq |\lambda_1| \geq |\lambda_2|$,

$$|\lambda_2| \leq \max\{s_1 s_j / |\lambda_1|, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1}\}.$$

Moreover

$$\prod_{i=2}^k |\lambda_i| \leq |\lambda_1|^{k-1} \leq \prod_{i=2}^k s_i, \quad k = 2, \dots, j-1,$$

$$\prod_{i=2}^k |\lambda_i| = \frac{1}{|\lambda_1|} \prod_{i=1}^k |\lambda_i| \leq \frac{s_1 s_j}{|\lambda_1|} \prod_{i=2, i \neq j}^k s_i, \quad k = j, \dots, m, \quad \text{by (1.1)}$$

$$\prod_{i=2}^{m+1} |\lambda_i| = \frac{s_1 s_j}{|\lambda_1|} \prod_{i=2, i \neq j}^{m+1} s_i \quad \text{by (1.2)}.$$

We consider two cases: (a) $\lambda_2 = 0$ and apply Case i. (b) $\lambda_2 \neq 0$ and apply the induction hypothesis in Case ii. For both cases, there exist $m \times m$ unitary matrices U_3, V_3 such that

$$U_3 \operatorname{diag} \left(\frac{s_1 s_2}{|\lambda_1|}, s_2, \dots, s_{j-1}, s_{j+1}, \dots, s_{m+1} \right) V_3$$

is upper triangular with diagonal $(\lambda_2, \dots, \lambda_{m+1})$. Then $A = U_4 A_1 V_4$ is the desired matrix, where $U_4 := 1 \oplus U_3, V_4 := 1 \oplus V_3$. \square

Acknowledgment: The author is very thankful to Roger Horn for many valuable comments and suggestions.

REFERENCES

- [1] N.N. Chan and K.H. Li. Diagonal elements and eigenvalues of a real symmetric matrix. *J. Math. Anal. Appl.*, **91** (1983) 562–566.
- [2] A. Horn. On the eigenvalues of a matrix with prescribed singular values. *Proc. Amer. Math. Soc.*, **5** (1954) 4–7.
- [3] A. Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *Amer. J. Math.*, **76** (1954), 620–630.
- [4] R.A. Horn and C.R. Johnson. *Topics in Matrix Analysis*. Cambridge Univ. Press, 1991.
- [5] C.K. Li and R. Mathias. Construction of matrices with prescribed singular values and eigenvalues. *BIT* **41** (2001) 115–126.
- [6] R.C. Thompson. Singular values, diagonal elements, and convexity. *SIAM J. Appl. Math.*, **32** (1977) 39–63.
- [7] H. Weyl. Inequalities between the two kinds of eigenvalues of a linear transformation. *Proc. Nat. Acad. Sci. U.S.A.*, **35** (1949) 408–411.
- [8] H. Zha and Z. Zhang. A note on constructing a symmetric matrix with specified diagonal entries and eigenvalues. *BIT*, **35** (1995) 448–452.