

2012

A four-vertex, quadratic, spanning forest polynomial identity

Aleksandar Vlasev

Karen Yeats

karen_yeats@sfu.ca

Follow this and additional works at: <http://repository.uwyo.edu/ela>

Recommended Citation

Vlasev, Aleksandar and Yeats, Karen. (2012), "A four-vertex, quadratic, spanning forest polynomial identity", *Electronic Journal of Linear Algebra*, Volume 23.

DOI: <https://doi.org/10.13001/1081-3810.1566>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact scholcom@uwyo.edu.

A FOUR-VERTEX, QUADRATIC, SPANNING FOREST POLYNOMIAL IDENTITY*

ALEKSANDAR VLASEV[†] AND KAREN YEATS[‡]

Abstract. The classical Dodgson identity can be interpreted as a quadratic identity of spanning forest polynomials, where the spanning forests used in each polynomial are defined by how three marked vertices are divided among the component trees. An analogous result with four marked vertices is proved.

Key words. Dodgson identity, Spanning forest polynomials.

AMS subject classifications. 05C31, 05C50.

1. Introduction. Let G be a connected graph with m vertices, n edges and let the i^{th} edge be assigned a variable α_i . Then we define the graph polynomial of G as

$$\Psi_G = \sum_{T \subseteq G} \prod_{e \notin T} \alpha_e,$$

where the sum runs over the spanning trees T of G . The reason why we pick edges not in the trees is that this form arises naturally in quantum field theory, see for example [3, 5, 10]. We can also obtain this polynomial via the matrix-tree theorem. Let A be the $n \times n$ diagonal matrix with the variables α_i . Orient the edges in the graph and let E be the signed $m \times n$ incidence matrix for this orientation. Let \widehat{E} be the matrix E with any row removed. Define the $m + n - 1$ by $m + n - 1$ block matrix

$$M_G = \begin{bmatrix} A & \widehat{E}^T \\ -\widehat{E} & 0 \end{bmatrix}.$$

Then the matrix-tree theorem states that

$$\Psi_G = \det(M_G).$$

*Received by the editors on January 17, 2012. Accepted for publication on November 22, 2012. Handling Editor: Ravi Bapat.

[†]Department of Mathematics, University of British Columbia, 1984 Mathematics Road, Vancouver BC, V6T 1Z2, Canada. Supported by an NSERC USRA.

[‡]Department of Mathematics, Simon Fraser University, 8888 University Dr, Burnaby BC, V5A 1S6, Canada. Supported by an NSERC discovery grant.

To put this in the usual form of the matrix-tree theorem, note that A is invertible, so we can calculate the determinant using the Schur complement; in this case,

$$\begin{aligned} \det(M) &= \alpha_1 \cdots \alpha_n \det(0 - (-\widehat{E}A^{-1}\widehat{E}^T)) \\ &= \alpha_1 \cdots \alpha_n \det(\widehat{E}A^{-1}\widehat{E}^T), \end{aligned}$$

where $\widehat{E}A^{-1}\widehat{E}^T$ is the graph Laplacian with a row and column removed and with inverted variables. See also Proposition 21 of [4].

There are two important ways to generalize Ψ_G – one via the polynomials and one via the matrix determinant. Let $P = P_1 \cup \cdots \cup P_k$ be a set partition of a subset of the vertices of G . Then define the spanning forest polynomial for G and P as

$$\Phi_G^P = \sum_{F \subseteq G} \prod_{e \notin F} \alpha_e,$$


where the sum runs over spanning forests F of G composed of tree components T_1, \dots, T_k where the vertices P_i are in tree T_i . Alternatively, let I, J, K be sets of indices with $|I| = |J|$. Define the Dodgson polynomial $\Psi_{G,K}^{I,J}$ as

$$\Psi_{G,K}^{I,J} = \det(M_G(I, J))_K,$$

where $M_G(I, J)$ is the submatrix obtained by removing the rows indexed by I and the columns indexed by J from M_G , and the subscript K indicates that we are setting the variables α indexed by K to 0. These two generalizations are related – every Dodgson polynomial can be expressed as a sum of signed spanning forest polynomials (see [7]). Thus, we can use determinant identities to derive identities for spanning forest polynomials. For any square matrix M , we have the classical Dodgson identity

$$\det(M(12, 12)) \det(M) = \det(M(1, 1)) \det(M(2, 2)) - \det(M(1, 2)) \det(M(2, 1))$$

which was popularized by Dodgson through his condensation algorithm [9]. Let G be a graph of the form



with two edges labelled 1 and 2, connecting three vertices v_1, v_2 and v_3 from top to bottom. The Dodgson identity gives the spanning forest polynomial identity (see Section 3)

$$\begin{aligned} \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) &= \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 2 \\ 3 \end{array} = \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 1 \\ 2 \end{array} + \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 1 \\ 2 \end{array} + \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 2 \\ 2 \end{array} + \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 2 \\ 1 \end{array} + \left(\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right) \begin{array}{c} 1 \\ 2 \\ 2 \end{array}, \quad (1.1) \end{aligned}$$

where for example, the graph with labels 1, 1, 2 on the vertices v_1, v_2, v_3 represents Φ_G^P with $P = \{v_1, v_2\} \cup \{v_3\}$.

This result can be interpreted as saying that if we transfer an extra edge from the left hand factor of the left hand side to the right hand factor of the left hand side, thus cutting a spanning tree into two in the left hand factor and joining two of the three trees together in the right hand factor, then we get all pairs of spanning forests with exactly two trees. However, it is subtle to see that the counting matches on both sides, and seems to require chains of edges to be transferred, along the lines of the combinatorial proof of the Dodgson identity due to Zeilberger [12].

Equation (1.1) and its combinatorial interpretation prompted us to investigate spanning forest polynomial identities of the form

$$\left(\begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \end{array} \right) = \left(\begin{array}{c} a_1 \\ a_2 \\ a_3 \\ a_4 \end{array} \right) \left(\begin{array}{c} b_1 \\ b_2 \\ b_3 \\ b_4 \end{array} \right) + \left(\begin{array}{c} c_1 \\ c_2 \\ c_3 \\ c_4 \end{array} \right) \left(\begin{array}{c} d_1 \\ d_2 \\ d_3 \\ d_4 \end{array} \right) + \dots + \left(\begin{array}{c} e_1 \\ e_2 \\ e_3 \\ e_4 \end{array} \right) \left(\begin{array}{c} f_1 \\ f_2 \\ f_3 \\ f_4 \end{array} \right).$$

Our work resulted in such an identity (Theorem 4.14) which is proved in this paper. For this result, we cannot simply interpret a classical determinantal identity; the Jacobi identity on M (see Corollary 4.5) naturally gives a cubic identity for such spanning forest polynomials, while the usual Dodgson identities on submatrices of M can only relate spanning forest polynomials whose degrees differ by at most 2. Rather, we need to combine classical identities in nontrivial ways.

The paper is organized as follows: In Section 2, we will set up our definitions. In Section 3, we will define spanning forest polynomials and give their relation to the minors of M . The main result itself is presented and proved in Section 4. Finally, in Section 5, we conclude with a discussion of the main result, its combinatorial interpretations, and possible extensions.

2. Graph polynomials.

DEFINITION 2.1. Let G be a connected graph and let M_G be a matrix built as in the previous section. Then we define

$$\Psi_G = \det(M_G).$$

By the matrix-tree theorem, Ψ_G is independent of the choice of M_G . We will call Ψ_G the graph polynomial or Kirchhoff polynomial of G . We fix a choice of $M = M_G$ for G .

DEFINITION 2.2. Let I, J , and K be subsets of the edges of G with $|I| = |J|$.

Let $M(I, J)_K$ be the matrix obtained from M by removing the rows indexed by edges of I , the columns indexed by edges of J , and setting $\alpha_i = 0$ for all $i \in K$. Then we define the Dodgson polynomials

$$\Psi_{G,K}^{I,J} = \det M(I, J)_K.$$

When G is clear, it will be suppressed from the notation. Also, if $K = \emptyset$, we may suppress it from the notation.

Up to sign these polynomials are independent of the choice of M (see [4]). By definition it is evident that $\Psi_{G,\emptyset}^{\emptyset,\emptyset} = \Psi_G$. Note that if any element of K appears in I or J then it does not appear in $M(I, J)$, so setting it to zero has no effect.

Contraction and deletion of edges is natural at the level of Dodgson polynomials. Define $\Psi_G = 0$ for G disconnected.

PROPOSITION 2.3. *Let G be a connected graph and let e_i denote the i -th edge in G . Then*

$$\Psi_G^{i,i} = \Psi_{G \setminus e_i} \quad \text{and} \quad \Psi_{G,i} = \Psi_{G/e_i},$$

where $G \setminus e_i$ and G/e_i are the graphs obtained from G by deleting and contracting the edge e_i , respectively.

Proof. The first identity follows immediately from the matrix definition of Ψ and the second from the sum of spanning trees definition. \square

The all-minors matrix-tree theorem [8] tells us that the monomials of any $\Psi_{G,K}^{I,J}$ result from spanning forests of G . For our purposes, it is most useful to organize these spanning forests with the following spanning forest polynomials.

DEFINITION 2.4. Let $P = P_1 \cup P_2 \cup \dots \cup P_k$ be a set partition of a subset of the vertices of G . Then we define

$$\Phi_G^P = \sum_F \prod_{e \notin F} \alpha_e,$$

where the sum runs over spanning forests $F = T_1 \cup T_2 \cup \dots \cup T_k$ with k component trees so that the vertices of P_i are in tree T_i . We note that we are allowing trees consisting of a single vertex.

The relation between Dodgson polynomials and spanning forest polynomials is given by the following proposition which is Proposition 12 in [7].

PROPOSITION 2.5. *Assume $I \cap J = \emptyset$. Then*

$$\Psi_{G,K}^{I,J} = \sum_P \pm \Phi_{G \setminus (I \cup J \cup K)}^P, \tag{2.1}$$

where the sum runs over all set partitions P of the end points of edges of I , J , and K with the property that all the forests of P become trees in both

$$G \setminus I / (J \cup K) \quad \text{and} \quad G \setminus J / (I \cup K).$$

Proof. Take a particular monomial m in $\Psi_{G,K}^{I,J}$. Let F be the set of edges of $G \setminus (I \cup J \cup K)$ which do not contribute to m , and N be the set of edges which do contribute to m . For any set S of edges of G , let $\widehat{E}[S]$ be the submatrix of \widehat{E} consisting of columns indexed by S .

From the form of M , the coefficient of m in $\Psi_K^{I,J}$ is

$$\det \begin{bmatrix} 0 & \widehat{E}[J \cup K \cup F]^T \\ -\widehat{E}[I \cup K \cup F] & 0 \end{bmatrix}.$$

By the matrix-tree theorem in its most stripped down form, see for example [4, Lemma 20], we have that a square matrix formed of columns of \widehat{E} has determinant ± 1 if the edges corresponding to those columns are a spanning tree of G , and has determinant 0 otherwise.

Thus, $J \cup K \cup F$ and $I \cup K \cup F$ are spanning trees of G and so F is a forest of G . Allowing trees consisting of a single vertex only we may view F as a spanning forest of G . Every spanning tree corresponding to the P appearing in the statement will appear in this way.


It remains to check the signs. The coefficient of m in $\Psi_{G,K}^{I,J}$ can be obtained from $\Psi_{G,K}^{I,J}$ by setting the variables in $F \cup K$ to 0 and taking the coefficient of the variables of N , that is the coefficient of m in $\Psi_{G,K}^{I,J}$ is

$$\Psi_{G \setminus N / F, K}^{I,J} = \det(\widehat{E}_{G \setminus N / (F \cup K)}[I]) \det(\widehat{E}_{G \setminus N / (F \cup K)}[J]).$$

But the only information from m left in $G \setminus N / F$ is which end point of edges of I , J , and K lie in the same tree of F . So all terms from the same $\Phi_{G \setminus (I \cup J \cup K)}^P$ must appear with the same sign in $\Psi_{G,K}^{I,J}$. \square

Note that if one forest of a partition P of the end points of I , J , and K becomes a tree in both the graph $G \setminus I / (J \cup K)$ and the graph $G \setminus J / (I \cup K)$, then necessarily all the forest of P must have this property. This is an additional consequence of proof.

3. The classical Dodgson identity. In this section, we interpret the classical Dodgson identity in terms of spanning forest polynomials. Consider the graph G



Apply the Dodgson determinant identity to the matrix M for G

$$\det(M(1, 1)) \det(M(2, 2)) - \det(M(1, 2)) \det(M(2, 1)) = \det(M) \det(M(12, 12)).$$

Interpreting this in terms of Dodgson polynomials gives

$$\Psi_G^{1,1} \Psi_G^{2,2} - \Psi_G^{1,2} \Psi_G^{1,2} = \Psi_G \Psi_G^{12,12},$$

and after setting the variables for edges 1 and 2 to 0, we obtain

$$\Psi_{G,2}^{1,1} \Psi_{G,1}^{2,2} - (\Psi_G^{1,2})^2 = \Psi_G \Psi_G^{12,12}.$$

For a generalization, see Corollary 4.6. Using the deletion-contraction relations we obtain

$$\Psi_{G \setminus e_1/e_2} \Psi_{G \setminus e_2/e_1} - (\Psi_G^{1,2})^2 = \Psi_G \Psi_{G \setminus \{e_1, e_2\}}.$$

Converting to spanning forest polynomials, using Proposition 2.5 for the square term, we find that

$$\begin{aligned} & \left(\Phi_H^{\{a,c\},\{b\}} + \Phi_H^{\{a\},\{b,c\}} \right) \left(\Phi_H^{\{a,b\},\{c\}} + \Phi_H^{\{a,c\},\{b\}} \right) - \left(\pm \Phi_H^{\{a,c\},\{b\}} \right)^2 \\ &= \Phi_H^{\{a,b,c\}} \Phi_H^{\{a\},\{b\},\{c\}}, \end{aligned}$$

where H is the graph with edges 1 and 2 removed, and the three vertices are labelled a, b, c from top to bottom. Rearranging and cancelling the squared term we find that

$$\Phi_H^{\{a,b,c\}} \Phi_H^{\{a\},\{b\},\{c\}} = \Phi_H^{\{a,b\},\{c\}} \Phi_H^{\{a,c\},\{b\}} + \Phi_H^{\{a,b\},\{c\}} \Phi_H^{\{a\},\{b,c\}} + \Phi_H^{\{a,c\},\{b\}} \Phi_H^{\{a\},\{b,c\}}$$

which is just equation (1.1) written in the spanning forest polynomial notation. See Proposition 22 in [7] for more details.

4. The main result. In Section 3, we gave the spanning forest polynomial version of the Dodgson identity. The main result of this paper is an analogous spanning forest polynomial identity for 4 marked vertices. Let us specialize our notation to this situation.

DEFINITION 4.1. Let $v_1, v_2, v_3,$ and v_4 be four distinct vertices of a graph G . We will write (c_1, c_2, c_3, c_4) with $c_i \in \{1, 2, 3, 4, -\}$ to denote the spanning forest

polynomial of the graph G defined by the partition of $\{v_i : c_i \neq -\}$ with one part p_ℓ for each distinct integer ℓ in (c_1, c_2, c_3, c_4) defined by $p_\ell = \{v_i : c_i = \ell\}$, and no other parts.

DEFINITION 4.2. Using the previous definition, we further abbreviate by defining the following special cases

$$\begin{aligned} A_1 &= (1, 1, 2, 3), & A_2 &= (1, 2, 1, 3), & A_3 &= (1, 2, 2, 3), \\ A_4 &= (1, 2, 3, 1), & A_5 &= (1, 2, 3, 2), & A_6 &= (1, 2, 3, 3) \end{aligned}$$

of 3 parts each, the following cases


$$\begin{aligned} B_1 &= (1, 1, 1, 2), & B_2 &= (1, 1, 2, 1), & B_3 &= (1, 2, 1, 1), \\ B_4 &= (1, 2, 2, 2), & B_5 &= (1, 1, 2, 2), & B_6 &= (1, 2, 1, 2), \\ & & B_7 &= (1, 2, 2, 1) \end{aligned}$$

of 2 parts each, and finally,

$$P = (1, 1, 1, 1).$$

The A_i and B_i are the different distinct ways in which we can partition four vertices in 3 and 2 sets, respectively. P is just Ψ_G for this G with four marked vertices.

EXAMPLE 4.3. Let G be a connected graph



with marked vertices v_1, v_2, v_3 , and v_4 . Then,

$$(1, 2, -, 1) = \Phi_G^{\{v_1, v_4\}, \{v_2\}}.$$

We are now ready to state our main result; see Theorem 4.14.

Let G be a connected graph with four marked vertices. Then the main result is

$$\begin{aligned}
 (1, 1, 1, 1)(1, 2, 3, 4) = & (1 - x_1 - x_2)A_4B_1 + x_7A_2B_4 + (1 - x_3 - x_2)A_5B_1 \\
 & + (1 - x_1 - x_4)A_6B_1 + x_2A_2B_2 + (x_3 + x_2 - x_5)A_3B_2 \\
 & + (1 - x_1 - x_6)A_6B_2 + x_1A_1B_3 + (x_1 - x_7 + x_4)A_3B_3 \\
 & + (x_1 - x_8 + x_6)A_5B_3 + x_5A_1B_4 + (x_1 - x_5 + x_4)A_3B_5 \\
 & + (x_1 - x_5 + x_6)A_5B_5 + x_3A_1B_6 + (x_3 + x_2 - x_7)A_3B_6 \\
 & + (1 - x_1 - x_2 + x_8 - x_6)A_4B_6 + (x_2 + x_7 - x_4)A_2B_7 \\
 & + (1 - x_1 - x_7 + x_8 - x_6)A_6B_6 + (x_1 + x_5 - x_3)A_1B_7 \\
 & + (1 + x_5 - x_3 - x_2 - x_8)A_5B_7 \\
 & + (1 - x_1 + x_7 - x_4 - x_8)A_6B_7 \\
 & + x_8A_4B_4 + x_4A_2B_5 + x_6A_4B_5
 \end{aligned} \tag{4.1}$$

for any x_1, \dots, x_8 .

This is the generalization of the classical Dodgson identity phrased in terms of spanning forest polynomials. It is possible to give a graphical representation of this identity like in equation (1.1) but it would take too much space.

To guide the reader through the lemmas and calculations which follow, here we will describe the outline of the proof of (4.1).

Let $E(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)$ be the right hand side of (4.1). We will first check that E does not depend on the values of the x_i by checking that the coefficient of each x_i in E is zero (Proposition 4.10). Then we will be free to use any choice of x_i which is algebraically convenient.

Next, using the classical Jacobi identity on an auxiliary graph with three extra edges we will obtain an expression for

$$(1, 2, 3, 4)^2(1, 1, 1, 1)$$

which is a linear combination of products of the form A_iA_j (Lemma 4.11).

Then we will calculate each PA_i as a linear combination of products of the form A_jB_k (Lemma 4.12). Using this calculation we will obtain an expression for

$$((1, 2, 3, 4)(1, 1, 1, 1))^2,$$

which we can then check is the same as

$$E(0, 0, 0, 0, 0, 0, 0, 0)E(0, 1, 0, 1, 1, 1, 1, 1)$$

(Lemma 4.13). The proof of (4.1) then will conclude by checking the sign.

Let us pause here for a brief word on the role of the x_i . The Dodgson identities give a number of quadratic identities between the A_i and B_i . Consequently, there cannot be a unique way to write $(1, 1, 1, 1)(1, 2, 3, 4)$ as a linear combination of products $A_i B_k$. The x_i 's describe this nonuniqueness. We can specialize to get more manageable equations, for example setting all $x_i = 0$ and collecting terms gives

$$(1, 2, 3, 4)(1, 1, 1, 1) = (1, 2, 3, 1)(1, -, 1, 2) + (1, 2, 3, 2)(-, 1, 1, 2) + (1, 2, 3, 3)(-, -, 1, 2)$$

but no such specialization is canonical, so we gave the general equation in (4.1).

Now we can proceed with the lemmas. We will need a particular form of the Jacobi determinantal identity and some further Dodgson identities which follow from it.

Let M be an $n \times n$ matrix. Let I and J be subsets of $\{1, 2, \dots, n\}$. Let $M(I, J)$ be the matrix obtained from M by removing rows I and columns J . Similarly, let $M[I, J]$ be the matrix where we only keep rows I and columns J . Finally, we let

$$s(I, J) = \sum_{x \in I} x + \sum_{x \in J} x.$$

THEOREM 4.4. *Let M be a nonsingular $n \times n$ matrix and let I and J be two sets in $\{1, 2, \dots, n\}$ with $|I| = |J| = t$. Let $A = \text{adj } M$ and define the matrix B by $b_{ij} = \det(M(i, j))$. Then*

$$\det(B[I, J]) = (\det M)^{t-1} \det(M(I, J)).$$

Proof. To remain self contained we will give a proof following the idea of the proof of Lemma 28 of [4]. Let I_n be the $n \times n$ identity matrix. Then

$$AM = I_n(\det M).$$

Take determinants to get

$$\det(A) = (\det M)^{n-1}.$$

Now if the k -th element of I is i_k and the k -th element of J is j_k let C be M with the j_k column replaced by \mathbf{e}_{i_k} , where \mathbf{e}_i is the i -th standard basis element of \mathbb{R}^n . Then multiplying out column by column we get that AC is the matrix D whose j -th column is

$$\begin{cases} (\det M)\mathbf{e}_j & \text{if } j \text{ is not in } J \\ A\mathbf{e}_{i_k} & \text{if } j \text{ is } j_k \text{ in } J \end{cases}.$$

Now notice that

$$\det C = (-1)^{s(I,J)} \det(M(I, J))$$

and

$$\begin{aligned} \det D &= (\det M)^{n-t} \det(A[J, I]) \\ &= (\det M)^{n-t} \det(B[I, J])(-1)^{s(I,J)}. \end{aligned}$$

The second equality holds since $A[J, I]$ can be converted to $B[I, J]$ by multiplying each row and each column which had an odd index in M by -1 and then taking a transpose; on determinants this changes the sign $s(I, J)$ times.

Finally, taking the determinant of $AC = D$, using the above calculations and dividing by $(\det M)^{n-t}$ gives us the result. \square

This formula can readily be translated into the Dodgson polynomials language.

COROLLARY 4.5. *Let G be a graph and M be its associated matrix. Let I, J and E be subsets of the edges, such that $|I| = |J| = k$. Then the k -level Dodgson identity is*

$$\det \left(\Psi_{G,E}^{I_i, J_j} \right)_{1 \leq i, j \leq k} = \Psi_{G,E}^{I,J} (\Psi_{G,E})^{k-1},$$

where $I = \{I_1, \dots, I_k\}$ and $J = \{J_1, \dots, J_k\}$.

Proof. Use Theorem 4.4. By definition $\det M = \Psi_G$ and $\det M(I, J) = \Psi_G^{I,J}$. Now $B[I, J]_{ij} = \det(M(I_i, J_j)) = \Psi^{I_i, J_j}$. Finally, we set $\alpha_e = 0$ for $e \in E$. \square

Careful book-keeping and application of the above identity yield the following corollary.

COROLLARY 4.6. *Let M be an associated matrix for the graph G . Let E, I, J, A and B be ordered sets indexing edges in G , such that $|A \cap I| = |B \cap J| = 0$, $|I| = |J| = k$ and $|A| = |B| = l$. Then the modified k -level Dodgson identity is*

$$\det \left(\Psi_{G,E}^{A \cup I_i, B \cup J_j} \right)_{1 \leq i, j \leq k} = \Psi_{G,E}^{A \cup I, B \cup J} \left(\Psi_{G,E}^{A,B} \right)^{k-1} \quad (4.2)$$

where $I = \{I_1, \dots, I_k\}$ and $J = \{J_1, \dots, J_k\}$.

Note that when $k = 2$ this gives the classical Dodgson identity.

We will use the following rearrangement of the $k = 2$ case.

PROPOSITION 4.7 (Brown, [4]). *Let I and J be subsets of edges of G with $|J| = |I| + 1$. Let a, b, x be edge indices with $a \notin I$, $b, x \notin I \cup J$, and $x < a < b$. Let*

$S = I \cup J \cup \{a, b, x\}$. Then

$$\Psi_S^{Ia,J} \Psi_S^{Ibx,Jx} - \Psi_S^{Iax,Jx} \Psi_S^{Ib,J} = \Psi_S^{Ix,J} \Psi_S^{Iab,Jx}. \quad (4.3)$$

Proof. This is equation (23) from [4]; the proof proceeds by applying the $k = 2$ case of (4.2) three times and rearranging. \square

We only need the signs relating Dodgson polynomials to spanning forest polynomials in two cases, given in the next lemma. The general formula is found in Proposition 16 of [7], but we give here a self contained proof of the cases we need.

LEMMA 4.8. *Fix an order and orientation of the edges of a graph G . Suppose edges 1, 2, and 3 have a common vertex v . Let w_1, w_2 , and w_3 be distinct and be the other end points of 1, 2, and 3, and let*

$$\epsilon(i, j) = \begin{cases} 1 & \text{if } i \text{ and } j \text{ are both oriented into } v \text{ or both oriented out of } v \\ -1 & \text{otherwise} \end{cases}$$

for $i \neq j \in \{1, 2, 3\}$. Then

$$\Psi^{1,2} = \epsilon(1, 2) \Phi^{\{v\}, \{w_1, w_2\}}$$

and

$$\Psi_k^{i,j} = \epsilon(i, j) (-1)^{i-j+1} \Phi^{\{v\}, \{w_i, w_j\}, \{w_k\}},$$

where $\{i, j, k\} = \{1, 2, 3\}$ in some order.

Proof. The first statement of the lemma follows from the second with $k = 3$ applied to the graph G with a new vertex w_3 added and a new edge 3 from v to w_3 . Consider the second statement. Let x be the vertex which was removed when forming M . We choose it to be disjoint from $\{v, w_i, w_j\}$.

Note that $\{v\}, \{w_i, w_j\}, \{w_k\}$ is the only set partition compatible with $\Psi_k^{i,j}$. From the observations preceding this lemma, if $\Psi_k^{i,j} = 0$, then there are no common spanning trees of $G \setminus i / \{j, k\}$ and $G \setminus j / \{i, k\}$ and so in particular there are no terms in $\Phi^{\{v\}, \{w_i, w_j\}, \{w_k\}}$. Thus,

$$\Psi_k^{i,j} = 0 \quad \Leftrightarrow \quad \Phi^{\{v\}, \{w_i, w_j\}, \{w_k\}} = 0.$$

By (2.1) we know that $\Psi_k^{i,j} = f \Phi^{\{v\}, \{w_i, w_j\}, \{w_k\}}$ for some $f \in \{-1, 1\}$, so it suffices to consider one term of $\Psi^{i,j}$. Pick a term t where the tree out of w_i and w_j

intersects x . Let F be the forest corresponding to t . The sign of t in $\Psi^{i,j}$ is $\det N$ where

$$N = \begin{bmatrix} 0 & \widehat{E}[\{i, k\} \cup F]^T \\ -\widehat{E}[\{j, k\} \cup F] & 0 \end{bmatrix}.$$

Let $B = \widehat{E}[\{k\} \cup F]$. Then $\widehat{E}[\{i, k\} \cup F]$ and $\widehat{E}[\{j, k\} \cup F]$ are formed by inserting the i -th and j -th columns respectively of \widehat{E} into B . If $\{i, j\} = \{1, 2\}$ the insertions are both made in the first column. Let i' be the index of the inserted column i and j' the index of the inserted column in j . Thus, if $\{i, j\} = \{1, 2\}$ then $i' = j' = 1$; if $\{i, j\} = \{1, 3\}$, then $\{i', j'\} = \{1, 2\}$; and if $\{i, j\} = \{2, 3\}$ then $i' = j' = 2$.

Consider B with the row corresponding to v removed. This is the same as the columns corresponding to edges of $\{k\} \cup F$ in the incidence matrix of the graph with v and x identified. This has determinant ± 1 since $\{k\} \cup F$ was chosen to be a tree in this graph. Likewise, removing the row corresponding to w_1 or w_2 we get a zero determinant since $\{k\} \cup F$ is not a tree in the graph with w_1 or w_2 identified with x .

Thus, if we expand $\det \widehat{E}[\{i, k\} \cup F]$ down the inserted column, only the cofactor coming from row v is retained, and likewise for $\widehat{E}[\{j, k\} \cup F]$. Thus,

$$\begin{aligned} \det N &= \det(\widehat{E}[\{i, k\} \cup F]) \det(\widehat{E}[\{j, k\} \cup F]) \\ &= e_{i,\ell} e_{j,\ell} (-1)^{i'+j'+2\ell} \det(\widehat{B})^2 \\ &= \epsilon(i, j) (-1)^{i-j+1}, \end{aligned}$$

where ℓ is the index of row v , \widehat{B} is B with row v removed and $e_{r,s}$ is the (r, s) entry of \widehat{E} . \square

Here is a catalogue of the instances of the Dodgson identity which we will need in the main argument, written in terms of the A_i and B_i from Definition 4.2.

LEMMA 4.9.

$$A_1(B_3 + B_7) + A_2(B_7 - B_5) - A_4(B_1 + B_5) = 0 \tag{4.4}$$

$$A_1(B_4 + B_7) + A_5(B_7 - B_5) - A_3(B_2 + B_5) = 0 \tag{4.5}$$

$$A_2(B_2 + B_7) + A_1(B_7 - B_6) - A_4(B_1 + B_6) = 0 \tag{4.6}$$

$$A_2(B_4 + B_7) + A_6(B_7 - B_6) - A_3(B_3 + B_6) = 0 \tag{4.7}$$

$$A_3(B_2 + B_6) + A_1(B_6 - B_7) - A_5(B_1 + B_7) = 0 \tag{4.8}$$


$$A_3(B_3 + B_5) + A_2(B_5 - B_7) - A_6(B_1 + B_7) = 0 \tag{4.9}$$

$$A_4(B_4 + B_5) + A_5(B_5 - B_7) - A_6(B_2 + B_7) = 0 \tag{4.10}$$

$$A_4(B_4 + B_6) + A_6(B_6 - B_7) - A_5(B_3 + B_7) = 0 \tag{4.11}$$

$$A_5(B_3 + B_5) + A_4(B_5 - B_6) - A_6(B_2 + B_6) = 0. \tag{4.12}$$

Proof. The equations differ only by permuting the four marked vertices, so it suffices to prove (4.5). Consider the graph



We use identity (4.3) with $x = 1$, $a = 2$, $b = 3$, $I = \emptyset$ and $J = \{2\}$, and by Lemma 4.8 we obtain

$$(1, -, 2, 3)(1, 2, 2, -) - (1, -, 2, -)(1, 2, 2, 3) = (1, 2, 3, 2)(1, -, 2, 2).$$

For the sign of $(1, -, 2, 2)$, note that the cutting happens first so that edges 1 and 3 become adjacent columns in the cut matrix. Expanding, $(1, -, 2, 3) = A_1 + A_3 + A_5$, $(1, 2, 2, -) = B_4 + B_7$, $(1, -, 2, -) = B_2 + B_4 + B_5 + B_7$, and $(1, -, 2, 2) = B_4 + B_5$. We substitute these in and rearranging gives us equation (4.5). \square

PROPOSITION 4.10. *All the free variables in (4.1) are explained by Dodgson identities.*

Proof. The coefficient of x_3 in equation (4.1) is the right hand side of equation (4.8), and thus is 0. Similarly, the coefficients of x_4 , x_5 , x_6 , x_7 , and x_8 are zero by (4.9), (4.5), (4.12), (4.7), and (4.11), respectively. The coefficient of x_2 is in a different form, but is also zero as it is the sum of the right hand sides of (4.6) and (4.8). Finally, the coefficient of x_1 is the sum of the right hand sides of (4.12), (4.9), and (4.4) and so is zero. \square

LEMMA 4.11.

$$(1, 1, 1, 1)(1, 2, 3, 4)^2 = \det \begin{pmatrix} A_1 + A_3 + A_5 & -A_3 & -A_5 \\ -A_3 & A_2 + A_3 + A_6 & -A_6 \\ -A_5 & -A_6 & A_4 + A_5 + A_6 \end{pmatrix}.$$


Proof. Let H be G with three new edges 1, 2 and 3 connecting vertex v_1 with the other 3 marked vertices. By Corollary 4.5 with $k = 3$ and $I = J = E = \{1, 2, 3\}$ we

have

$$(\Psi_{H,123})^2 \Psi_H^{123,123} = \det \begin{pmatrix} \Psi_{H,23}^{1,1} & \Psi_{H,3}^{1,2} & \Psi_{H,2}^{1,3} \\ \Psi_{H,3}^{1,2} & \Psi_{H,13}^{2,2} & \Psi_{H,1}^{2,3} \\ \Psi_{H,2}^{1,3} & \Psi_{H,1}^{2,3} & \Psi_{H,12}^{3,3} \end{pmatrix}, \quad (4.13)$$


where $\Psi_H^{123,123}$ is the graph polynomial of G with the edges 1,2 and 3 removed, namely $\Psi_H^{123,123} = P = (1, 1, 1, 1)$; $\Psi_{H,123}$ is the spanning forest polynomial of G where each of the four vertices is in a separate tree, namely $\Psi_{H,123} = (1, 2, 3, 4)$.

The Dodgson polynomials on the main diagonal are just spanning forest polynomials of G where one of the edges is removed and the other two contracted. By inspection, these are precisely the terms in the diagonal of the matrix in the result. The Dodgson polynomials on the off-diagonals require more care. We orient the edges like this: edge 2 goes towards vertex 1 and the other two away from it.



This ensures all the off-diagonal signs are negative (by Lemma 4.8) and that each Dodgson polynomial gives the desired spanning forest polynomial. The result follows. \square

Note that the matrix in Lemma 4.11 is the Laplacian matrix with row and column 1 removed for the following graph




where the edge labels are the A 's. This is not a coincidence and there is a general identity which we leave out for brevity. However, the statement is analogous.

To complete the calculation we need to multiply the whole expression by P and use the following

LEMMA 4.12.

$$\begin{aligned} PA_1 &= B_1B_2 + B_1B_5 + B_2B_5 + B_5B_6 + B_5B_7 - B_6B_7 \\ PA_2 &= B_1B_3 + B_1B_6 + B_3B_6 + B_5B_6 - B_5B_7 + B_6B_7 \\ PA_3 &= B_1B_4 + B_1B_7 + B_4B_7 - B_5B_6 + B_5B_7 + B_6B_7 \\ PA_4 &= B_2B_3 + B_2B_7 + B_3B_7 - B_5B_6 + B_5B_7 + B_6B_7 \\ PA_5 &= B_2B_4 + B_2B_6 + B_4B_6 + B_5B_6 - B_5B_7 + B_6B_7 \\ PA_6 &= B_3B_4 + B_3B_5 + B_4B_5 + B_5B_6 + B_5B_7 - B_6B_7. \end{aligned}$$

Proof. By symmetry of the four vertices it suffices to prove the formula for PA_1 . Consider the graph



Then

$$\begin{aligned} PA_1 &= -\Psi^{123,123}\Psi_2^{1,3} \quad \text{by Lemma 4.8} \\ &= \Psi^{12,32}\Psi_2^{13,13} - \Psi^{12,31}\Psi^{13,23} \quad \text{by (4.2) with } A = \{1\}, B = \{3\}, I = \{2, 3\}, \\ &\quad J = \{1, 2\}, \text{ and } E = \{1, 2, 3\} \\ &= (1, 1, 2, -)(-, -, 1, 2) - (1, -, 2, 1)(-, 1, 2, 1) \quad \text{by Lemma 4.8} \\ &= (B_2 + B_5)(B_1 + B_2 + B_6 + B_7) - (B_2 + B_7)(B_2 + B_6) \\ &= B_1B_2 + B_1B_5 + B_2B_5 + B_5B_6 + B_5B_7 - B_6B_7. \quad \square \end{aligned}$$

Now we find out what happens when we multiply the equation in Lemma 4.11 by P .

LEMMA 4.13.

$$((1, 1, 1, 1)(1, 2, 3, 4))^2 = E(0, 0, 0, 0, 0, 0, 0, 0)E(0, 1, 0, 1, 1, 1, 1, 1),$$

where $E(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)$ is the right hand side of (4.1).

Proof. By definition

$$E(0, 0, 0, 0, 0, 0, 0, 0) = (A_5 + A_6)(B_1 + B_7) + A_6(B_2 + B_6) + A_4(B_1 + B_6) \quad (4.14)$$

and

$$E(0, 1, 0, 1, 1, 1, 1, 1) = (A_1 + A_2)(B_4 + B_7) + A_2(B_2 + B_5) + A_4(B_4 + B_5). \quad (4.15)$$

Use Lemma 4.11 and 4.12 to calculate $((1, 1, 1, 1)(1, 2, 3, 4))^2$. With some trial and error we chose which lines of Lemma 4.12 to use so that the final result would look as much as possible like the product of (4.14) and (4.15). The term $((1, 1, 1, 1)(1, 2, 3, 4))^2$ equals

$$\begin{aligned} & (A_1 + A_2)(PA_3)(A_5 + A_6) + A_2(PA_1)(A_5 + A_6) + (A_1 + A_2)(PA_5)A_6 \\ & \quad + A_4(A_1 + A_2 + A_5 + A_6)(PA_3) + A_4(A_2 + A_6)(PA_1 + PA_5) \\ = & (A_1 + A_2)(A_5 + A_6)(B_1B_7 + B_4B_7 + B_1B_4) \\ & \quad + A_2(A_5 + A_6)(B_1B_2 + B_2B_5 + B_1B_5 + B_5B_7) \\ & \quad + (A_1 + A_2)A_6(B_2B_6 + B_4B_6 + B_2B_4 + B_6B_7) \\ & \quad + A_4(A_1 + A_2 + A_5 + A_6)(B_1B_7 + B_4B_7 + B_1B_4 + B_5B_7 + B_6B_7) \\ & \quad + A_4(A_2 + A_6)(B_1B_5 + B_2B_5 + B_1B_2 + B_2B_6 + B_4B_6 + B_2B_4 + B_5B_6) \\ & \quad + B_5B_6A_2A_6 - B_5B_6A_1A_5 - B_5B_6A_4(A_1 + A_5) \\ & \quad + B_5B_7(A_1 + A_2)A_5 + B_6B_7A_1(A_5 + A_6). \end{aligned}$$

Now we consider the difference between this expression and (4.14) times (4.15)

$$\begin{aligned} & A_1A_5(-B_7^2 - B_5B_6 + B_5B_7 + B_6B_7) + A_1A_6(-B_7^2 + B_2B_6 - B_2B_7 + B_6B_7) \\ & \quad + A_2A_5(-B_7^2 + B_2B_5 - B_2B_7 + B_5B_7) + A_2A_6(-B_7^2 - 2B_2B_7 - B_2^2) \\ & \quad + A_4A_5(B_1B_7 + B_6B_7 - B_1B_5 - B_5B_6) \\ & \quad + A_4A_6(B_1B_7 + B_6B_7 + B_1B_2 + B_2B_6) \\ & \quad + A_1A_4(B_4B_7 + B_5B_7 - B_6B_4 - B_5B_6) \\ & \quad + A_2A_4(B_4B_7 + B_5B_7 + B_2B_5 + B_2B_4) - A_4^2(B_1 + B_6)(B_4 + B_5) \end{aligned}$$

$$\begin{aligned}
 &= -(A_6(B_2 + B_7) - A_5(B_5 - B_7))(A_2(B_2 + B_7) - A_1(B_6 - B_7)) \\
 &\quad - A_4A_5(B_1 + B_6)(B_5 - B_7) + A_4A_6(B_1 + B_6)(B_2 + B_7) \\
 &\quad - A_1A_4(B_4 + B_5)(B_6 - B_7) + A_2A_4(B_2 + B_7)(B_5 + B_4) \\
 &\quad - A_4^2(B_1 + B_6)(B_4 + B_5) \\
 &= -A_4(B_4 + B_5)(A_2(B_2 + B_7) - A_1(B_6 - B_7)) \quad \text{by (4.10)} \\
 &\quad - A_4A_5(B_1 + B_6)(B_5 - B_7) + A_4A_6(B_1 + B_6)(B_2 + B_7) \\
 &\quad - A_1A_4(B_4 + B_5)(B_6 - B_7) + A_2A_4(B_2 + B_7)(B_5 + B_4) \\
 &\quad - A_4^2(B_1 + B_6)(B_4 + B_5) \\
 &= A_4(B_1 + B_6)(A_6(B_2 + B_7) + A_5(B_7 - B_5) - A_4(B_4 + B_5)) \\
 &= 0 \quad \text{by (4.10)}. \quad \square
 \end{aligned}$$

We now have everything needed for our main theorem

THEOREM 4.14. *Let G be a connected graph with four marked vertices. Then (4.1) holds for all x_1, \dots, x_8 , that is*

$$\begin{aligned}
 (1, 1, 1, 1)(1, 2, 3, 4) = & (1 - x_1 - x_2)A_4B_1 + x_7A_2B_4 + (1 - x_3 - x_2)A_5B_1 \\
 & + (1 - x_1 - x_4)A_6B_1 + x_2A_2B_2 + (x_3 + x_2 - x_5)A_3B_2 \\
 & + (1 - x_1 - x_6)A_6B_2 + x_1A_1B_3 + (x_1 - x_7 + x_4)A_3B_3 \\
 & + (x_1 - x_8 + x_6)A_5B_3 + x_5A_1B_4 + (x_1 - x_5 + x_4)A_3B_5 \\
 & + (x_1 - x_5 + x_6)A_5B_5 + x_3A_1B_6 + (x_3 + x_2 - x_7)A_3B_6 \\
 & + (1 - x_1 - x_2 + x_8 - x_6)A_4B_6 + (x_2 + x_7 - x_4)A_2B_7 \\
 & + (1 - x_1 - x_7 + x_8 - x_6)A_6B_6 + (x_1 + x_5 - x_3)A_1B_7 \\
 & + (1 + x_5 - x_3 - x_2 - x_8)A_5B_7 \\
 & + (1 - x_1 + x_7 - x_4 - x_8)A_6B_7 \\
 & + x_8A_4B_4 + x_4A_2B_5 + x_6A_4B_5
 \end{aligned}$$

for all x_1, \dots, x_8 .

Proof. Let $E(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8)$ be the right hand side of (4.1).

By Lemma 4.13 we know that

$$E(0, 0, 0, 0, 0, 0, 0, 0)E(0, 1, 0, 1, 1, 1, 1, 1) = ((1, 2, 3, 4)(1, 1, 1, 1))^2$$

and by Proposition 4.10 we know that E does not depend on the x_i . Thus, we have

$$E(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8) = \pm(1, 2, 3, 4)(1, 1, 1, 1).$$

It remains to check the sign. Note that $(1, 1, 1, 1) = P = \Psi_G$ and $(1, 2, 3, 4)$ is Ψ for G with v_1, v_2, v_3 , and v_4 identified. Since both $(1, 1, 1, 1)$ and $(1, 2, 3, 4)$ are Kirchhoff

polynomials of graphs all monomials appear with nonnegative coefficients. Looking at (4.14), we see that the sign is 1 and the proof is complete. \square

5. Conclusions. Theorem 4.14 gives a nice generalization of (1.1). Equation (1.1) itself is crucial to the combinatorial and algebro-geometric approach to understanding the periods of Feynman integrals [11, 2, 7, 6, 1]. In such work, having a good intuition of how to massage the polynomials which occur is crucial, and it is the second author's experience that spanning forest polynomials and their identities are very useful in this regard.

We can ask for an edge-transferring interpretation of Theorem 4.14, comparable to what we discussed for (1.1) in the introduction. Consider (4.14), which is the result of setting the free variables to 0 in our main theorem. Collecting terms this gives

$$(1, 2, 3, 4)(1, 1, 1, 1) = (1, 2, 3, 1)(1, -, 1, 2) + (1, 2, 3, 2)(-, 1, 1, 2) + (1, 2, 3, 3)(-, -, 1, 2)$$

which says that we can choose to transfer an edge from any spanning forests contributing to $(1, 1, 1, 1)$ to one of those contributing to $(1, 2, 3, 4)$, so that we always merge the tree of the last vertex from $(1, 2, 3, 4)$ into one of the other trees, and always split the last and second last vertices of $(1, 1, 1, 1)$ into separate trees. Furthermore, the identity describes precisely how the split trees will interact with the remaining vertices. We know of no direct combinatorial proof which follows this interpretation.

We initially obtained (4.1) by a numerical calculation. We first picked a graph on which to perform the calculations – we picked K_4 , K_5 and K_6 . Then we calculated each A_i and B_i on this graph and then formed all possible products of A 's and B 's and formed the sum $\sum_{s,t} x_{st} A_s B_t$, where x_{st} is a constant, $1 \leq s \leq 6$ and $1 \leq t \leq 7$ for a total of 42 constants, and solved the linear system. The initial numerical calculation could, a priori, have had spurious degrees of freedom, but it could not miss any true identity of the desired form. Consequently, (4.1) is the most general quadratic formula involving 4 marked vertices.

A natural question is what do formulae for more marked vertices look like. Numerical calculations show that for 5 and 6 marked vertices the formulae have 15 and 24 free variables. For the classical Dodgson identity, the A 's and B 's are the same. If we treat the A 's and the B 's as different, we have a formula with 3 free variables. Trivially, a formula for 2 marked vertices has no free variables. For $n = 2, 3, 4, 5$ and 6 the identities so far point to expressions having 0, 3, 8, 15 and 24 variables in formulae for n marked vertices. These numbers are generated by $n(n-2)$ for $n = 2, 3, 4, 5$ and 6.

REFERENCES

- [1] P. Aluffi and M. Marcolli. Parametric Feynman integrals and determinant hypersurfaces. *Adv. Theor. Math. Phys.*, 14(3):911–964, 2010. Preprint arXiv:0901.2107.
- [2] S. Bloch, H. Esnault, and D. Kreimer. On motives associated to graph polynomials. *Commun. Math. Phys.*, 267:181–225, 2006. Preprint arXiv:math/0510011v1 [math.AG].
- [3] C. Bogner and S. Weinzierl. Feynman graph polynomials. Preprint arXiv:1002.3458.
- [4] F. Brown. On the periods of some Feynman integrals. Preprint arXiv:0910.0114.
- [5] F. Brown. The massless higher-loop two-point function. *Commun. Math. Phys.*, 287:925–958, 2009. Preprint arXiv:0804.1660.
- [6] F. Brown and O. Schnetz. A K3 in ϕ^4 . *Duke Math J.*, 161(10):1817–1862, 2012. Preprint arXiv:1006.4064.
- [7] F. Brown and K. Yeats. Spanning forest polynomials and the transcendental weight of Feynman graphs. *Commun. Math. Phys.*, 301(2):357–382, 2011. Preprint arXiv:0910.5429.
- [8] S. Chaiken. A combinatorial proof of the all minors matrix tree theorem. *SIAM J. Alg. Disc. Meth.*, 3(3):319–329, 1982.
- [9] C.L. Dodgson. Condensation of determinants, being a new and brief method for computing their arithmetic values. *Proc. Roy. Soc. Ser. A*, 15:150–155, 1866.
- [10] D. Kreimer. The core Hopf algebra. In *Quanta of Maths*, Clay Mathematics Proceedings, 11:313–321, 2010. arXiv:0902.1223.
- [11] O. Schnetz. Quantum periods: A census of ϕ^4 -transcendentals. *Communications in Number Theory and Physics*, 4(1):1–48, 2010. Preprint arXiv:0801.2856.
- [12] D. Zeilberger. Dodgson’s determinant-evaluation rule proved by two-timing men and women. *Electron. J. Combin.*, 4(2), 1997.