

1-1-2006

# Sign Patterns That Allow a Positive or Nonnegative Left Inverse

I. J. Kim

D. D. Olesky

Bryan L. Shader

University of Wyoming, [bshader@uwyo.edu](mailto:bshader@uwyo.edu)

P. Van den Driessche

Follow this and additional works at: [http://repository.uwyo.edu/math\\_facpub](http://repository.uwyo.edu/math_facpub)



Part of the [Mathematics Commons](#)

---

## Publication Information

Kim, I. J.; Olesky, D. D.; Shader, Bryan L.; and Van den Driessche, P. (2006). "Sign Patterns That Allow a Positive or Nonnegative Left Inverse." *SIAM Journal on Matrix Analysis and Applications* 29.2, 554-565.

This Article is brought to you for free and open access by the Mathematics at Wyoming Scholars Repository. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Wyoming Scholars Repository. For more information, please contact [scholcom@uwyo.edu](mailto:scholcom@uwyo.edu).

## SIGN PATTERNS THAT ALLOW A POSITIVE OR NONNEGATIVE LEFT INVERSE\*

IN-JAE KIM<sup>†</sup>, D. D. OLESKY<sup>‡</sup>, B. L. SHADER<sup>§</sup>, AND P. VAN DEN DRIESSCHE<sup>¶</sup>

**Abstract.** An  $m$  by  $n$  sign pattern  $\mathcal{S}$  is an  $m$  by  $n$  matrix with entries in  $\{+, -, 0\}$ . Such a sign pattern allows a positive (resp., nonnegative) left inverse, provided that there exist an  $m$  by  $n$  matrix  $A$  with the sign pattern  $\mathcal{S}$  and an  $n$  by  $m$  matrix  $B$  with only positive (resp., nonnegative) entries satisfying  $BA = I_n$ , where  $I_n$  is the  $n$  by  $n$  identity matrix. For  $m > n \geq 2$ , a characterization of  $m$  by  $n$  sign patterns with no rows of zeros that allow a positive left inverse is given. This leads to a characterization of all  $m$  by  $n$  sign patterns with  $m \geq n \geq 2$  that allow a positive left inverse, giving a generalization of the known result for the square case, which involves a related bipartite digraph. For  $m \geq n$ ,  $m$  by  $n$  sign patterns with all entries in  $\{+, 0\}$  and  $m$  by 2 sign patterns with  $m \geq 2$  that allow a nonnegative left inverse are characterized, and some necessary or sufficient conditions for a general  $m$  by  $n$  sign pattern to allow a nonnegative left inverse are presented.

**Key words.** bipartite digraph, nonnegative left inverse, positive left inverse, positive left null-vector, sign pattern, strong Hall

**AMS subject classifications.** 15A09, 15A48, 05C20, 05C50

**DOI.** 10.1137/060660916

**1. Introduction.** An  $m$  by  $n$  sign pattern  $\mathcal{S} = [s_{ij}]$  is an  $m$  by  $n$  matrix with entries in  $\{+, -, 0\}$ . If a sign pattern  $\mathcal{S}$  has all entries in  $\{+, 0\}$ , then  $\mathcal{S}$  is a *nonnegative* sign pattern. A *subpattern* of  $\mathcal{S}$  is an  $m$  by  $n$  sign pattern  $\mathcal{U} = [u_{ij}]$  such that  $u_{ij} = 0$  whenever  $s_{ij} = 0$ . If  $\mathcal{U}$  is a subpattern of  $\mathcal{S}$ , then  $\mathcal{S}$  is a *superpattern* of  $\mathcal{U}$ . The *sign pattern class*  $Q(\mathcal{S})$  of an  $m$  by  $n$  sign pattern  $\mathcal{S}$  is the set of  $m$  by  $n$  matrices  $A = [a_{ij}]$  such that  $\text{sgn}(a_{ij}) = s_{ij}$  for all  $i, j$ . If  $A \in Q(\mathcal{S})$ , then  $A$  is a *realization* of  $\mathcal{S}$ .

Let  $A = [a_{ij}]$  be an  $m$  by  $n$  matrix. If each entry of  $A$  is positive (resp., nonnegative), then  $A$  is *positive* (resp., *nonnegative*), written  $A > 0$  (resp.,  $A \geq 0$ ). A *left inverse* of an  $m$  by  $n$  matrix  $A$  is an  $n$  by  $m$  matrix  $B$  such that  $BA = I_n$ , where  $I_n$  denotes the  $n$  by  $n$  identity matrix. If  $B > 0$ , then  $B$  is a *positive* left inverse (abbreviated as *PLI*) of  $A$ . If  $B \geq 0$ , then  $B$  is a *nonnegative* left inverse (abbreviated as *NLI*) of  $A$ . In general, neither a PLI nor an NLI of  $A$  is unique. It is easily verified that  $A$  has a left inverse if and only if  $\text{rank } A = n$ ; thus, if  $A$  has a left inverse, then necessarily  $m \geq n$ . An  $m$  by  $n$  sign pattern  $\mathcal{S}$  *allows a positive (resp., nonnegative) left inverse*, provided there exist  $A \in Q(\mathcal{S})$  and a matrix  $B > 0$  (resp.,  $B \geq 0$ ) such that  $BA = I_n$ . Note that if  $P_1$  and  $P_2$  are permutation matrices, then  $\mathcal{S}$  allows a PLI (resp., an NLI) if and only if  $P_1\mathcal{S}P_2$  allows a PLI (resp., an NLI).

\*Received by the editors May 25, 2006; accepted for publication (in revised form) by R. Bhatia October 5, 2006; published electronically April 20, 2007.

<http://www.siam.org/journals/simax/29-2/66091.html>

<sup>†</sup>Department of Mathematics and Statistics, Minnesota State University, Mankato, MN 56001 (in-jae.kim@mnsu.edu). The research of this author was done while he was a postdoctoral fellow at the University of Victoria.

<sup>‡</sup>Department of Computer Science, University of Victoria, P.O. Box 3055, Victoria, BC, Canada V8W 3P6 (dolesky@cs.uvic.ca). The research of this author was supported in part by an NSERC Discovery Grant.

<sup>§</sup>Department of Mathematics, University of Wyoming, Laramie, WY 82071 (bshader@uwyo.edu).

<sup>¶</sup>Department of Mathematics and Statistics, University of Victoria, P.O. Box 3045, Victoria, BC, Canada V8W 3P4 (pvdd@math.uvic.ca). The research of this author was supported in part by an NSERC Discovery Grant.

A motivation for studying PLIs and NLIs comes from determining the qualitative behavior of solutions of  $A^T x = b$  with  $A$  an  $m$  by  $n$  matrix; see, for example, [2, Chapter 1] and [5] for applications in economics. Specifically,  $A$  has a PLI (resp., an NLI) if and only if for each  $n$  by 1 nonzero vector  $b \geq 0$  there exists an  $m$  by 1 vector  $x > 0$  (resp.,  $x \geq 0$ ) satisfying  $A^T x = b$  or equivalently  $x^T A = b^T$ ; see Proposition 4.1 for a proof.

Square sign patterns with entries in  $\{+, -\}$  that allow a positive (left) inverse are characterized in [6], and this characterization is extended to arbitrary square sign patterns in [4]. These results are summarized in [2, section 9.2]. In section 2, we characterize nonsquare sign patterns that allow a PLI, and combine the square and nonsquare characterizations. In section 3, we discuss sign patterns that allow an NLI. More specifically, we characterize nonnegative sign patterns and  $m$  by 2 sign patterns with  $m \geq 2$  that allow an NLI, and present some necessary or sufficient conditions for general  $m$  by  $n$  sign patterns with  $m \geq n$  to allow an NLI. We conclude with some remarks in section 4.

**2. Positive left inverses.** We begin this section with a necessary and sufficient condition for a column sign pattern to allow a PLI or an NLI.

PROPOSITION 2.1. *Let  $\mathcal{S} = (s_1, s_2, \dots, s_m)^T$  be an  $m$  by 1 sign pattern. Then the following are equivalent:*

- (i)  $\mathcal{S}$  has a + entry.
- (ii)  $\mathcal{S}$  allows a PLI.
- (iii)  $\mathcal{S}$  allows an NLI.

*Proof.* Suppose there is an index  $i \in \{1, 2, \dots, m\}$  with  $s_i = +$ . For  $j \in \{1, \dots, m\}$ , set

$$a_j = \begin{cases} 1 & \text{if } j \neq i \text{ and } s_j = +, \\ -1 & \text{if } j \neq i \text{ and } s_j = -, \\ 0 & \text{if } j \neq i \text{ and } s_j = 0, \\ 1 + \sum_{k \neq i} |a_k| & \text{if } j = i. \end{cases}$$

Then  $A = (a_1, \dots, a_m)^T \in Q(\mathcal{S})$ , and  $(1, 1, \dots, 1)A = 1 + \sum_{k \neq i} (|a_k| + a_k) = c > 0$ . This implies that  $\frac{1}{c}(1, 1, \dots, 1)$  is a PLI of  $A$ . Thus,  $\mathcal{S}$  allows a PLI.

It is clear that (ii) implies (iii). Next, suppose that the sign pattern  $\mathcal{S}$  allows an NLI. Then there exist  $A = (a_1, \dots, a_m)^T \in Q(\mathcal{S})$  and  $B = (b_1, \dots, b_m) \geq 0$  such that  $BA = 1$ , i.e.,  $\sum_{j=1}^m b_j a_j = 1 > 0$ . This implies that there exists an  $i$  with  $b_i a_i > 0$ . Since  $b_i \geq 0$ , it follows that  $b_i > 0$ ; hence  $a_i > 0$  and thus  $s_i = +$ .  $\square$

We now consider  $m \geq n \geq 2$ . The following two lemmas give necessary conditions for a sign pattern to allow a PLI.

LEMMA 2.2. *Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with  $n \geq 2$ . If  $\mathcal{S}$  allows a PLI, then each column of  $\mathcal{S}$  has a + and a - entry.*

*Proof.* Suppose that there exist  $A \in Q(\mathcal{S})$  and an  $n$  by  $m$  positive matrix  $B$  such that  $BA = I_n$ . Let  $i \in \{1, 2, \dots, n\}$ . Since the  $(i, i)$ -entry of  $BA$  is 1 and each entry of  $B$  is positive, it follows that some entry in column  $i$  of  $A$  is positive. Hence, column  $i$  of  $\mathcal{S}$  has a + entry.

Since  $n \geq 2$ , there exists  $j \in \{1, \dots, n\}$  with  $j \neq i$ . The  $(j, i)$ -entry of  $BA$  is 0, so since  $B > 0$  and at least one entry of column  $i$  of  $A$  is positive, it follows that at least one entry of column  $i$  of  $A$  must be negative. Thus, column  $i$  of  $\mathcal{S}$  has a - entry.  $\square$

An  $m$  by  $n$  sign pattern  $\mathcal{S}$  with  $n \geq 2$  is *strong Hall*, provided that for every nonempty proper subset  $\gamma$  of  $\{1, 2, \dots, n\}$  the submatrix of  $\mathcal{S}$  consisting of the columns

indexed by  $\gamma$  has nonzero entries in at least  $|\gamma| + 1$  rows (see [3]). Note that if  $\mathcal{S}$  is strong Hall, then necessarily  $m \geq n$ . Also, for  $m \geq n$ ,  $\mathcal{S}$  is not strong Hall if and only if there exist permutation matrices  $P_1$  and  $P_2$  such that

$$(2.1) \quad P_1 \mathcal{S} P_2 = \begin{bmatrix} \mathcal{S}_{11} & \mathcal{S}_{12} \\ O & \mathcal{S}_{22} \end{bmatrix},$$

where  $\mathcal{S}_{11}$  is a  $k$  by  $\ell$  sign pattern for some integers  $k, \ell$  with  $n > \ell \geq 1$  and  $k \leq \ell$ .

LEMMA 2.3. *Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with  $n \geq 2$ . If  $\mathcal{S}$  allows a PLI, then  $\mathcal{S}$  is strong Hall.*

*Proof.* To prove the contrapositive, assume that  $\mathcal{S}$  is not strong Hall. If  $m < n$ , then it is clear that  $\mathcal{S}$  does not allow a PLI. Otherwise, without loss of generality, we may assume that  $\mathcal{S}$  has the form (2.1). If  $k < \ell$ , then the first  $\ell$  columns of each realization of  $\mathcal{S}$  are linearly dependent, and hence  $\mathcal{S}$  does not allow a PLI.

Otherwise,  $k = \ell < n$ . Suppose that there exists a matrix  $A = \begin{bmatrix} A_{11} & A_{12} \\ O & A_{22} \end{bmatrix} \in Q(\mathcal{S})$  with a left inverse  $B = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$ , where  $B_{11}$  is an  $\ell$  by  $\ell$  matrix. Clearly, the  $\ell$  by  $\ell$  matrix  $A_{11}$  is invertible, and by  $BA = I_n$ , it follows that  $\begin{bmatrix} B_{21} & B_{22} \end{bmatrix} \begin{bmatrix} A_{11} \\ O \end{bmatrix} = O$ . Thus,  $B_{21}A_{11} = O$ , and since  $A_{11}$  is invertible, the  $(n - \ell)$  by  $\ell$  matrix  $B_{21} = O$ . Since  $n - \ell \geq 1$  and  $\ell \geq 1$ , every left inverse of a matrix in  $Q(\mathcal{S})$  has a zero entry, and hence  $\mathcal{S}$  does not allow a PLI.  $\square$

Note that if  $\mathcal{S}$  is a square sign pattern of order  $n \geq 2$ , then  $\mathcal{S}$  is strong Hall if and only if  $\mathcal{S}$  is fully indecomposable (see [3]), and  $\mathcal{S}$  allows a PLI if and only if  $\mathcal{S}$  allows a positive inverse. The next theorem, first proved in [4], provides a characterization of square sign patterns that allow a positive inverse. In order to recall this characterization, we use the following definition as in [1] and [2]. Let  $\mathcal{S} = [s_{ij}]$  be an  $m$  by  $n$  sign pattern. The *bipartite digraph*  $D(\mathcal{S})$  of  $\mathcal{S}$  is the digraph with row vertices  $u_1, \dots, u_m$ , column vertices  $v_1, \dots, v_n$ , an arc  $u_i \rightarrow v_j$  if  $s_{ij} = +$ , and an arc  $v_j \rightarrow u_i$  if  $s_{ij} = -$ . Note that there exists at most one arc between  $u_i$  and  $v_j$ .

THEOREM 2.4 (see [2, Theorem 9.2.1]). *An  $n$  by  $n$  square sign pattern  $\mathcal{S}$  with  $n \geq 2$  allows a positive (left) inverse if and only if  $\mathcal{S}$  is strong Hall and the bipartite digraph  $D(\mathcal{S})$  of  $\mathcal{S}$  is strongly connected.*

Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern and let  $D(\mathcal{S})$  be its bipartite digraph. A *strong component* of  $D(\mathcal{S})$  is a maximal strongly connected subdigraph of  $D(\mathcal{S})$ . If  $\alpha$  is a strong component of  $D(\mathcal{S})$ , then  $|\alpha|$  denotes the number of vertices in  $\alpha$ .

Remark 2.5. Let  $\alpha$  be a strong component of  $D(\mathcal{S})$ . Since  $D(\mathcal{S})$  is a bipartite digraph with no cycles of length 2, it follows that if  $|\alpha| \geq 2$ , then  $\alpha$  has at least two row vertices and at least two column vertices.

Let  $\alpha_1, \alpha_2, \dots, \alpha_t$  be the strong components of  $D(\mathcal{S})$ . The *condensed* digraph  $CD(\mathcal{S})$  of  $\mathcal{S}$  has vertices  $\alpha_1, \alpha_2, \dots, \alpha_t$  and an arc  $\alpha_i \rightarrow \alpha_j$  if and only if  $i \neq j$  and  $D(\mathcal{S})$  has at least one arc from a vertex in  $\alpha_i$  to a vertex in  $\alpha_j$ . A strong component  $\alpha_i$  of  $D(\mathcal{S})$  is a *source* if there is no arc coming into  $\alpha_i$  in  $CD(\mathcal{S})$  and there is at least one arc coming out of  $\alpha_i$  in  $CD(\mathcal{S})$ ;  $\alpha_i$  is a *sink* if there is no arc coming out of  $\alpha_i$  in  $CD(\mathcal{S})$  and there is at least one arc coming into  $\alpha_i$  in  $CD(\mathcal{S})$ ; and  $\alpha_i$  is *isolated* if there are no arcs coming into or out of  $\alpha_i$  in  $CD(\mathcal{S})$ .

LEMMA 2.6. *Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern which has a  $+$  and a  $-$  entry in each column and no rows of zeros. Then the following hold for  $D(\mathcal{S})$ :*

- (i) *Each sink and source strong component of  $D(\mathcal{S})$  has at least one row vertex.*
- (ii) *Each isolated strong component has at least two row vertices.*

*Proof.* (i) Let  $\alpha$  be a sink or source strong component. If  $|\alpha| = 1$ , then since each column of  $\mathcal{S}$  has a  $+$  and a  $-$  entry, it follows that no sink or source strong component

consists of exactly one column vertex. Hence,  $\alpha$  is a row vertex. If  $|\alpha| \geq 2$ , then Remark 2.5 implies that  $\alpha$  has at least one row vertex.

(ii) By the assumptions on the rows and columns of  $\mathcal{S}$ , there is no isolated strong component with exactly one vertex. Hence, by Remark 2.5, each isolated strong component has at least two row vertices.  $\square$

Let  $A$  be an  $m$  by  $n$  matrix with  $m \geq n$ . If there exists an  $m$  by 1 vector  $y > 0$  satisfying  $y^T A = 0$ , then  $y^T$  is a *positive left nullvector* of  $A$ . The following theorem gives a characterization of nonsquare sign patterns with no rows of zeros that allow a PLI. Note that conditions for such a sign pattern to allow a PLI are weaker than those for square sign patterns (Theorem 2.4), although the bipartite digraph is used in our proof for a nonsquare sign pattern.

**THEOREM 2.7.** *For  $m > n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with no rows of zeros. Then the following are equivalent:*

- (i) *There exists a matrix  $A \in Q(\mathcal{S})$  with a PLI and a positive left nullvector.*
- (ii)  *$\mathcal{S}$  allows a PLI.*
- (iii) *Each column of  $\mathcal{S}$  has a + and a - entry, and  $\mathcal{S}$  is strong Hall.*

*Proof.* Clearly, (i) implies (ii). By Lemmas 2.2 and 2.3, (ii) implies (iii).

To prove that (iii) implies (i), assume that  $\mathcal{S}$  is strong Hall and that  $\mathcal{S}$  has a + and a - entry in each column. We claim that it suffices to show that there exists an  $m$  by  $(m - n)$  sign pattern  $\mathcal{C}$  so that the  $m$  by  $m$  sign pattern  $[\mathcal{S} \mid \mathcal{C}]$  allows a positive (left) inverse. To prove this claim, suppose there exists an  $m$  by  $m$  matrix  $[A \mid C] \in Q([\mathcal{S} \mid \mathcal{C}])$  with a positive (left) inverse  $\begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$  where  $B_1$  is an  $n$  by  $m$  positive matrix and  $B_2$  is an  $(m - n)$  by  $m$  positive matrix. Then  $B_1 A = I_n$  and hence  $B_1$  is a PLI of  $A$ , implying that  $\mathcal{S}$  allows a PLI. In addition, since  $B_2 A = O$  and  $B_2$  has at least one positive row,  $A$  has a positive left nullvector. Therefore, by Theorem 2.4, it suffices to find an  $m$  by  $(m - n)$  sign pattern  $\mathcal{C}$  such that the  $m$  by  $m$  sign pattern  $[\mathcal{S} \mid \mathcal{C}]$  is strong Hall and its bipartite digraph  $D([\mathcal{S} \mid \mathcal{C}])$  is strongly connected.

Consider the bipartite digraph  $D(\mathcal{S})$  of  $\mathcal{S}$ . Let  $\alpha_1, \alpha_2, \dots, \alpha_t$  be its strong components, where  $\alpha_1, \dots, \alpha_k$  are sinks,  $\alpha_{k+1}, \dots, \alpha_{k+\ell}$  are sources,  $\alpha_{k+\ell+1}, \dots, \alpha_{k+\ell+r}$  are isolated, and  $\alpha_{k+\ell+r+1}, \dots, \alpha_t$  are neither sinks, sources, nor isolated strong components. By Lemma 2.6 (i), each sink and source strong component has a row vertex. Let  $r_i$  be a fixed row vertex of  $\alpha_i$  for each  $i \in \{1, \dots, k + \ell\}$ . Also, by Lemma 2.6 (ii), each isolated strong component has at least two row vertices. Let  $r_i^+, r_i^-$  be distinct fixed row vertices of  $\alpha_i$  for each  $i \in \{k + \ell + 1, \dots, k + \ell + r\}$ . Let  $\mathcal{C}_{n+1}$  be the  $m$  by 1 column sign pattern with nonzero  $j$ th coordinate:

$$(2.2) \quad \begin{cases} + & \text{if } u_j \in \{r_1, \dots, r_k\} \cup \{r_{k+\ell+1}^-, \dots, r_{k+\ell+r}^-\}, \\ - & \text{if } u_j \in \{r_{k+1}, \dots, r_{k+\ell}\} \cup \{r_{k+\ell+1}^+, \dots, r_{k+\ell+r}^+\}, \\ + & \text{otherwise.} \end{cases}$$

Then  $D([\mathcal{S} \mid \mathcal{C}_{n+1}])$  is obtained from  $D(\mathcal{S})$  by appending a new column vertex  $c_{n+1}$ , and arcs  $r_j \rightarrow c_{n+1}$  if  $r_j$  is in a sink component;  $c_{n+1} \rightarrow r_j$  if  $r_j$  is in a source component;  $r_j^- \rightarrow c_{n+1}$  and  $c_{n+1} \rightarrow r_j^+$  if  $r_j^-$  and  $r_j^+$  are in the same isolated component; as well as some additional arcs coming into vertex  $c_{n+1}$ .

To prove that  $D([\mathcal{S} \mid \mathcal{C}_{n+1}])$  is strongly connected, we show that for each vertex  $w$  of  $D(\mathcal{S})$  there exists in  $D([\mathcal{S} \mid \mathcal{C}_{n+1}])$  a walk from  $c_{n+1}$  to  $w$  and a walk from  $w$  to  $c_{n+1}$ . Note that if  $w$  is not in an isolated strong component of  $D(\mathcal{S})$ , then there is a walk from  $w$  to a vertex in a sink strong component  $\alpha_i$  of  $D(\mathcal{S})$  ( $i \in \{1, \dots, k\}$ ). Since  $\alpha_i$  is strongly connected, this walk from  $w$  can be extended to the fixed row vertex  $r_i$  of  $\alpha_i$ . By (2.2), there is an arc  $r_i \rightarrow c_{n+1}$  in  $D([\mathcal{S} \mid \mathcal{C}_{n+1}])$ . Hence, there is

a walk from  $w$  to  $c_{n+1}$ . Similarly, there is a walk from  $c_{n+1}$  to  $w$ .


Next, suppose that  $w$  is a vertex in an isolated strong component  $\alpha_i$  in  $D(\mathcal{S})$  ( $i \in \{k + \ell + 1, \dots, k + \ell + r\}$ ). Since  $\alpha_i$  is strongly connected, there is a walk from  $w$  to the fixed row vertex  $r_i^-$  of  $\alpha_i$ . By (2.2), there are arcs  $r_i^- \rightarrow c_{n+1}$  and  $c_{n+1} \rightarrow r_i^+$  in  $D([\mathcal{S} \mid \mathcal{C}_{n+1}])$ . Since  $\alpha_i$  is strongly connected, there is a walk from  $r_i^+$  to  $w$ . Thus, there exist a walk from  $w$  to  $c_{n+1}$  and a walk from  $c_{n+1}$  to  $w$ .

Finally, define  $\mathcal{C}_{n+2}, \dots, \mathcal{C}_m$  to be  $m$  by 1 column sign patterns, each having no zeros, at least one +, and at least one - entry. Then it is easily verified that  $D([\mathcal{S} \mid \mathcal{C}_{n+1} \mid \dots \mid \mathcal{C}_m])$  is strongly connected. Since  $\mathcal{S}$  is strong Hall and  $[\mathcal{C}_{n+1} \mid \dots \mid \mathcal{C}_m]$  has no zeros, it is clear that  $[\mathcal{S} \mid \mathcal{C}_{n+1} \mid \dots \mid \mathcal{C}_m]$  is strong Hall, completing the proof.  $\square$

*Example 2.8.* Consider the 6 by 4 sign pattern

$$\mathcal{S} = \begin{bmatrix} + & - & 0 & 0 \\ - & + & 0 & 0 \\ + & - & 0 & 0 \\ 0 & 0 & + & - \\ 0 & 0 & - & + \\ 0 & 0 & 0 & - \end{bmatrix}$$

with



Each column of  $\mathcal{S}$  has a + and a - entry, and  $\mathcal{S}$  is strong Hall. Thus, by Theorem 2.7,  $\mathcal{S}$  allows a PLI. However,  $D(\mathcal{S})$  is not strongly connected, illustrating a distinction between the nonsquare and square cases (see Theorem 2.4). In fact,  $D(\mathcal{S})$  has one sink strong component  $\alpha_1$  that consists of vertex  $u_6$ , one source strong component  $\alpha_2$  with vertices  $u_4, u_5, v_3, v_4$ , and one isolated strong component  $\alpha_3$  with vertices  $u_1, u_2, u_3, v_1, v_2$ . Taking  $r_1 = u_6$ ,  $r_2 = u_5$ ,  $r_3^+ = u_1$ , and  $r_3^- = u_2$  in the proof of Theorem 2.7, it follows that

$$\mathcal{C}_5 = \begin{bmatrix} - \\ + \\ + \\ + \\ - \\ + \end{bmatrix}.$$

The last column  $\mathcal{C}_6$  can be taken to be any 6 by 1 column having a + and a - entry, and no zeros. Let  $\mathcal{C} = [\mathcal{C}_5 \mid \mathcal{C}_6]$ . In order to determine a matrix  $[A \mid C] \in Q([\mathcal{S} \mid \mathcal{C}])$  with a positive (left) inverse  $\begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$ , the algorithm described in the proof of [2, Theorem 9.2.1] can be used. Then  $B_1$  is a PLI of  $A$ , and the rows of  $B_2$  are positive left nullvectors of  $A$ .

The next lemma is used to prove Theorem 2.10, in which square and nonsquare cases are combined.

LEMMA 2.9. *Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with  $n \geq 2$  and let  $\mathcal{T}$  be the sign pattern obtained from  $\mathcal{S}$  by deleting the rows of zeros in  $\mathcal{S}$ . Then*

- (i)  $\mathcal{S}$  is strong Hall if and only if  $\mathcal{T}$  is strong Hall, and
- (ii)  $\mathcal{S}$  allows a positive (nonnegative) left inverse if and only if  $\mathcal{T}$  allows a positive (nonnegative) left inverse.

*Proof.* Without loss of generality, assume that  $\mathcal{S} = \begin{bmatrix} \mathcal{T} \\ \mathcal{O} \end{bmatrix}$ . The proof of (i) follows immediately from the definition of strong Hall.

To prove (ii), suppose first that  $\mathcal{S}$  allows a PLI. Let  $A_1 \in Q(\mathcal{T})$  and  $A = \begin{bmatrix} A_1 \\ \mathcal{O} \end{bmatrix} \in Q(\mathcal{S})$  have  $B = [B_1 \ B_2]$  as a PLI. Then  $B_1 A_1 = I_n$  and hence  $\mathcal{T}$  allows a PLI. Next, suppose that  $\mathcal{T}$  allows a PLI. Let  $A_1 \in Q(\mathcal{T})$  have  $B_1$  as a PLI. With  $J$  denoting the all 1's matrix, it is easily verified that  $B = [B_1 \ J]$  is a PLI for  $A = \begin{bmatrix} A_1 \\ \mathcal{O} \end{bmatrix} \in Q(\mathcal{S})$ . Hence,  $\mathcal{S}$  allows a PLI. The nonnegative case can be shown by a similar argument to that above.  $\square$

THEOREM 2.10. *Let  $m \geq n \geq 2$ . The  $m$  by  $n$  sign pattern  $\mathcal{S}$  allows a PLI if and only if*

- (i) each column of  $\mathcal{S}$  has a + and a - entry;
- (ii)  $\mathcal{S}$  is strong Hall; and
- (iii) the bipartite digraph  $D(\mathcal{S}_1)$  of  $\mathcal{S}_1$  is strongly connected whenever  $\mathcal{S}$  is permutationally equivalent to  $\begin{bmatrix} \mathcal{S}_1 \\ \mathcal{O} \end{bmatrix}$  and  $\mathcal{S}_1$  is an  $n$  by  $n$  sign pattern.

*Proof.* For the necessity, suppose that  $\mathcal{S}$  allows a PLI. Then (i) and (ii) follow from Lemmas 2.2 and 2.3, and (iii) follows from Theorem 2.4 and Lemma 2.9 (ii).

For the sufficiency, first assume  $m = n$ . Then  $\mathcal{S}$  is permutationally equivalent to  $\mathcal{S}_1$ , and by Theorem 2.4 the result follows from (ii) and (iii). Next, suppose that  $m > n$ . If  $\mathcal{S}$  has no rows of zeros, then, by Theorem 2.7, the result follows from (i) and (ii). Otherwise, without loss of generality, assume that  $\mathcal{S} = \begin{bmatrix} \mathcal{T} \\ \mathcal{O} \end{bmatrix}$ , where  $\mathcal{T}$  has no rows of zeros. By Lemma 2.9 (i), it follows from (ii) that  $\mathcal{T}$  is strong Hall. Thus, if  $\mathcal{T}$  is an  $n$  by  $n$  sign pattern, then (iii) and Theorem 2.4 imply that  $\mathcal{T}$  allows a PLI. By Lemma 2.9 (ii), this implies that  $\mathcal{S}$  allows a PLI. Otherwise, since it follows from (i) that each column of  $\mathcal{T}$  has a + and a - entry, Theorem 2.7 implies that  $\mathcal{T}$  allows a PLI. Therefore, by Lemma 2.9 (ii),  $\mathcal{S}$  allows a PLI.  $\square$

Remark 2.11. For  $m \geq n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern. Then the following hold:

- (i) If  $\mathcal{S}$  satisfies (i), (ii), and (iii) in Theorem 2.10, then so does every superpattern of  $\mathcal{S}$ . Hence, if  $\mathcal{S}$  allows a PLI, then every superpattern of  $\mathcal{S}$  allows a PLI.
- (ii) Suppose that  $\mathcal{S} = \begin{bmatrix} \mathcal{S}_1 \\ \mathcal{O} \end{bmatrix}$ , where  $\mathcal{S}_1$  is a square sign pattern, satisfies (iii) in Theorem 2.10. Then, in contrast with Theorem 2.7 (i), there is no matrix  $A = \begin{bmatrix} A_1 \\ \mathcal{O} \end{bmatrix} \in Q(\mathcal{S})$  with a PLI that also has a positive left nullvector, since the equation  $[y^T \ z^T] \begin{bmatrix} A_1 \\ \mathcal{O} \end{bmatrix} = 0$  and the fact that  $A_1$  is nonsingular together imply that  $y = 0$ .

The following theorem gives sufficient conditions for an  $m$  by  $n$  sign pattern with  $m > n \geq 1$  to have a realization with a PLI and a positive left nullvector.

THEOREM 2.12. *Let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with  $m > n$  and let  $\mathcal{T}$  be the  $t$  by  $n$  sign pattern obtained from  $\mathcal{S}$  by deleting the rows of zeros in  $\mathcal{S}$ .*

- (i) *If  $n = 1$  and  $\mathcal{T}$  has a + and a - entry, then there exists a matrix in  $Q(\mathcal{S})$  with a PLI and a positive left nullvector.*
- (ii) *If  $t > n \geq 2$  and  $\mathcal{T}$  allows a PLI, then there exists a matrix in  $Q(\mathcal{S})$  with a PLI and a positive left nullvector.*

*Proof.* (i) By Proposition 2.1, a + entry implies the existence of  $A \in Q(\mathcal{S})$  with a

PLI. Since  $A$  has a positive and a negative entry, it can be easily verified that  $A$  has a positive left nullvector.

(ii) When  $m = t$ , the result follows by Theorem 2.7. If  $m > t > n$ , then Theorem 2.7 implies that there exists a matrix  $A \in Q(\mathcal{T})$  with a PLI  $B$  and a positive left nullvector  $y^T$ . Note that the positive matrix  $[B \mid J]$  is a PLI and the vector  $[y^T \ 1 \cdots 1]$  is a positive left nullvector of the matrix  $\begin{bmatrix} A \\ O \end{bmatrix} \in Q(\mathcal{S})$ . Hence, the result follows.  $\square$

**3. Nonnegative left inverses.** In this section we determine structures of nonsquare sign patterns that allow an NLI, as well as structures of NLIs.

For  $m \geq n$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with a realization of rank  $n$ . Then, by induction, it can be shown that  $\mathcal{S}$  is permutationally equivalent to

$$(3.1) \quad \begin{bmatrix} \mathcal{S}_{11} & \mathcal{S}_{12} & \cdots & \mathcal{S}_{1k} \\ O & \mathcal{S}_{22} & \cdots & \mathcal{S}_{2k} \\ \vdots & & \ddots & \vdots \\ O & \cdots & O & \mathcal{S}_{kk} \end{bmatrix},$$

where  $k \geq 1$ ,  $\mathcal{S}_{ii}$  is a square fully indecomposable sign pattern for each  $i \in \{1, \dots, k-1\}$ , and  $\mathcal{S}_{kk}$  is strong Hall. Note that  $\mathcal{S}$  is strong Hall if and only if  $k = 1$ . If  $\mathcal{S}$  is an  $n$  by  $n$  fully indecomposable sign pattern, then  $\mathcal{S}$  allows a nonnegative (left) inverse if and only if  $\mathcal{S}$  allows a positive (left) inverse; see [2, Theorems 9.2.1 and 9.2.3]. In addition, [2, Theorem 9.2.6] provides a complete characterization of  $n$  by  $n$  partly decomposable sign patterns that allow a nonnegative (left) inverse.

*Remark 3.1.* Suppose  $m > n$ . Let  $\mathcal{S}'$  be the square submatrix of  $\mathcal{S}$  obtained by deleting the columns and rows associated with  $\mathcal{S}_{kk}$ . Suppose that  $\mathcal{S}$  allows an NLI. Then the square sign pattern  $\mathcal{S}'$  also allows an NLI. Hence, for  $k = 2$ ,  $\mathcal{S}'$  is fully indecomposable and must satisfy one of the equivalent conditions in [2, Theorem 9.2.1] (see also Theorem 2.4), and for  $k \geq 3$ ,  $\mathcal{S}'$  is partly decomposable and must satisfy the conditions in [2, Theorem 9.2.6]. Furthermore, by an argument similar to that in the proof of Lemma 2.3, it is easily verified that an NLI  $B$  of a matrix in  $Q(\mathcal{S})$  has the block form  $B = [B_{ij}]$  with  $1 \leq i, j \leq k$  and the  $(i, j)$ -block  $B_{ij} = O$  whenever  $i > j$ . Thus, it follows that the strong Hall sign pattern  $\mathcal{S}_{kk}$  also allows an NLI.

We now investigate various necessary and/or sufficient conditions for a strong Hall nonsquare sign pattern to allow an NLI. We first consider strong Hall sign patterns with a  $+$  and a  $-$  entry in each column.

**PROPOSITION 3.2.** *For  $m > n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  strong Hall sign pattern with a  $+$  and a  $-$  entry in each column, and let  $\mathcal{T}$  be the  $t$  by  $n$  sign pattern obtained from  $\mathcal{S}$  by deleting the rows of zeros in  $\mathcal{S}$ . If  $t > n$ , then  $\mathcal{S}$  allows an NLI. If  $t = n$ , then  $\mathcal{S}$  allows an NLI if and only if  $D(\mathcal{T})$  is strongly connected.*

*Proof.* The result follows directly from Theorem 2.10 and the fact that if  $\mathcal{S}$  allows a PLL, then  $\mathcal{S}$  allows an NLI.  $\square$

Let  $\mathcal{I}_n$  denote the  $n$  by  $n$  sign pattern with  $I_n$  as a realization, i.e.,  $I_n \in Q(\mathcal{I}_n)$ . Clearly,  $\mathcal{I}_n$  allows an NLI. Thus, in order to allow an NLI, an  $m$  by  $n$  sign pattern with  $m \geq n$  need not have a  $-$  entry in each column as is required to allow a PLI (see Lemma 2.2), but clearly must have a  $+$  entry in each column. We first consider the case that  $\mathcal{S}$  has a nonnegative column having only  $+$  or  $0$  entries. For ease of notation, we sometimes use  $(M)_{ij}$  to denote the  $(i, j)$ -entry of a matrix  $M$ .

**PROPOSITION 3.3.** *For  $m \geq n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern with at least one nonnegative column. If  $\mathcal{S}$  allows an NLI, then each nonnegative column has at most  $m - n + 1$  positive entries.*



*Proof.* Let  $B$  be an NLI of  $A \in Q(\mathcal{S})$ , and let  $t$  be the number of positive entries in any nonnegative column of  $A$ . Without loss of generality, assume that the first column of  $A$  is a nonnegative column with its first  $t$  entries positive. Since  $(BA)_{h1} = 0$  for each  $h \in \{2, \dots, n\}$ , it follows that  $B$  has the block form  $B = [B_{ij}]$  with  $1 \leq i, j \leq 2$ , where the  $(2, 1)$ -block  $B_{21}$  is the  $(n - 1)$  by  $t$  zero matrix. Hence, the equality  $\text{rank } B = n$  implies that the rank of the  $(n - 1)$  by  $(m - t)$  matrix  $B_{22}$  is  $n - 1$ . Thus,  $n - 1 \leq m - t$  and the result follows.  $\square$

If all columns are nonnegative, then the following result gives a necessary and sufficient condition for such a sign pattern to allow an NLI.

**THEOREM 3.4.** *For  $m \geq n \geq 1$ , let  $\mathcal{S}$  be an  $m$  by  $n$  nonnegative sign pattern. Then  $\mathcal{S}$  allows an NLI if and only if  $\mathcal{S}$  is permutationally equivalent to*

$$\begin{bmatrix} \mathcal{I}_n \\ \mathcal{T} \end{bmatrix},$$

where  $\mathcal{T}$  is an  $(m - n)$  by  $n$  nonnegative sign pattern.

*Proof.* The case  $n = 1$  follows directly from Proposition 2.1. Suppose that  $n \geq 2$ .

For the sufficiency, assume without loss of generality that

$$\mathcal{S} = \begin{bmatrix} \mathcal{I}_n \\ \mathcal{T} \end{bmatrix}.$$

Let  $T \in Q(\mathcal{T})$  and  $A = \begin{bmatrix} \mathcal{I}_n \\ T \end{bmatrix} \in Q(\mathcal{S})$ . Since  $[I_n \mid O]A = I_n$ , it follows that  $\mathcal{S}$  allows an NLI.

For the necessity, suppose that  $\mathcal{S} = [s_{ij}]$  allows an NLI; i.e., there exist  $A = [a_{ij}] \in Q(\mathcal{S})$  and an  $n$  by  $m$  nonnegative matrix  $B = [b_{ij}]$  such that  $BA = I_n$ . Let  $i \in \{1, \dots, n\}$ . Since  $(BA)_{ii} = 1$ , there exists  $j_i \in \{1, \dots, m\}$  such that  $b_{ij_i} a_{j_i i} > 0$ . This implies that  $s_{j_i i} = +$ . Also, for each  $k \in \{1, \dots, n\} \setminus \{i\}$ ,  $(BA)_{ik} = 0$  implies that  $b_{ij_i} a_{j_i k} = 0$ . Thus, row  $j_i$  of  $\mathcal{S}$  is equal to row  $i$  of  $\mathcal{I}_n$ . As this holds for each  $i \in \{1, \dots, n\}$ , the result follows.  $\square$

*Remark 3.5.* Let  $\mathcal{S} = \begin{bmatrix} \mathcal{I}_n \\ \mathcal{J} \end{bmatrix}$  be the  $m$  by  $n$  nonnegative sign pattern with  $m \geq n \geq 2$ , where  $\mathcal{J}$  is the sign pattern with all entries positive. Then, by Theorem 3.4,  $\mathcal{S}$  allows an NLI. However, in contrast with Remark 2.11 (i), Theorem 3.4 implies that no nonnegative superpattern of  $\mathcal{S}$  (except  $\mathcal{S}$  itself) allows an NLI.

Next, we consider sign patterns that have at least one nonnegative column and at least one column with a  $+$  and a  $-$  entry. We use  $e_i$  to denote the  $i$ th column vector of an identity matrix.

**THEOREM 3.6.** *For  $m \geq n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern that has  $p \geq 1$  nonnegative columns and  $n - p \geq 1$  columns with a  $+$  and a  $-$  entry. Suppose that  $\mathcal{S}$  allows an NLI. Then  $\mathcal{S}$  is permutationally equivalent to a matrix of the form*

$$(3.2) \quad \begin{bmatrix} \mathcal{I}_p & \mathcal{S}_{12} \\ \mathcal{S}_{21} & \mathcal{S}_{22} \\ O & \mathcal{S}_{32} \end{bmatrix},$$

where  $\mathcal{S}_{21}$  is an  $r$  by  $p$  nonnegative sign pattern with no rows of zeros,  $O$  is an  $s$  by  $p$  zero matrix with  $s \geq 1$ , and each of the last  $n - p$  columns of  $\mathcal{S}$  has a  $+$  and a  $-$  entry. Furthermore, if  $\mathcal{S}$  is strong Hall, then  $\mathcal{S}_{21}$  is not vacuous and has no column of zeros.

*Proof.* Without loss of generality, we may assume that the first  $p$  columns of  $\mathcal{S}$  are nonnegative, and that each of the last  $n - p$  columns of  $\mathcal{S}$  has a  $+$  and a  $-$  entry.

Since  $\mathcal{S}$  allows an NLI, so does the  $m$  by  $p$  nonnegative sign pattern consisting of the first  $p$  columns of  $S$ . Therefore, by Theorem 3.4, we may permute the rows of  $\mathcal{S}$  to obtain a matrix of the form (3.2), where  $S_{21}$  is a nonnegative matrix with no row of zeros,  $O$  is an  $s$  by  $p$  zero matrix with  $s \geq 0$ , and each of the last  $n - p$  columns has a  $+$  and a  $-$  entry.

Let  $A$  be a matrix in  $Q(\mathcal{S})$  that has an NLI, say  $B$ . Since  $BA = I_n$ , each of the vectors  $e_1^T, \dots, e_n^T$  is a nontrivial, nonnegative linear combination of the rows of  $A$ . Since the first  $p$  columns of  $A$  are nonnegative and  $n > p$ , this requires that  $s \geq 1$ , and we conclude that  $\mathcal{S}$  has the desired form.

If  $S_{21}$  is vacuous or has a column of zeros, then  $\mathcal{S}$  has an  $(m - 1)$  by 1 zero submatrix. Hence  $\mathcal{S}$  is not strong Hall, and the result follows by taking the contrapositive.  $\square$

PROPOSITION 3.7. *For  $m \geq n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern that has  $p \geq 1$  nonnegative columns and  $n - p \geq 1$  columns with a  $+$  and a  $-$  entry. Suppose that  $\mathcal{S}$  allows an NLI and has the form (3.2). Let*

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \\ O & A_{32} \end{bmatrix} \in Q(\mathcal{S})$$

have an NLI  $B = \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{21} & B_{22} & B_{23} \end{bmatrix}$ , where each of  $A_{21}, A_{22}, B_{12}$ , and  $B_{22}$  may be vacuous if  $\mathcal{S}$  is not strong Hall. Then the following hold:

- (i)  $B_{11}$  is a diagonal matrix, and  $B_{21}$  and  $B_{22}$  are zero matrices.
- (ii) The sign pattern  $\mathcal{S}_{32}$  allows an NLI.
- (iii) If row  $q$  of  $S_{21}$  has at least two positive entries, then column  $q$  of  $B_{12}$  is a zero column.
- (iv) Each column of  $B_{12}$  has at most one positive entry. Furthermore, the sign pattern of  $B_{12}$  is a subpattern of  $S_{21}^T$ .

*Proof.* Assume that  $S_{21}$  is not vacuous.

Since  $BA = I_n$ , it follows that  $B_{21}A_{11} + B_{22}A_{21} = O$ . Moreover, since  $B_{21}, B_{22}, A_{11}$ , and  $A_{21}$  are nonnegative, and no row of  $A_{11}$  or  $A_{21}$  is all zeros,  $B_{21} = O$  and  $B_{22} = O$ . Also,  $BA = I_n$  implies that  $B_{11}A_{11} + B_{12}A_{21} = I_p$ . Since  $B_{11}, B_{12}, A_{11}$ , and  $A_{21}$  are nonnegative, both  $B_{11}A_{11}$  and  $B_{12}A_{21}$  are diagonal matrices. Since  $A_{11} \in Q(\mathcal{I}_p)$ ,  $A_{11}$  is an invertible diagonal matrix, and hence  $B_{11}$  is a diagonal matrix. Thus, (i) is proven.

Since  $B_{21}$  and  $B_{22}$  are zero matrices, and  $BA = I_n$ ,  $B_{23}$  is an NLI of  $A_{32}$ , and (ii) is proven.

Since  $B_{12}A_{21}$  is a diagonal matrix and  $B_{12}$  is nonnegative, the  $i$ th row of  $B_{12}A_{21}$  is a nonnegative linear combination of the rows of  $A_{21}$  (weighted by the entries of the  $i$ th row of  $B_{12}$ ). As the  $i$ th row of  $B_{12}A_{21}$  is a nonnegative multiple of  $e_i^T$ , and  $A_{21}$  is a nonnegative matrix with no row of zeros, it follows that if the  $(i, j)$ -entry of  $B_{12}$  is nonzero, then the  $j$ th row of  $A_{21}$  is a multiple of  $e_i^T$ . In particular, this implies that each column of  $B_{12}$  has at most one nonzero entry. If the  $j$ th row of  $A_{21}$  has at least two positive entries, then column  $j$  of  $B_{12}$  is a column of zeros, proving (iii). If the  $(i, j)$ -entry of  $B_{12}$  is nonzero, then the  $(j, i)$ -entry of  $A_{21}$  is nonzero, completing the proof of (iv).

If  $S_{21}$  is vacuous, then  $A_{21}, A_{22}, B_{12}$ , and  $B_{22}$  are vacuous, in which case the proofs of (i) for  $B_{11}, B_{21}$  and (ii) are similar, but statements (i) for  $B_{22}$ , (iii), and (iv) are vacuous.  $\square$

For  $m \geq 2$ , Proposition 3.2, Theorem 3.4, and the following theorem completely characterize the  $m$  by 2 sign patterns that allow an NLI.

**THEOREM 3.8.** *For  $m \geq 2$ , let  $\mathcal{S}$  be an  $m$  by 2 sign pattern such that the first column is nonnegative and the second column has a + and a - entry. Then  $\mathcal{S}$  allows an NLI if and only if the first column of  $\mathcal{S}$  has a + entry and  $[0 +]$  is a row of  $\mathcal{S}$ .*

*Proof.* Suppose that  $\mathcal{S}$  allows an NLI. Then the first column of  $\mathcal{S}$  also allows an NLI. Hence, Theorem 3.4 implies that the first column of  $\mathcal{S}$  has a + entry. By Theorem 3.6, we may assume without loss of generality that  $\mathcal{S}$  is of the form (3.2). Since  $\mathcal{S}_{32}$  is a column sign pattern, Propositions 3.7 (ii) and 2.1 imply that  $\mathcal{S}_{32}$  has a + entry. Hence,  $[0 +]$  is a row of  $\mathcal{S}$ .

For the converse, suppose that the first column of  $\mathcal{S}$  has a + entry and  $[0 +]$  is a row of  $\mathcal{S}$ . Suppose that  $[+ -]$  is also a row of  $\mathcal{S}$ . Then without loss of generality,  $A \in \mathcal{S}$  has the form

$$\begin{bmatrix} a & -b \\ u & v \\ 0 & c \end{bmatrix},$$

where  $a, b, c > 0$ , and  $u$  and  $v$  are  $(m - 2)$  by 1 vectors. It is easy to verify that

$$\begin{bmatrix} 1/a & O & b/ac \\ 0 & O & 1/c \end{bmatrix}$$

is an NLI of  $A$ .

Next suppose that  $[+ -]$  is not a row of  $\mathcal{S}$ . Then without loss of generality,  $A \in \mathcal{S}$  has the form

$$\begin{bmatrix} a & b \\ u & v \\ 0 & -c \\ 0 & d \end{bmatrix},$$

where  $a, c, d > 0$ ,  $b \geq 0$ , and  $u$  and  $v$  are  $(m - 3)$  by 1 vectors. It is easy to verify that

$$\begin{bmatrix} 1/a & O & b/ac & 0 \\ 0 & O & 1/c & 2/d \end{bmatrix}$$

is an NLI of  $A$ .

Hence,  $\mathcal{S}$  allows an NLI.  $\square$

Note that the proof of Theorem 3.8 actually shows that if  $\mathcal{S}$  is an  $m$  by 2 matrix whose first column is nonnegative, second column has a + and a - entry, and  $[0 +]$  is one of its rows, then *each* matrix with sign pattern  $\mathcal{S}$  has an NLI.

*Example 3.9.* The strong Hall sign pattern

$$\mathcal{S} = \begin{bmatrix} + & - \\ + & - \\ 0 & + \end{bmatrix}$$

does not allow a PLI (by Lemma 2.2), but does allow an NLI (by Theorem 3.8) since

$$\begin{bmatrix} 1 & 0 & 1/2 \\ 0 & 0 & 1/2 \end{bmatrix} \begin{bmatrix} 1 & -1 \\ 1 & -1 \\ 0 & 2 \end{bmatrix} = I_2.$$

In general (as noted in the introduction) an NLI is not unique. For instance,

$$\begin{bmatrix} 1/2 & 1/2 & 1/2 \\ 0 & 0 & 1/2 \end{bmatrix}$$

is another NLI of the above matrix.

In the next theorem, it is shown that if a sign pattern  $\mathcal{S}$  of the form (3.2) has a  $(3, 2)$ -block  $\mathcal{S}_{32}$  that allows an NLI or PLI, then some conditions on the negative entries in  $\mathcal{S}_{12}$  insure that  $\mathcal{S}$  allows an NLI.

**THEOREM 3.10.** *For  $m \geq n \geq 2$ , let  $\mathcal{S}$  be an  $m$  by  $n$  sign pattern of the form (3.2) with  $p \geq 1$ ,  $n - p \geq 1$ , and  $\mathcal{S}_{21}$ ,  $\mathcal{S}_{22}$  arbitrary. Then the following hold:*

- (i) *If  $\mathcal{S}_{32}$  allows an NLI and  $\mathcal{S}_{12}$  has only 0 or  $-$  entries, then  $\mathcal{S}$  allows an NLI.*
- (ii) *If  $\mathcal{S}_{32}$  allows a PLI and each row of  $\mathcal{S}_{12}$  has a  $-$  entry, then  $\mathcal{S}$  allows an NLI.*

*Proof.* (i) Let

$$(3.3) \quad A = \begin{bmatrix} I_p & A_{12} \\ A_{21} & A_{22} \\ O & A_{32} \end{bmatrix} \in Q(\mathcal{S}),$$

where  $-A_{12} \geq 0$  and  $A_{32}$  has  $B_{23}$  as an NLI. Let

$$(3.4) \quad B = \begin{bmatrix} I_p & O & B_{13} \\ O & O & B_{23} \end{bmatrix}$$

with  $B_{13} = -A_{12}B_{23}$ , which is a nonnegative matrix. Then  $B \geq 0$ ,  $BA = I_n$ , and hence the result follows.

(ii) Let  $A \in Q(\mathcal{S})$  be of the form (3.3) and let  $B$  be of the form (3.4). If  $B_{23}$  is a PLI of  $A_{32}$  and  $B_{13} = -A_{12}B_{23}$ , then  $B_{13} > 0$ , provided that the negative entries of  $A_{12}$  are sufficiently large in magnitude, and  $BA = I_n$  as required.  $\square$

**4. Concluding remarks.** In section 3, we have characterized nonnegative sign patterns, strong Hall sign patterns with each column having a  $+$  and a  $-$  entry, and  $m$  by 2 sign patterns that allow an NLI. For other cases, we have given some necessary or sufficient conditions for  $\mathcal{S}$  to allow an NLI. A characterization for the blocks of the last column of a sign pattern  $\mathcal{S}$  of the form (3.1) with  $k \geq 2$  that allows an NLI remains open. We conclude by showing (in Theorem 4.2) that some conditions on the submatrix  $\mathcal{S}_{kk}$  of a sign pattern  $\mathcal{S}$  of the form (3.1) with  $k \geq 2$  insure that  $\mathcal{S}$  allows an NLI for arbitrary  $\mathcal{S}_{1k}, \dots, \mathcal{S}_{k-1,k}$ .

Let  $\mathcal{S}$  allow a PLI and  $A \in Q(\mathcal{S})$ . The following proposition, which is used to prove Theorem 4.2, describes a relation between a PLI of  $A$  and the qualitative behavior of solutions of  $x^T A = b^T$ . The latter equation is given in the introduction as motivation for studying PLIs and NLIs.

**PROPOSITION 4.1.** *For  $m \geq n$ , let  $A$  be an  $m$  by  $n$  matrix. Then  $A$  has a PLI if and only if for each  $n$  by 1 nonzero vector  $b \geq 0$  there exists an  $m$  by 1 vector  $x > 0$  satisfying  $x^T A = b^T$ .*

*Proof.* Suppose that an  $n$  by  $m$  matrix  $B > 0$  is a PLI of  $A$ . For an  $n$  by 1 nonzero vector  $b \geq 0$ , it is clear that  $(b^T B)A = b^T$  and  $b^T B > 0$ . Hence, the result follows.

Next, suppose that for each  $n$  by 1 nonzero vector  $b \geq 0$  there exists an  $m$  by 1 vector  $x > 0$  satisfying  $x^T A = b^T$ . Take  $b$  to be the  $i$ th column  $e_i$  of  $I_n$  and let  $x_i > 0$

be a solution of  $x^T A = e_i^T$ . Then the matrix

$$B = \begin{bmatrix} x_1^T \\ \vdots \\ x_n^T \end{bmatrix}$$

is a PLI of  $A$ .  $\square$

**THEOREM 4.2.** For  $m > s \geq 1$ ,  $n > t \geq 1$ , and  $m > n$ , let  $\mathcal{S}_{11}$  be an  $s$  by  $t$  sign pattern that allows an NLI and let  $\mathcal{S}_{22}$  be an  $(m-s)$  by  $(n-t)$  sign pattern that allows a PLI. Suppose that if  $n-t=1$ , then  $\mathcal{S}_{22}$  has a  $-$  entry, and if  $n-t \geq 2$ , then  $\mathcal{S}_{22}$  is not permutationally equivalent to the sign pattern  $\begin{bmatrix} \mathcal{T} \\ \mathcal{O} \end{bmatrix}$  in which  $\mathcal{T}$  is a square sign pattern. Then, for an arbitrary  $s$  by  $(n-t)$  sign pattern  $\mathcal{S}_{12}$ , the sign pattern  $\mathcal{S} = \begin{bmatrix} \mathcal{S}_{11} & \mathcal{S}_{12} \\ \mathcal{O} & \mathcal{S}_{22} \end{bmatrix}$  allows an NLI.

*Proof.* Let  $A_{11}$  be a matrix in  $Q(\mathcal{S}_{11})$  with  $B_{11}$  as an NLI. By Theorem 2.12, there exists  $A_{22} \in Q(\mathcal{S}_{22})$  that has a PLI  $B_{22}$  and a positive left nullvector  $y^T$ . Let  $A_{12} \in Q(\mathcal{S}_{12})$ . Then  $A_{12}$  can be written as  $A_{12} = V_1 - V_2$ , where  $V_1, V_2 \geq 0$  and the entrywise (Hadamard) product  $V_1 \circ V_2 = O$ . Let  $v_i^T \geq 0$  for  $1 \leq i \leq s$  denote row  $i$  of  $V_1$ . If  $v_i \neq 0$ , then by Proposition 4.1 there exists an  $(m-s)$  by 1 vector  $x_i > 0$  such that  $x_i^T A_{22} = v_i^T$ . If  $v_i = 0$ , then  $x_i^T A_{22} = v_i^T = 0$  when  $x_i^T = y^T$ . Thus,  $K_1 = [x_1, \dots, x_s]^T > 0$  and  $K_1 A_{22} = V_1$ . Similarly, there exists  $K_2 > 0$  such that  $K_2 A_{22} = V_2$ .

Let  $A_{12}(\epsilon) = \epsilon V_1 - V_2 = (\epsilon K_1 - K_2) A_{22}$  for a sufficiently small  $\epsilon > 0$  such that  $K_2 - \epsilon K_1 > 0$ . Note that  $V_1 \circ V_2 = O$  implies that  $A_{12}(\epsilon) \in Q(\mathcal{S}_{12})$ . Let  $B_{12} = B_{11}(K_2 - \epsilon K_1)$ . Since  $K_2 - \epsilon K_1 > 0$  and  $B_{11} \geq 0$  with no rows of zeros, it follows that  $B_{12} > 0$ . It can be easily verified that  $\begin{bmatrix} B_{11} & B_{12} \\ \mathcal{O} & B_{22} \end{bmatrix} \begin{bmatrix} A_{11} & A_{12}(\epsilon) \\ \mathcal{O} & A_{22} \end{bmatrix} = I_n$ . Hence, the result follows.  $\square$

**Remark 4.3.** Take  $\mathcal{S}_{11}$  and  $\mathcal{S}_{22}$  in Theorem 4.2 to be  $\mathcal{S}'$  in Remark 3.1 and  $\mathcal{S}_{kk}$  in the form (3.1) with  $k \geq 2$ , respectively. Then the conditions on  $\mathcal{S}_{kk}$  in Theorem 4.2 insure that the sign pattern  $\mathcal{S}$  of the form (3.1) with  $k \geq 2$  allows an NLI for arbitrary  $\mathcal{S}_{1k}, \dots, \mathcal{S}_{k-1,k}$ .

**Acknowledgment.** The authors thank the referee for comments that improved the readability of the paper.

#### REFERENCES

- [1] A. BERMAN AND B. D. SAUNDERS, *Matrices with zero line sums and maximal rank*, Linear Algebra Appl., 40 (1981), pp. 229–235.
- [2] R. A. BRUALDI AND B. L. SHADER, *Matrices of Sign-Solvable Linear Systems*, Cambridge University Press, Cambridge, UK, 1995.
- [3] R. A. BRUALDI AND B. L. SHADER, *Strong Hall matrices*, SIAM J. Matrix Anal. Appl., 15 (1994), pp. 359–365.
- [4] M. FIEDLER AND R. GRONE, *Characterizations of sign patterns of inverse positive matrices*, Linear Algebra Appl., 40 (1981), pp. 237–245.
- [5] T. FUJIMOTO AND R. R. RANADE, *Two characterizations of inverse-positive matrices: The Hawkins-Simon Condition and the Le Chatelier-Braun Principle*, Electron. J. Linear Algebra, 11 (2004), pp. 59–65.
- [6] C. R. JOHNSON, F. T. LEIGHTON, AND H. A. ROBINSON, *Sign patterns of inverse positive matrices*, Linear Algebra Appl., 24 (1979), pp. 75–83.