

1999

## The possible numbers of zeros in a orthogonal matrix

Gi-Sang Cheon  
gscheon@road.daejin.ac.kr

Charles R. Johnson  
crjohnso@math.wm.edu

Sang-Gu Lee  
sglee@yurim.skku.ac.kr

Ethan J. Pribble  
pribble@gladstone.uoregon.edu

Follow this and additional works at: <http://repository.uwyo.edu/ela>

---

### Recommended Citation

Cheon, Gi-Sang; Johnson, Charles R.; Lee, Sang-Gu; and Pribble, Ethan J.. (1999), "The possible numbers of zeros in a orthogonal matrix", *Electronic Journal of Linear Algebra*, Volume 5.  
DOI: <https://doi.org/10.13001/1081-3810.1028>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact [scholcom@uwyo.edu](mailto:scholcom@uwyo.edu).

## THE POSSIBLE NUMBERS OF ZEROS IN AN ORTHOGONAL MATRIX\*

G.-S. CHEON<sup>†</sup>, C. R. JOHNSON<sup>‡</sup>, S.-G. LEE<sup>§</sup>, AND E. J. PRIBBLE<sup>¶</sup>

**Abstract.** It is shown that for  $n \geq 2$  there is an  $n \times n$  indecomposable orthogonal matrix with exactly  $k$  entries equal to zero if and only if  $0 \leq k \leq (n-2)^2$ .

**Key words.** orthogonal matrix, indecomposable matrix, zero-nonzero pattern

**AMS subject classifications.** 15A57, 05C50

**1. Introduction.** By a *pattern* we simply mean the arrangement of zero and nonzero entries in a matrix. An  $n \times n$  pattern  $P$  is called *orthogonal* if there is a (real) orthogonal matrix  $U$  whose pattern is  $P$ . By  $\#(U)$  or  $\#(P)$  we mean the number of zero entries in the matrix  $U$  or pattern  $P$ . An  $n \times n$  pattern (or matrix)  $P$  is called *indecomposable* if it has no  $r \times q$  zero submatrix,  $r + q = n$ ; equivalently, there do not exist permutation matrices  $Q_1$  and  $Q_2$  such that

$$Q_1 P Q_2 = \begin{bmatrix} P_{11} & O \\ P_{21} & P_{22} \end{bmatrix},$$

in which  $P_{11}$  and  $P_{22}$  are square and nonempty (or, equivalently the bipartite graph of  $P$  is connected). If  $P$  were an orthogonal pattern and there were such reducing blocks, then an elementary calculation shows that  $P_{21} = O$  also. Since an  $n \times n$  orthogonal matrix  $U$  is invertible,  $\#(U) \leq n(n-1)$  (which is sharp because the identity is orthogonal), but to be indecomposable,  $U$  must have more nonzero entries. In [BBS], it was observed that the maximum number of zero entries in an  $n \times n$  indecomposable orthogonal matrix,  $n \geq 2$ , is  $(n-2)^2$ , in response to a query made by [F].

What, then, about smaller numbers of zeros? It should be noted that if any single entry is changed to a nonzero in any indecomposable orthogonal pattern  $P$  that realizes  $(n-2)^2$  zeros,  $n \geq 5$ , the resulting pattern is no longer orthogonal. Nonetheless,  $(n-2)^2 - 1$  zeros can occur in an  $n \times n$  indecomposable orthogonal matrix. It is our purpose here to show that there is an  $n \times n$  indecomposable orthogonal matrix

---

\*Received by the editors on 24 September 1998. Accepted for publication on 27 January 1999.  
Handling Editor: Daniel Hershkowitz.

<sup>†</sup>Department of Mathematics, Daejin University, Pocheon 487-711, Korea (gscheon@road.daejin.ac.kr).

<sup>‡</sup>Department of Mathematics, College of William & Mary, Williamsburg, VA 23187-8795, USA (crjohnso@math.wm.edu). The work of this author was supported by the National Science Foundation through NSF REU grant DMS-96-19577.

<sup>§</sup>Department of Mathematics, Sungkyunkwan University, Suwon 440-746, Korea, BSRI-98-1420 (sglee@yurim.skku.ac.kr).

<sup>¶</sup>Department of Mathematics, University of Oregon, Eugene, OR 97403-1222, USA (pribble@gladstone.uoregon.edu). The work of this author was supported by the National Science Foundation through NSF REU grant DMS-96-19577.

$U$  such that  $\#(U) = k$  if and only if  $0 \leq k \leq (n-2)^2$ , thereby greatly strengthening earlier observations. The same is true for complex unitary matrices.

**2. Numbers of Zeros from 0 to  $\frac{1}{2}(n-2)(n-1)$ .** Let  $P$  be an  $n \times n$  indecomposable orthogonal matrix with columns  $p_1, \dots, p_n$ , and let

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

be a  $2 \times 2$  orthogonal matrix with no zero entries. Then it is easy to show that the matrix

$$D_i(P) = \begin{bmatrix} p_1 & \cdots & p_{i-1} & ap_i & bp_i & p_{i+1} & \cdots & p_n \\ 0 & \cdots & 0 & c & d & 0 & \cdots & 0 \end{bmatrix}$$

is an  $(n+1) \times (n+1)$  indecomposable orthogonal matrix. This idea comes from the notions of matrix weaving and woven matrices which can be found in [C].

It should be clear at this point that the above notion may as well be applied to orthogonal patterns. Thus we obtain the following lemma.

**LEMMA 2.1.** *If  $P$  is an  $n \times n$  indecomposable orthogonal pattern, then  $D_i(P)$  is an  $(n+1) \times (n+1)$  indecomposable orthogonal pattern.*

Since for each  $\theta$ ,  $0 < \theta < \frac{\pi}{2}$ ,

$$B(\theta) = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

is an orthogonal matrix, it is clear that there are full (i.e. indecomposable)  $2 \times 2$  orthogonal matrices and that there are ones arbitrarily close to the identity matrix  $I_2$ . It follows that for any  $B(\theta)$  with a sufficiently small  $\theta$  and for any vector  $v \in \mathbb{R}^2$  with no zero components, the row vector  $v^T B(\theta)$  has no zero components.

We denote by  $K_{n,i}$  the  $n \times n$  pattern whose only zero entries are the first  $i$  entries of the last row.

**LEMMA 2.2.** *For  $n \geq 2$ , each  $K_{n,i}$ ,  $i = 0, \dots, n-2$ , is an indecomposable orthogonal pattern.*

*Proof.* First we show that if  $K_{n,i}$  is an orthogonal pattern for  $n \geq 2$  and some integer  $i$  satisfying  $1 \leq i \leq n-2$ , then  $K_{n,i-1}$  is also an orthogonal pattern. For  $n \geq 2$ , suppose there exists an  $n \times n$  orthogonal matrix  $A = (a_{pq})$  and an integer  $i$  satisfying  $1 \leq i \leq n-2$  so that  $A$  has pattern  $K_{n,i}$ . Define  $R_j(\theta)$  to be the  $n \times n$  orthogonal matrix with entries equal to the identity matrix except that

$$R_j(\theta)[\{j, j+1\}] = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

where the notation  $A[\alpha]$  denotes the principal submatrix of  $A$  whose rows and columns are indexed by the set  $\alpha$ .

Now form the product  $AR_i(\theta)$ . Note that  $A$  and  $AR_i(\theta)$  are entrywise equal except for columns  $i$  and  $i + 1$ . These two columns of  $AR_i(\theta)$  are

$$\begin{bmatrix} a_{1,i} \cos(\theta) + a_{1,i+1} \sin(\theta) \\ \vdots \\ a_{n-1,i} \cos(\theta) + a_{n-1,i+1} \sin(\theta) \\ a_{n,i+1} \sin(\theta) \end{bmatrix} \text{ and } \begin{bmatrix} -a_{1,i} \sin(\theta) + a_{1,i+1} \cos(\theta) \\ \vdots \\ -a_{n-1,i} \sin(\theta) + a_{n-1,i+1} \cos(\theta) \\ a_{n,i+1} \cos(\theta) \end{bmatrix},$$

respectively. Since both  $A$  and  $R_i(\theta)$  are orthogonal, the product  $AR_i(\theta)$  is orthogonal. Now we only need to choose some  $\theta$  sufficiently close to 0 so that we do not create any extra zero entries in  $AR_i(\theta)$ . Thus  $AR_i(\theta)$  is an orthogonal matrix with pattern  $K_{n,i-1}$ .

Next we prove the lemma using the above result. We proceed by induction. Assume that for  $n \geq 2$  there exists a full  $n \times n$  orthogonal pattern  $P$ . Note that there is such a pattern for  $n = 2$ . By Lemma 2.1,  $D_n(P)$  is also an orthogonal pattern.  $D_n(P)$  has pattern  $K_{n+1,(n+1)-2}$ . By the above result,  $K_{n+1,i}$ ,  $i = 0, \dots, (n+1) - 2$ , is also an orthogonal pattern. And  $K_{n+1,0}$  is an  $(n+1) \times (n+1)$  full orthogonal pattern, which completes the induction. Note that for  $i$  satisfying  $0 \leq i \leq n-2$ ,  $K_{n,i}$  is indecomposable as well.  $\square$

We now know that iterative application of the operator  $D_j()$  to a  $K_{n,i}$ ,  $0 \leq i \leq n-2$ , will produce indecomposable orthogonal patterns. Certain of these will be of particular interest.

For  $2 \leq m \leq n$  and  $0 \leq i \leq m-2$ , we let

$$H_{n,m,i} = D_{n-1}(D_{n-2}(\cdots D_m(K_{m,i}) \cdots)).$$

Then we obtain the following immediate corollary to Lemmas 2.1 and 2.2.

**COROLLARY 2.3.** *Each  $H_{n,m,i}$ ,  $2 \leq m \leq n$ ,  $0 \leq i \leq m-2$  is an indecomposable orthogonal pattern.*

We note that since  $H_{n,2,0}$  is the full  $n \times n$  (upper) Hessenberg pattern, it follows that this pattern with  $\#(H_{n,2,0}) = \frac{1}{2}(n-2)(n-1)$  is orthogonal. This is the sparsest pattern among the  $H_{n,m,i}$  and its indecomposable orthogonality will also be used in the next section.

**COROLLARY 2.4.** *For each  $k = 0, \dots, \frac{1}{2}(n-2)(n-1)$ , there is an  $n \times n$  indecomposable orthogonal matrix with exactly  $k$  zero entries.*

*Proof.* We count the number of zeros in each  $H_{n,m,i}$  where  $2 \leq m \leq n$  and  $0 \leq i \leq m-2$ .  $K_{m,i}$  has  $i$  zeros,  $D_m(K_{m,i})$  has  $i + ((m+1) - 2)$  zeros and so on. So we have

$$\begin{aligned} \#(H_{n,m,i}) &= i + ((m+1) - 2) + ((m+2) - 2) + \cdots + ((m + (n-m)) - 2) \\ &= i + (m-1) + m + \cdots + (n-2) \\ &= i + \frac{1}{2}(n-2)(n-1) - \frac{1}{2}(m-2)(m-1). \end{aligned}$$

Now it is clear that we do indeed get all numbers of zeros between 0 and  $\frac{1}{2}(n-2)(n-1)$  as we let  $m$  and  $i$  vary.  $\square$

**3. Remaining Numbers of Zeros.** From Corollary 2.3 we know that  $H_{n,2,0}$ , the  $n \times n$  full upper Hessenberg pattern, is an indecomposable orthogonal pattern. Note that column  $i$  of  $H_{n,2,0}$  has exactly  $n - 1 - i$  zeros as long as  $1 \leq i \leq n - 1$ . We will need this fact in the proof of the next lemma.

LEMMA 3.1. *For  $n \geq 2$ , there exists an  $n \times n$  indecomposable orthogonal matrix with  $k$  zeros,  $k = \frac{1}{2}(n - 2)(n - 1), \dots, (n - 2)^2$ .*

*Proof.* We proceed by induction. Suppose that there exists an  $n \times n$  indecomposable orthogonal pattern  $P_k$  with exactly  $k$  zeros,  $k = \frac{1}{2}(n - 2)(n - 1), \dots, (n - 2)^2$ . Also suppose that  $P_k$  has a column, namely column  $j(k)$ , with exactly  $n - 2$  zeros. It is easily verified that these conditions hold for  $n = 2$ .

First note that we may take  $P_{\frac{1}{2}(n-2)(n-1)}$  to be  $H_{n,2,0}$ . Form  $D_i(H_{n,2,0})$ ,  $i = 1, \dots, n - 1$ . Now we count zeros.  $H_{n,2,0}$  has  $\frac{1}{2}(n - 2)(n - 1)$  zeros, we double a column with  $n - 1 - i$  zeros and we add  $n - 1$  zeros along the bottom of the pattern.

$$\begin{aligned} \#(D_i(H_{n,2,0})) &= (n - 1 - i) + (n - 1) + \#(H_{n,2,0}) \\ &= -i + (n - 1) + (n - 1) + \frac{1}{2}(n - 2)(n - 1) \\ &= -i + (n - 1) + \frac{1}{2}(n - 1)(n) \\ &= -i + ((n + 1) - 2) + \frac{1}{2}((n + 1) - 2)((n + 1) - 1). \end{aligned}$$

Since  $i$  ranges from 1 to  $n - 1$ ,  $\#(D_i(H_{n,2,0}))$  ranges from  $\frac{1}{2}((n + 1) - 2)((n + 1) - 1)$  to  $((n + 1) - 3) + \frac{1}{2}((n + 1) - 2)((n + 1) - 1)$ . Also note that the last row of  $D_i(H_{n,2,0})$  has  $(n + 1) - 2$  zeros so that  $(D_i(H_{n,2,0}))^T$  is an indecomposable orthogonal pattern with a column that has exactly  $(n + 1) - 2$  zeros,  $i = 1, \dots, n - 1$ .

Next, for each  $k = \frac{1}{2}(n - 2)(n - 1) + 1, \dots, (n - 2)^2$ , form  $D_{j(k)}(P_k)$ . Again we count zeros.  $P_k$  has  $k$  zeros, we double a column with  $n - 2$  zeros and we add  $n - 1$  zeros along the bottom of the pattern.

$$\#(D_{j(k)}(P_k)) = k + (n - 1) + (n - 2).$$

Since  $k$  ranges from  $\frac{1}{2}(n - 2)(n - 1) + 1$  up to  $(n - 2)^2$ , we have that  $\#(D_{j(k)}(P_k))$  ranges from

$$\begin{aligned} \frac{1}{2}(n - 2)(n - 1) + 1 + (n - 1) + (n - 2) &= \frac{1}{2}(n - 1)(n) + (n - 1) \\ &= \frac{1}{2}((n + 1) - 2)((n + 1) - 1) + ((n + 1) - 2) \end{aligned}$$

up to

$$\begin{aligned} (n - 2)^2 + (n - 1) + (n - 2) &= (n^2 - 4n + 4) + (n - 1) + (n - 2) \\ &= n^2 - 2n + 1 \\ &= (n - 1)^2 \\ &= ((n + 1) - 2)^2. \end{aligned}$$

Note that since  $D_{j(k)}(P_k)$  has a row with exactly  $(n+1) - 2$  zeros,  $(D_{j(k)}(P_k))^T$  is an indecomposable orthogonal pattern that has a column with exactly  $(n+1) - 2$  zeros.

Combining the two ranges of constructed  $(n+1) \times (n+1)$  indecomposable orthogonal patterns gives us matrices with numbers of zeros from  $\frac{1}{2}((n+1) - 2)((n+1) - 1)$  up to  $((n+1) - 2)^2$ . And since each of the transposes of these matrices has a column with exactly  $(n+1) - 2$  zeros, the induction is complete.  $\square$

**THEOREM 3.2.** *For  $n \geq 2$ , there is an  $n \times n$  indecomposable orthogonal matrix with exactly  $k$  zeros if and only if  $0 \leq k \leq (n - 2)^2$ .*

*Proof.* The theorem follows immediately from Corollary 2.4, Lemma 3.1 and the result of [BBS].  $\square$

**REMARK 3.3.** It follows from Theorem 3.2 that for  $n \geq 4$ , there exists an  $n \times n$  orthogonal matrix with exactly  $k$  zeros if and only if  $0 \leq k \leq n(n - 1) - 4$  or  $k = n(n - 1) - 2$  or  $k = n(n - 1)$ .

#### REFERENCES

- [BBS] L. B. Beasley, R. A. Brualdi and B. L. Shader, Combinatorial Orthogonality, in *Combinatorial and Graph-Theoretical Problems in Linear Algebra*, R. A. Brualdi, S. Friedland and V. Klee, eds., Springer-Verlag, New York, pp. 207-218, 1993.
- [C] R. Craigen, The craft of weaving matrices, *Congressus Numerantium*, 92:9-28, 1993.
- [F] M. Fiedler, A question raised about the sparsity of orthogonal matrices, Oral communication during the IMA Linear Algebra year, Minneapolis, Minn., 1991.