

2000

On convergence of infinite matrix products

Olga Holtz
holtz@cs.wisc.edu

Follow this and additional works at: <http://repository.uwyo.edu/ela>

Recommended Citation

Holtz, Olga. (2000), "On convergence of infinite matrix products", *Electronic Journal of Linear Algebra*, Volume 7.
DOI: <https://doi.org/10.13001/1081-3810.1054>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact scholcom@uwyo.edu.

ON CONVERGENCE OF INFINITE MATRIX PRODUCTS*

OLGA HOLTZ†

Abstract. A necessary and sufficient condition for the convergence of an infinite right product of matrices of the form

$$A := \begin{bmatrix} I & B \\ 0 & C \end{bmatrix},$$

with (uniformly) contracting submatrices C , is proven.

Key words. Infinite matrix products, RCP sets.

AMS subject classifications. 15A60, 15A99

1. Introduction. Consider the set of all matrices in $\mathbb{C}^{d \times d}$ of the form

$$(1) \quad A := \begin{bmatrix} I_s & B \\ 0 & C \end{bmatrix},$$

where I_s denotes the identity matrix of order $s < d$.

Matrices (1) are known to form an LCP set whenever the submatrices B are uniformly bounded and the submatrices C are uniformly contracting, that is, satisfy the condition $\|C\| \leq r$ for some fixed matrix (i.e., submultiplicative) norm $\|\cdot\|$ on $\mathbb{C}^{(d-s) \times (d-s)}$ and some constant $r < 1$; see, e.g., [1]. To recall, a set Σ has the LCP (RCP) property if all left (right) infinite products formed from matrices in Σ are convergent.

Matrices of the form (1) with uniformly bounded submatrices B and uniformly contracting submatrices C do not necessarily form an RCP set. (They do form such a set if and only if they satisfy a very stringent condition given in Corollary 2.3 below.) However, there exists a simple criterion that can be used to check whether a *particular* right infinite product formed from such matrices converges.

2. A convergence test.

THEOREM 2.1. *Let $(A_n)_{n \in \mathbb{N}}$ be a sequence of matrices of the form (1) and let*

$$\|C_n\| \leq r < 1 \quad \text{for all } n \in \mathbb{N}$$

for some matrix norm $\|\cdot\|$. The sequence $(P_n := A_1 A_2 \cdots A_n)$ converges if and only if so does the sequence $(B_n(I - C_n)^{-1})$. In this event,

$$\lim_{n \rightarrow \infty} P_n = \begin{bmatrix} I & \lim_{n \rightarrow \infty} B_n(I - C_n)^{-1} \\ 0 & 0 \end{bmatrix}.$$

*Received by the editors on 6 September 2000. Accepted for publication on 21 September 2000.
 Handling Editor: Daniel Hershkowitz.

† Department of Computer Sciences, University of Wisconsin, Madison, Wisconsin 53706 U.S.A.
 (holtz@cs.wisc.edu). This work was supported in part by the Clay Mathematics Institute.

Proof. To prove the necessity, partition P_n conformably with A_n . Then

$$P_n = \begin{bmatrix} I & X_n \\ 0 & C_1 C_2 \cdots C_n \end{bmatrix}, \quad \text{where} \quad X_n := \sum_{i=0}^n B_{n-i} (C_{n+1-i} C_{n+2-i} \cdots C_n).$$

If (P_n) converges, then $\lim_{n \rightarrow \infty} (X_n - X_{n-1}) = 0$. Also, $\|(I - C_n)^{-1}\| \leq 1/(1 - r)$ for all $n \in \mathbb{N}$. But $X_n = B_n + X_{n-1} C_n$, and thus

$$B_n (I - C_n)^{-1} - X_{n-1} = (X_n - X_{n-1}) (I - C_n)^{-1} \rightarrow 0 \quad \text{as} \quad n \rightarrow \infty.$$

Hence $\lim_{n \rightarrow \infty} B_n (I - C_n)^{-1} = \lim_{n \rightarrow \infty} X_n$.

To prove the sufficiency, without loss of generality one can assume that $s = d - s$. Indeed, simply replace each A_n by

$$\widetilde{A}_n := \begin{bmatrix} I_{\max\{s, d-s\}} & \widetilde{B}_n \\ 0 & \widetilde{C}_n \end{bmatrix},$$

where

$$\widetilde{B}_n := \begin{cases} \begin{bmatrix} B_n & 0_{s \times (2s-d)} \end{bmatrix} & \text{if } s \geq d - s \\ \begin{bmatrix} B_n \\ 0_{(d-2s) \times (d-s)} \end{bmatrix} & \text{if } s < d - s, \end{cases}$$

$$\widetilde{C}_n := \begin{cases} \begin{bmatrix} C_n & 0_{(d-s) \times (2s-d)} \\ 0_{(2s-d) \times (d-s)} & 0_{2s-d} \end{bmatrix} & \text{if } s \geq d - s \\ C_n & \text{if } s < d - s. \end{cases}$$

Then the matrices \widetilde{A}_n satisfy all the assumptions of the theorem and the sequence $(B_n (I - C_n)^{-1})$ (the product P_n) converges if and only if so does the sequence $(\widetilde{B}_n (I - \widetilde{C}_n)^{-1})$ (the product \widetilde{P}_n).

Thus, assume that $s = d - s$. Note that if the sequence $(B_n (I - C_n)^{-1})$ converges, then the sequence (B_n) is bounded, since $\|I - C_n\| \leq 1 + r$ for all n . Now, let

$$D_n := X_n - B_n (I - C_n)^{-1}$$

$$Y_n := B_{n+1} (I - C_{n+1})^{-1} - B_n (I - C_n)^{-1}$$

for all $n \in \mathbb{N}$. Then

$$(2) \quad D_{n+1} = (D_n - Y_n) C_{n+1},$$

hence

$$\|D_{n+1}\| \leq (\|D_n\| + \|Y_n\|) \|C_{n+1}\| \leq (\|D_n\| + \|Y_n\|) r.$$

Repeated use of this inequality gives

$$\|D_n\| \leq \sum_{i=1}^{n-1} \|Y_{n-i}\| r^i.$$

This implies, in particular, that

$$S := \limsup_{n \rightarrow \infty} \|D_n\| < \infty.$$

Since $\lim_{n \rightarrow \infty} Y_n = 0$, the identity (2) and the upper bound on $\|C_n\|$ imply that $S \leq rS$, therefore $S = 0$, that is, $\lim_{n \rightarrow \infty} D_n = 0$. \square

The obtained criterion of convergence can be used to make two more observations in the same spirit.

COROLLARY 2.2. *Let $(A_n)_{n \in \mathbb{N}}$ be a sequence of matrices of the form (1) such that the sequence (C_n) converges to a matrix C with spectral radius smaller than 1. Then the sequence $(P_n := A_1 A_2 \cdots A_n)$ converges if and only if so does the sequence (B_n) . In this event,*

$$\lim_{n \rightarrow \infty} P_n = \begin{bmatrix} I & \lim_{n \rightarrow \infty} B_n (I - C)^{-1} \\ 0 & 0 \end{bmatrix}.$$

Proof. If $\rho(C) < 1$, then there exists a matrix norm $\|\cdot\|$ on $\mathbb{C}^{(d-s) \times (d-s)}$ such that $\|C\| < 1$; see, e.g., [2, p. 297, Lemma 5.6.10]. Thus, $\|C_n\| \leq r$ for all $n \geq N$ for some $r < 1$ and some $N \in \mathbb{N}$, and the assumption of Theorem 2.1 is then satisfied. The product P_n converges whenever the product $A_N A_{N+1} \cdots$ converges, therefore (P_n) has a limit whenever (B_n) has one. On the other hand, the sequence $((I - C_n)^{-1})_{n=N}^\infty$ is bounded, so the necessity argument from the proof of Theorem 2.1 shows that the convergence of (B_n) is also necessary. \square

COROLLARY 2.3. *A set Σ consisting of matrices of the form (1) with uniformly contracting submatrices C is an RCP set if and only if*

$$(3) \quad B_1(I - C_1)^{-1} = B_2(I - C_2)^{-1} \quad \text{for all } A_1, A_2 \in \Sigma,$$

where

$$A_i = \begin{bmatrix} I & B_i \\ 0 & C_i \end{bmatrix}, \quad i = 1, 2.$$

Proof. Given $A_1, A_2 \in \Sigma$, apply Theorem 2.1 to the product $A_1 A_2 A_1 A_2 \cdots$ to see that the condition (3) is necessary and sufficient for the convergence of such a product. But if it is satisfied for all pairs of matrices from Σ , then it is sufficient for the convergence of any right product of matrices from Σ . \square



Acknowledgements. I am grateful to Professor Hans Schneider for his critical reading of the manuscript and to the referee for valuable suggestions.

REFERENCES

- [1] I. Daubechies and J. Lagarias. Sets of matrices all infinite products of which converge. *Linear Algebra Appl.*, 161:227–263, 1992.
- [2] R. A. Horn and C. R. Johnson. *Matrix analysis*. Cambridge University Press, Cambridge, 1985.