

2014

Simultaneous decomposition of two EP matrices with applications

Julio Benitez
jbenitez@mat.upv.es

Xiaoji Liu

Follow this and additional works at: <http://repository.uwyo.edu/ela>

Recommended Citation

Benitez, Julio and Liu, Xiaoji. (2014), "Simultaneous decomposition of two EP matrices with applications", *Electronic Journal of Linear Algebra*, Volume 27.
DOI: <https://doi.org/10.13001/1081-3810.1627>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact scholcom@uwyo.edu.

SIMULTANEOUS DECOMPOSITION OF TWO EP MATRICES WITH APPLICATIONS*

JULIO BENÍTEZ[†] AND XIAOJI LIU[‡]

Abstract. We find a simple canonical form for EP complex matrices \mathbf{A} and \mathbf{B} under simultaneous unitary equivalence.

Key words. EP matrices, Unitarily equivalence, Generalized inverses.

AMS subject classifications. 15B99, 15A23, 15A09.

1. Introduction. For an $m \times n$ complex matrix \mathbf{A} , the symbols \mathbf{A}^* , $\mathcal{R}(\mathbf{A})$, $\mathcal{N}(\mathbf{A})$, and $\text{rk}(\mathbf{A})$ will stand for the conjugate transpose, the column space, the null space, and the rank of \mathbf{A} , respectively. An $n \times n$ matrix \mathbf{P} is an orthogonal projector if $\mathbf{P} = \mathbf{P}^2 = \mathbf{P}^*$. The symbol \mathbf{I}_n will denote the identity matrix of order n . The zero matrix of order $n \times m$ will be denoted by $\mathbf{0}_{n,m}$, and $\mathbf{0}_n$ will be used instead of $\mathbf{0}_{n,n}$. When there is no danger of confusion with the size, a zero matrix will be denoted, simply, by $\mathbf{0}$. Furthermore, \mathbf{A}^\dagger will stand for the Moore-Penrose inverse of $\mathbf{A} \in \mathbb{C}_{n,m}$ i.e., the unique matrix satisfying the four equations

$$\mathbf{A}\mathbf{A}^\dagger\mathbf{A} = \mathbf{A}, \quad \mathbf{A}^\dagger\mathbf{A}\mathbf{A}^\dagger = \mathbf{A}^\dagger, \quad (\mathbf{A}\mathbf{A}^\dagger)^* = \mathbf{A}\mathbf{A}^\dagger, \quad (\mathbf{A}^\dagger\mathbf{A})^* = \mathbf{A}^\dagger\mathbf{A}.$$

It is known that any matrix $\mathbf{A} \in \mathbb{C}_{n,m}$ has a Moore-Penrose inverse (see e.g. [21]). Moreover, it can be easily proved that $\mathbf{A}\mathbf{A}^\dagger$ is the orthogonal projector onto $\mathcal{R}(\mathbf{A})$ and $\mathbf{A}^\dagger\mathbf{A}$ is the orthogonal projector onto $\mathcal{R}(\mathbf{A}^*)$.

The following lemma, which we will use below, establishes a canonical form for a pair of orthogonal projectors (see e.g., [8, 13, 16]).

LEMMA 1.1 (CS decomposition). *Let $\mathbf{P}_1, \mathbf{P}_2 \in \mathbb{C}_{n,n}$ be two orthogonal projectors.*

*Received by the editors on November 16, 2012. Accepted for publication on June 2, 2014.
Handling Editor: Bryan L. Shader.

[†]Universidad Politécnica de Valencia, Instituto de Matemática Multidisciplinar, Valencia, 46022, Spain (jbenitez@mat.upv.es).

[‡]Faculty of Science, Guangxi University for Nationalities, Nanning 530006, China, and Guangxi Key Laboratory of Hybrid Computational and IC Design Analysis, Nanning 530006, China (xiaojiлию72@126.com).

and

$$\mathbf{B} = \mathbf{U} \begin{bmatrix} \mathbf{CB}_1\mathbf{C} & \mathbf{CB}_1\mathbf{S} & \mathbf{CB}_2 & \mathbf{0} \\ \mathbf{S}^*\mathbf{B}_1\mathbf{C} & \mathbf{S}^*\mathbf{B}_1\mathbf{S} & \mathbf{S}^*\mathbf{B}_2 & \mathbf{0} \\ \mathbf{B}_3\mathbf{C} & \mathbf{B}_3\mathbf{S} & \mathbf{B}_4 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*, \quad (2.2)$$

where

- (i) $\mathbf{A}_1 \in \mathbb{C}_{r+s, r+s}$, $\mathbf{A}_5 \in \mathbb{C}_{q-s, q-s}$, $\mathbf{A}_9 \in \mathbb{C}_{p, p}$, $\mathbf{B}_1 \in \mathbb{C}_{r+s, r+s}$, $\mathbf{B}_2 \in \mathbb{C}_{r+s, p}$, $\mathbf{B}_3 \in \mathbb{C}_{p, r+s}$, and $\mathbf{B}_4 \in \mathbb{C}_{p, p}$.
- (ii) $\mathbf{C} = \text{diag}(\cos \theta_1, \dots, \cos \theta_r) \oplus \mathbf{0}_s$, $\mathbf{S} = \begin{bmatrix} \text{diag}(\sin \theta_1, \dots, \sin \theta_r) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_s & \mathbf{0}_{s, q-s} \end{bmatrix}$,
- (iii) $\begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \mathbf{A}_3 \\ \mathbf{A}_4 & \mathbf{A}_5 & \mathbf{A}_6 \\ \mathbf{A}_7 & \mathbf{A}_8 & \mathbf{A}_9 \end{bmatrix}$ and $\begin{bmatrix} \mathbf{B}_1 & \mathbf{B}_2 \\ \mathbf{B}_3 & \mathbf{B}_4 \end{bmatrix}$ are nonsingular.

Proof. By Lemma 1.1 applied to the orthogonal projectors $\mathbf{P}_\mathbf{A}$ and $\mathbf{P}_\mathbf{B}$, there exist $\theta_1, \dots, \theta_r \in]0, \pi/2[$, $p, q, r, s \in \{0, 1, \dots, n\}$, and a unitary matrix $\mathbf{U} \in \mathbb{C}_{n, n}$ such that

$$\mathbf{P}_\mathbf{A} = \mathbf{U} \left(\begin{bmatrix} \mathbf{I}_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \oplus \mathbf{I}_p \oplus \mathbf{I}_q \oplus \mathbf{0} \oplus \mathbf{0} \right) \mathbf{U}^*$$

and

$$\mathbf{P}_\mathbf{B} = \mathbf{U} \left(\begin{bmatrix} \widehat{\mathbf{C}}^2 & \widehat{\mathbf{C}}\widehat{\mathbf{S}} \\ \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \widehat{\mathbf{S}}^2 \end{bmatrix} \oplus \mathbf{I}_p \oplus \mathbf{0} \oplus \mathbf{I}_s \oplus \mathbf{0} \right) \mathbf{U}^*,$$

where $\widehat{\mathbf{C}} = \text{diag}(\cos \theta_1, \dots, \cos \theta_r)$ and $\widehat{\mathbf{S}} = \text{diag}(\sin \theta_1, \dots, \sin \theta_r)$. Evidently, we have

$$\widehat{\mathbf{C}}\widehat{\mathbf{S}} = \widehat{\mathbf{S}}\widehat{\mathbf{C}} \quad \text{and} \quad \widehat{\mathbf{C}}^2 + \widehat{\mathbf{S}}^2 = \mathbf{I}_r. \quad (2.3)$$

It is clear that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{P}_\mathbf{A}) = r + p + q$. Also, it is easy to see that the matrix $\begin{bmatrix} \widehat{\mathbf{C}} & -\widehat{\mathbf{S}} \\ \widehat{\mathbf{S}} & \widehat{\mathbf{C}} \end{bmatrix}$ is unitary and in particular is nonsingular. By using

$$\begin{bmatrix} \widehat{\mathbf{C}} & -\widehat{\mathbf{S}} \\ \widehat{\mathbf{S}} & \widehat{\mathbf{C}} \end{bmatrix} \begin{bmatrix} \widehat{\mathbf{C}} & \widehat{\mathbf{S}} \\ \mathbf{0} & \mathbf{0}_r \end{bmatrix} = \begin{bmatrix} \widehat{\mathbf{C}}^2 & \widehat{\mathbf{C}}\widehat{\mathbf{S}} \\ \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \widehat{\mathbf{S}}^2 \end{bmatrix}$$

and $\text{rk} \left(\begin{bmatrix} \widehat{\mathbf{C}} & \widehat{\mathbf{S}} \\ \mathbf{0} & \mathbf{0}_r \end{bmatrix} \right) = r$, we get $\text{rk}(\mathbf{B}) = \text{rk}(\mathbf{P}_\mathbf{B}) = r + p + s$. By the rank hypothesis, we obtain $s \leq q$. We shall denote $\mathbf{N} = [\mathbf{I}_s \mid \mathbf{0}_{s, q-s}] \in \mathbb{C}_{s, q}$,

$$\mathbf{C} = \begin{bmatrix} \widehat{\mathbf{C}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s \end{bmatrix} \in \mathbb{C}_{r+s, r+s} \quad \text{and} \quad \mathbf{S} = \begin{bmatrix} \widehat{\mathbf{S}} & \mathbf{0} \\ \mathbf{0} & \mathbf{N} \end{bmatrix} \in \mathbb{C}_{r+s, r+q}.$$

Evidently,

$$\begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{S}^*\mathbf{C} & \mathbf{S}^*\mathbf{S} \end{bmatrix} = \begin{bmatrix} \widehat{\mathbf{C}}^2 & \mathbf{0} & \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s & \mathbf{0} & \mathbf{0}_{s, q} \\ \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \mathbf{0} & \widehat{\mathbf{S}}^2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{q, s} & \mathbf{0} & \mathbf{N}^*\mathbf{N} \end{bmatrix}, \quad \mathbf{N}^*\mathbf{N} = \begin{bmatrix} \mathbf{I}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{q-s} \end{bmatrix}.$$

By means of a suitable permutation, we can assume that

$$\mathbf{P}_A = \mathbf{U} \left(\left[\begin{array}{cc} \mathbf{I}_{r+s} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+s} \oplus \mathbf{I}_{q-s} \end{array} \right] \oplus \mathbf{I}_p \oplus \mathbf{0} \right) \mathbf{U}^* \quad (2.4)$$

and

$$\mathbf{P}_B = \mathbf{U} \left(\left[\begin{array}{cc} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{S}^* \mathbf{C} & \mathbf{S}^* \mathbf{S} \end{array} \right] \oplus \mathbf{I}_p \oplus \mathbf{0} \right) \mathbf{U}^*. \quad (2.5)$$

Let $m = 2r + s + q$. Observe that the first summands in (2.4) and (2.5) are $m \times m$ matrices. We partition \mathbf{A} as follows:

$$\mathbf{A} = \mathbf{U} \left[\begin{array}{ccc} \mathbf{A}_1 & \mathbf{A}_2 & \mathbf{A}_3 \\ \mathbf{A}_4 & \mathbf{A}_5 & \mathbf{A}_6 \\ \mathbf{A}_7 & \mathbf{A}_8 & \mathbf{A}_9 \end{array} \right] \mathbf{U}^*, \quad \mathbf{A}_1 \in \mathbb{C}_{m,m}, \mathbf{A}_5 \in \mathbb{C}_{p,p}. \quad (2.6)$$

Since \mathbf{A} is EP, it follows $\mathbf{A} = \mathbf{A} \mathbf{A}^\dagger \mathbf{A} = \mathbf{P}_A \mathbf{A}$ and $\mathbf{A} = \mathbf{A} \mathbf{P}_A$. From (2.4) and (2.6) we obtain the blocks $\mathbf{A}_3, \mathbf{A}_6, \mathbf{A}_7, \mathbf{A}_8, \mathbf{A}_9$ are zero, and if we denote

$$\mathbf{P} = \left[\begin{array}{cc} \mathbf{I}_{r+s} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+s} \oplus \mathbf{I}_{q-s} \end{array} \right] \in \mathbb{C}_{m,m}, \quad (2.7)$$

then we get

$$\mathbf{A}_1 = \mathbf{P} \mathbf{A}_1 = \mathbf{A}_1 \mathbf{P}, \quad \mathbf{A}_2 = \mathbf{P} \mathbf{A}_2, \quad \mathbf{A}_4 = \mathbf{A}_4 \mathbf{P}.$$

By employing these equalities, $\mathbf{A}_1, \mathbf{A}_2$ and \mathbf{A}_4 can be written as

$$\mathbf{A}_1 = \left[\begin{array}{ccc} \mathbf{A}_{11} & \mathbf{0} & \mathbf{A}_{12} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_{21} & \mathbf{0} & \mathbf{A}_{22} \end{array} \right], \quad \mathbf{A}_2 = \left[\begin{array}{c} \mathbf{A}_{13} \\ \mathbf{0} \\ \mathbf{A}_{23} \end{array} \right], \quad \mathbf{A}_4 = \left[\begin{array}{ccc} \mathbf{A}_{31} & \mathbf{0} & \mathbf{A}_{32} \end{array} \right],$$

where $\mathbf{A}_{11} \in \mathbb{C}_{r+s,r+s}, \mathbf{A}_{22} \in \mathbb{C}_{q-s,q-s}, \mathbf{A}_{13} \in \mathbb{C}_{r+s,p}, \mathbf{A}_{23} \in \mathbb{C}_{q-s,p}, \mathbf{A}_{31} \in \mathbb{C}_{p,r+s}$, and $\mathbf{A}_{32} \in \mathbb{C}_{p,q-s}$. Thus, (by a renaming of the subindexes) \mathbf{A} can be written as in (2.1). Furthermore, the size of $\mathbf{\Delta} = \left[\begin{array}{ccc} \mathbf{A}_{11} & \mathbf{A}_{12} & \mathbf{A}_{13} \\ \mathbf{A}_{21} & \mathbf{A}_{22} & \mathbf{A}_{23} \\ \mathbf{A}_{31} & \mathbf{A}_{32} & \mathbf{A}_5 \end{array} \right]$ is $r + q + p$, which is equal to the rank of \mathbf{A} . Also it is evident that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{\Delta})$, which implies that $\mathbf{\Delta}$ is nonsingular.

We partition \mathbf{B} as follows:

$$\mathbf{B} = \mathbf{U} \left[\begin{array}{ccc} \mathbf{B}_1 & \mathbf{B}_2 & \mathbf{B}_3 \\ \mathbf{B}_4 & \mathbf{B}_5 & \mathbf{B}_6 \\ \mathbf{B}_7 & \mathbf{B}_8 & \mathbf{B}_9 \end{array} \right] \mathbf{U}^*, \quad \mathbf{B}_1 \in \mathbb{C}_{m,m}, \mathbf{B}_5 \in \mathbb{C}_{p,p}.$$

As for \mathbf{A} , the blocks $\mathbf{B}_3, \mathbf{B}_6, \mathbf{B}_7, \mathbf{B}_8$, and \mathbf{B}_9 are zero, and if we denote

$$\mathbf{Q} = \begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{S}^*\mathbf{C} & \mathbf{S}^*\mathbf{S} \end{bmatrix}, \quad (2.8)$$

then

$$\mathbf{B}_1 = \mathbf{QB}_1, \quad \mathbf{B}_1 = \mathbf{B}_1\mathbf{Q}, \quad \mathbf{B}_2 = \mathbf{QB}_2, \quad \mathbf{B}_4 = \mathbf{B}_4\mathbf{Q}. \quad (2.9)$$

Let us partition

$$\mathbf{B}_1 = \begin{bmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{bmatrix}, \quad \mathbf{B}_{11} \in \mathbb{C}_{r+s, r+s}.$$

From the first equality of (2.9), we obtain

$$\mathbf{B}_{11} = \mathbf{C}(\mathbf{CB}_{11} + \mathbf{SB}_{21}), \quad \mathbf{B}_{12} = \mathbf{C}(\mathbf{CB}_{12} + \mathbf{SB}_{22}), \quad \mathbf{B}_{21} = \mathbf{S}^*(\mathbf{CB}_{11} + \mathbf{SB}_{21}),$$

and

$$\mathbf{B}_{22} = \mathbf{S}^*(\mathbf{CB}_{12} + \mathbf{SB}_{22}).$$

If we denote $\mathbf{H}_1 = \mathbf{CB}_{11} + \mathbf{SB}_{21}$ and $\mathbf{H}_2 = \mathbf{CB}_{12} + \mathbf{SB}_{22}$, then $\mathbf{B}_1 = \begin{bmatrix} \mathbf{CH}_1 & \mathbf{CH}_2 \\ \mathbf{S}^*\mathbf{H}_1 & \mathbf{S}^*\mathbf{H}_2 \end{bmatrix}$. From the second equality of (2.9) we get $\mathbf{CH}_1 = \mathbf{C}(\mathbf{H}_1\mathbf{C} + \mathbf{H}_2\mathbf{S}^*)\mathbf{C}$, $\mathbf{CH}_2 = \mathbf{C}(\mathbf{H}_1\mathbf{C} + \mathbf{H}_2\mathbf{S}^*)\mathbf{S}$, $\mathbf{S}^*\mathbf{H}_1 = \mathbf{S}^*(\mathbf{H}_1\mathbf{C} + \mathbf{H}_2\mathbf{S}^*)\mathbf{C}$, and $\mathbf{S}^*\mathbf{H}_2 = \mathbf{S}^*(\mathbf{H}_1\mathbf{C} + \mathbf{H}_2\mathbf{S}^*)\mathbf{S}$. If we define $\mathbf{H} = \mathbf{H}_1\mathbf{C} + \mathbf{H}_2\mathbf{S}^*$, then \mathbf{B}_1 can be written as

$$\mathbf{B}_1 = \begin{bmatrix} \mathbf{CHC} & \mathbf{CHS} \\ \mathbf{S}^*\mathbf{HC} & \mathbf{S}^*\mathbf{HS} \end{bmatrix}, \quad \mathbf{H} \in \mathbb{C}_{r+s, r+s}.$$

In the same way, by using the third and fourth equalities of (2.9), there exist $\mathbf{E} \in \mathbb{C}_{r+s, p}$ and $\mathbf{F} \in \mathbb{C}_{p, r+s}$ such that

$$\mathbf{B}_2 = \begin{bmatrix} \mathbf{CE} \\ \mathbf{S}^*\mathbf{E} \end{bmatrix}, \quad \mathbf{B}_4 = \begin{bmatrix} \mathbf{FC} & \mathbf{FS} \end{bmatrix}.$$

Thus, by renaming the blocks, \mathbf{B} can be written as in (2.2).

To find the canonical angles between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$, we appeal to Definition 1.2. From (2.4), (2.5), (2.7) and (2.8), obviously,

$$\mathbf{P}_A\mathbf{P}_B = \mathbf{U}(\mathbf{PQ} \oplus \mathbf{I}_p \oplus \mathbf{0})\mathbf{U}^*, \quad (2.10)$$

and if we denote $\mathbf{X} = \mathbf{0}_{r+s} \oplus \mathbf{I}_{q-s}$, then

$$\mathbf{PQ} = \begin{bmatrix} \mathbf{I}_{r+s} & \mathbf{0} \\ \mathbf{0} & \mathbf{X} \end{bmatrix} \begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{S}^*\mathbf{C} & \mathbf{S}^*\mathbf{S} \end{bmatrix} = \begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{XS}^*\mathbf{C} & \mathbf{XS}^*\mathbf{S} \end{bmatrix}.$$

But we have

$$\mathbf{XS}^* = \begin{bmatrix} \mathbf{0}_r & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_{q-s} \end{bmatrix} \begin{bmatrix} \widehat{\mathbf{S}} & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_s \\ \mathbf{0} & \mathbf{0}_{q-s,s} \end{bmatrix} = \mathbf{0}.$$

Furthermore, from the definitions of \mathbf{C} and \mathbf{S} , and (2.3), we have

$$\mathbf{C}^2 + \mathbf{SS}^* = \begin{bmatrix} \widehat{\mathbf{C}}^2 & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s \end{bmatrix} + \begin{bmatrix} \widehat{\mathbf{S}}^2 & \mathbf{0} \\ \mathbf{0} & \mathbf{NN}^* \end{bmatrix} = \mathbf{I}_r \oplus \mathbf{I}_s = \mathbf{I}_{r+s},$$

$$\begin{aligned} \mathbf{S}(\mathbf{C} \oplus \mathbf{I}_{q-s}) &= \begin{bmatrix} \widehat{\mathbf{S}} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_s & \mathbf{0}_{s,q-s} \end{bmatrix} \begin{bmatrix} \widehat{\mathbf{C}} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_{q-s} \end{bmatrix} \\ &= \begin{bmatrix} \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s & \mathbf{0}_{s,q-s} \end{bmatrix} = \begin{bmatrix} \widehat{\mathbf{C}}\widehat{\mathbf{S}} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{s,q} \end{bmatrix} = \mathbf{CS}, \end{aligned}$$

and

$$\mathbf{S}^*\mathbf{S} + (\mathbf{C}^2 \oplus \mathbf{I}_{q-s}) = \begin{bmatrix} \widehat{\mathbf{S}}^2 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0}_{q-s} \end{bmatrix} + \begin{bmatrix} \widehat{\mathbf{C}}^2 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_{q-s} \end{bmatrix} = \mathbf{I}_{r+q},$$

which imply that $\begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \oplus \mathbf{I}_{q-s} \end{bmatrix}$ is unitary. From now on, we let

$$\mathbf{W} = \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \oplus \mathbf{I}_{q-s} \end{bmatrix}.$$

Hence, the fact that \mathbf{C} is diagonal and

$$\mathbf{PQ} = \begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{0} & \mathbf{0}_{r+q} \end{bmatrix} = \begin{bmatrix} \mathbf{C} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+q} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \oplus \mathbf{I}_{q-s} \end{bmatrix}$$

yield that $\cos\theta_1, \dots, \cos\theta_r$, and 0 (repeated $s+r+q$ times) are the singular values of \mathbf{PQ} . From (2.10), we obtain that $\cos\theta_1, \dots, \cos\theta_r$; 0 (repeated $s+r+q$ times); and 1 (repeated p times) are singular values of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B}$. Since there must be $\min\{\text{rk}(\mathbf{A}), \text{rk}(\mathbf{B})\} = \text{rk}(\mathbf{B}) = r+p+s$ canonical angles between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$, these canonical angles are 0 (repeated p times), $\theta_1, \dots, \theta_r$, and $\pi/2$ (repeated s times).

It remains to prove that (maintaining the notation of the proof) that $\mathbf{\Lambda} = \begin{bmatrix} \mathbf{H} & \mathbf{E} \\ \mathbf{F} & \mathbf{B}_5 \end{bmatrix}$ is nonsingular. To this end, since $\mathbf{\Lambda}$ is $(r+s+p) \times (r+s+p)$, it is sufficient to prove

$\text{rk}(\mathbf{A}) = r + s + p$. Observe that

$$\begin{aligned} \mathbf{U}^* \mathbf{B} \mathbf{U} &= \begin{bmatrix} \mathbf{C} \mathbf{H} \mathbf{C} & \mathbf{C} \mathbf{H} \mathbf{S} & \mathbf{C} \mathbf{E} & \mathbf{0} \\ \mathbf{S}^* \mathbf{H} \mathbf{C} & \mathbf{S}^* \mathbf{H} \mathbf{S} & \mathbf{S}^* \mathbf{E} & \mathbf{0} \\ \mathbf{F} \mathbf{C} & \mathbf{F} \mathbf{S} & \mathbf{B}_5 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{C} & -\mathbf{S} & \mathbf{0} & \mathbf{0} \\ \mathbf{S}^* & \mathbf{C} \oplus \mathbf{I}_{q-s} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_p & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{I} \end{bmatrix} \begin{bmatrix} \mathbf{H} & \mathbf{0} & \mathbf{E} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+q} & \mathbf{0} & \mathbf{0} \\ \mathbf{F} & \mathbf{0} & \mathbf{B}_5 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} & \mathbf{0} & \mathbf{0} \\ -\mathbf{S}^* & \mathbf{C} \oplus \mathbf{I}_{q-s} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{I}_p & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{I} \end{bmatrix}, \end{aligned}$$

where \mathbf{I} denotes an identity matrix of suitable size. In addition, since \mathbf{W} is unitary and in particular is nonsingular, we get $r + p + s = \text{rk}(\mathbf{B}) = \text{rk} \left(\begin{bmatrix} \mathbf{H} & \mathbf{E} \\ \mathbf{F} & \mathbf{B}_5 \end{bmatrix} \right)$. \square

Two particular cases are described in next two results.

COROLLARY 2.2. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices. The following statements are equivalent:*

- (i) $\mathbf{P}_\mathbf{A} \mathbf{P}_\mathbf{B} = \mathbf{P}_\mathbf{B} \mathbf{P}_\mathbf{A}$.
- (ii) *The matrices \mathbf{A} and \mathbf{B} can be written as*

$$\mathbf{A} = \mathbf{U} \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_3 & \mathbf{A}_4 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*, \quad \mathbf{B} = \mathbf{U} \begin{bmatrix} \mathbf{0}_q & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{B}_4 & \mathbf{B}_3 & \mathbf{0} \\ \mathbf{0} & \mathbf{B}_2 & \mathbf{B}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*, \quad (2.11)$$

where $\mathbf{U} \in \mathbb{C}_{n,n}$ is unitary, $\mathbf{A}_1 \in \mathbb{C}_{q,q}$, $\mathbf{A}_4, \mathbf{B}_4 \in \mathbb{C}_{p,p}$, $\mathbf{B}_1 \in \mathbb{C}_{s,s}$, $\begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \\ \mathbf{A}_3 & \mathbf{A}_4 \end{bmatrix}$ and $\begin{bmatrix} \mathbf{B}_4 & \mathbf{B}_3 \\ \mathbf{B}_2 & \mathbf{B}_1 \end{bmatrix}$ are nonsingular.

Proof. The proof of (ii) \Rightarrow (i) is trivial. Let us prove the converse. We can clearly assume that $\text{rk}(\mathbf{B}) \geq \text{rk}(\mathbf{A})$. Since $\mathbf{P}_\mathbf{A} \mathbf{P}_\mathbf{B}$ is an orthogonal projector, its singular values are 0 or 1. Hence, there is no canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$ in $]0, \pi/2[$. By Theorem 2.1, we can write matrices \mathbf{A} and \mathbf{B} as in (2.1) and (2.2) with $r = 0$, and therefore, $\mathbf{C} = \mathbf{0}_s$ and $\mathbf{S} = [\mathbf{I}_s \ \mathbf{0}_{s,q-s}]$. Evidently, we get

$$\mathbf{S}^* \mathbf{B}_1 \mathbf{S} = \begin{bmatrix} \mathbf{B}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{q-s} \end{bmatrix}, \quad \mathbf{B}_3 \mathbf{S} = [\mathbf{B}_3 \ \mathbf{0}_{p,q-s}], \quad \mathbf{S}^* \mathbf{B}_2 = \begin{bmatrix} \mathbf{B}_2 \\ \mathbf{0}_{q-s,p} \end{bmatrix}.$$

By a simultaneous permutation of the rows and the columns of \mathbf{A} and \mathbf{B} , we get

$$\mathbf{A} = \mathbf{U} \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 & \mathbf{A}_3 & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_4 & \mathbf{A}_5 & \mathbf{A}_6 & \mathbf{0} & \mathbf{0} \\ \mathbf{A}_7 & \mathbf{A}_8 & \mathbf{A}_9 & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0}_s & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*, \quad \mathbf{B} = \mathbf{U} \begin{bmatrix} \mathbf{0}_s & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{q-s} & \mathbf{0} & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{B}_4 & \mathbf{B}_3 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{B}_2 & \mathbf{B}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*.$$

By joining some blocks of the matrices \mathbf{A} and \mathbf{B} , and renaming the blocks of these matrices, we obtain the theorem. \square

Based on Theorem 2.1, we now give a simpler proof of Corollary 2.3 than that given in [4, Corollary 3.9]

COROLLARY 2.3. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices. The following statements are equivalent:*

- (i) $\mathbf{AB} = \mathbf{BA}$.
- (ii) *The matrices \mathbf{A} and \mathbf{B} can be written as*

$$\mathbf{A} = \mathbf{U}(\mathbf{A}_1 \oplus \mathbf{A}_2 \oplus \mathbf{0} \oplus \mathbf{0})\mathbf{U}^*, \quad \mathbf{B} = \mathbf{U}(\mathbf{0} \oplus \mathbf{B}_1 \oplus \mathbf{B}_2 \oplus \mathbf{0})\mathbf{U}^*,$$

where $\mathbf{U} \in \mathbb{C}_{n,n}$ is unitary, $\mathbf{A}_1 \in \mathbb{C}_{q,q}$, $\mathbf{A}_2, \mathbf{B}_1 \in \mathbb{C}_{p,p}$, and $\mathbf{B}_2 \in \mathbb{C}_{s,s}$ are nonsingular matrices such that $\mathbf{A}_1\mathbf{B}_1 = \mathbf{B}_1\mathbf{A}_1$.

Proof. We only prove (i) \Rightarrow (ii) because the other implication is trivial. Since \mathbf{A} is EP, there exists a unitary matrix \mathbf{V} such that $\mathbf{A} = \mathbf{V}(\mathbf{K} \oplus \mathbf{0})\mathbf{V}^*$, where $\mathbf{K} \in \mathbb{C}_{k,k}$ is nonsingular (see e.g. [9, Section 4.3]). From $\mathbf{AB} = \mathbf{BA}$ we get that \mathbf{B} can be written as $\mathbf{B} = \mathbf{V}(\mathbf{X} \oplus \mathbf{Y})\mathbf{V}^*$ with $\mathbf{X} \in \mathbb{C}_{k,k}$. Since $\mathbf{P}_\mathbf{A} = \mathbf{V}(\mathbf{I}_k \oplus \mathbf{0})\mathbf{V}^*$ and $\mathbf{P}_\mathbf{B} = \mathbf{V}(\mathbf{X}\mathbf{X}^\dagger \oplus \mathbf{Y}\mathbf{Y}^\dagger)\mathbf{V}^*$ we obtain $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} = \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$. Corollary 2.2 yields that matrices \mathbf{A} and \mathbf{B} can be written as in (2.11). From $\mathbf{AB} = \mathbf{BA}$ we get $\mathbf{A}_2\mathbf{B}_4 = \mathbf{0}$, $\mathbf{A}_2\mathbf{B}_3 = \mathbf{0}$, $\mathbf{A}_4\mathbf{B}_3 = \mathbf{0}$, $\mathbf{B}_4\mathbf{A}_3 = \mathbf{0}$, $\mathbf{B}_2\mathbf{A}_3 = \mathbf{0}$, and $\mathbf{B}_2\mathbf{A}_4 = \mathbf{0}$. From these, we get

$$\begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \\ \mathbf{A}_3 & \mathbf{A}_4 \end{bmatrix} \begin{bmatrix} \mathbf{0} \\ \mathbf{B}_3 \end{bmatrix} = \begin{bmatrix} \mathbf{B}_4 & \mathbf{B}_3 \\ \mathbf{B}_2 & \mathbf{B}_1 \end{bmatrix} \begin{bmatrix} \mathbf{A}_3 \\ \mathbf{0} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \end{bmatrix}$$

and

$$\begin{bmatrix} \mathbf{0} & \mathbf{B}_2 \end{bmatrix} \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \\ \mathbf{A}_3 & \mathbf{A}_4 \end{bmatrix} = \begin{bmatrix} \mathbf{A}_2 & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{B}_4 & \mathbf{B}_3 \\ \mathbf{B}_2 & \mathbf{B}_1 \end{bmatrix} = \begin{bmatrix} \mathbf{0} & \mathbf{0} \end{bmatrix}.$$

The nonsingularity of $\begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \\ \mathbf{A}_3 & \mathbf{A}_4 \end{bmatrix}$ and $\begin{bmatrix} \mathbf{B}_4 & \mathbf{B}_3 \\ \mathbf{B}_2 & \mathbf{B}_1 \end{bmatrix}$ imply that \mathbf{A}_2 , \mathbf{A}_3 , \mathbf{B}_2 , and \mathbf{B}_3 are null. By a renaming blocks, the proof is concluded. \square

3. Some applications. In this section, we obtain several results based on Theorem 2.1.

The following lemma concerns representations based on Theorem 2.1 of two EP matrices and their Moore-Penrose inverses under certain conditions and will be used extensively in the computations below.

LEMMA 3.1. *Let \mathbf{A} and $\mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices such that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular. Then $\mathbf{C} = \text{diag}(\cos \theta_1, \dots, \cos \theta_r) \oplus \mathbf{0}_s$, $\mathbf{S} =$*

$\text{diag}(\sin \theta_1, \dots, \sin \theta_r) \oplus \mathbf{I}_s,$

$$\mathbf{A} = \mathbf{U} \begin{bmatrix} \mathbf{A}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+s} \end{bmatrix} \mathbf{U}^*, \quad \mathbf{B} = \mathbf{U} \begin{bmatrix} \mathbf{CB}_1\mathbf{C} & \mathbf{CB}_1\mathbf{S} \\ \mathbf{S}^*\mathbf{B}_1\mathbf{C} & \mathbf{S}^*\mathbf{B}_1\mathbf{S} \end{bmatrix} \mathbf{U}^*, \quad (3.1)$$

and

$$\mathbf{A}^\dagger = \mathbf{U} \begin{bmatrix} \mathbf{A}_1^{-1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_{r+s} \end{bmatrix} \mathbf{U}^*, \quad \mathbf{B}^\dagger = \mathbf{U} \begin{bmatrix} \mathbf{CB}_1^{-1}\mathbf{C} & \mathbf{CB}_1^{-1}\mathbf{S} \\ \mathbf{S}^*\mathbf{B}_1^{-1}\mathbf{C} & \mathbf{S}^*\mathbf{B}_1^{-1}\mathbf{S} \end{bmatrix} \mathbf{U}^*. \quad (3.2)$$

Proof. We write \mathbf{A} and \mathbf{B} as in Theorem 2.1. Since $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular, $2r + q + s = n$ and $p = 0$. Since $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$, then $q = s$. Therefore, \mathbf{C} , \mathbf{S} , \mathbf{A} , and \mathbf{B} can be expressed as in the statement of the theorem. The representation of \mathbf{A}^\dagger is evident. Since $\begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix}$ is unitary,

$$\begin{aligned} \mathbf{B}^\dagger &= \left(\mathbf{U} \begin{bmatrix} \mathbf{C} & -\mathbf{S} \\ \mathbf{S}^* & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{B}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix} \mathbf{U}^* \right)^\dagger \\ &= \mathbf{U} \begin{bmatrix} \mathbf{C} & -\mathbf{S} \\ \mathbf{S}^* & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{B}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}^\dagger \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix} \mathbf{U}^* \\ &= \mathbf{U} \begin{bmatrix} \mathbf{C} & -\mathbf{S} \\ \mathbf{S}^* & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{B}_1^{-1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix} \mathbf{U}^* \\ &= \mathbf{U} \begin{bmatrix} \mathbf{CB}_1^{-1}\mathbf{C} & \mathbf{CB}_1^{-1}\mathbf{S} \\ \mathbf{S}^*\mathbf{B}_1^{-1}\mathbf{C} & \mathbf{S}^*\mathbf{B}_1^{-1}\mathbf{S} \end{bmatrix} \mathbf{U}^*. \end{aligned} \quad (3.3)$$

The proof is finished. \square

Let \mathbf{A} and \mathbf{B} be two EP matrices, and let p, r, s and q have the same meaning as in Theorem 2.1. If $\text{rk}(\mathbf{A}) \leq \text{rk}(\mathbf{B})$, then

$$\text{rk}(\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}) = \text{rk}(\mathbf{A}) + \text{rk}(\mathbf{B}) - 2p = 2r + q + s \quad (3.4)$$

and

$$\text{rk}(\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}) = 2r. \quad (3.5)$$

The reader is referred to [11] for a deeper insight of a pair of orthogonal projectors, concretely, the equalities (3.4) and (3.5) appeared as part of Theorem 26 and Corollary 55 in [11]. In particular, we have that the nonsingularity of $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ implies $p = 0$; and the nonsingularity of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ implies $q = s = p = 0$, $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B}) = r$. Hence, the nonsingularity of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ yields the nonsingularity of $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ (observe that we can drop $\text{rk}(\mathbf{A}) \leq \text{rk}(\mathbf{B})$). In next Corollary 3.2 we will give a kind of converse. To establish this converse, we need the concept of co-EP matrices, which

will be immediately defined. Following [6], we say that $\mathbf{X} \in \mathbb{C}_{n,n}$ is *co-EP* when $\mathbf{X}\mathbf{X}^\dagger - \mathbf{X}^\dagger\mathbf{X}$ is nonsingular.

COROLLARY 3.2. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices such that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular. Then the following statements are equivalent:*

- (i) $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ is nonsingular.
- (ii) $\mathbf{AB} - \mathbf{BA}$ is nonsingular.
- (iii) \mathbf{AB} is *co-EP*.

Under this equivalence one has $\mathbf{AB}(\mathbf{AB})^\dagger = \mathbf{P}_\mathbf{A}$ and $(\mathbf{AB})^\dagger\mathbf{AB} = \mathbf{P}_\mathbf{B}$.

Proof. Maintaining the notation of Theorem 2.1, the hypotheses of the corollary yield $q = s$ and $2r + q + s = n$.

(i) \Leftrightarrow (ii): By (3.1), we have

$$\mathbf{AB} - \mathbf{BA} = \mathbf{U} \begin{bmatrix} \mathbf{A}_1\mathbf{CB}_1\mathbf{C} - \mathbf{CB}_1\mathbf{CA}_1 & \mathbf{A}_1\mathbf{CB}_1\mathbf{S} \\ -\mathbf{S}^*\mathbf{B}_1\mathbf{CA}_1 & \mathbf{0} \end{bmatrix} \mathbf{U}^*.$$

Hence, $\det(\mathbf{AB} - \mathbf{BA}) = \det(\mathbf{A}_1)^2 \det(\mathbf{B}_1)^2 \det(\mathbf{C})^2 \det(\mathbf{S})^2$. Recall that Theorem 2.1 implies that \mathbf{A}_1 and \mathbf{B}_1 are both nonsingular. Lemma 3.1 yields that \mathbf{S} is nonsingular. Hence, $\mathbf{AB} - \mathbf{BA}$ is nonsingular $\Leftrightarrow \mathbf{C}$ is nonsingular $\Leftrightarrow s = 0 \Leftrightarrow 2r = n \Leftrightarrow \mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ is nonsingular.

(i) \Rightarrow (iii): If we set $\mathbf{X} = \mathbf{A}_1\mathbf{CB}_1$, then (3.1) yields

$$\mathbf{AB} = \mathbf{U} \begin{bmatrix} \mathbf{XC} & \mathbf{XS} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{X} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix} \mathbf{U}^*. \quad (3.6)$$

Since $\begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix}$ is unitary, it follows

$$(\mathbf{AB})^\dagger = \mathbf{U} \begin{bmatrix} \mathbf{C} & -\mathbf{S} \\ \mathbf{S}^* & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{X}^\dagger & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{CX}^\dagger & \mathbf{0} \\ \mathbf{S}^*\mathbf{X}^\dagger & \mathbf{0} \end{bmatrix} \mathbf{U}^*. \quad (3.7)$$

Since $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ is nonsingular, $s = 0$. Hence, \mathbf{C} is nonsingular, and from the definition of matrix \mathbf{X} , we can see that \mathbf{X} is nonsingular. Equalities (2.4) and (2.5), and the above representations of \mathbf{AB} and $(\mathbf{AB})^\dagger$ lead to

$$\mathbf{AB}(\mathbf{AB})^\dagger = \mathbf{U} \begin{bmatrix} \mathbf{I}_r & \mathbf{0} \\ \mathbf{0} & \mathbf{0}_r \end{bmatrix} \mathbf{U}^* = \mathbf{P}_\mathbf{A}, \quad (\mathbf{AB})^\dagger\mathbf{AB} = \mathbf{U} \begin{bmatrix} \mathbf{C}^2 & \mathbf{CS} \\ \mathbf{S}^*\mathbf{C} & \mathbf{S}^*\mathbf{S} \end{bmatrix} \mathbf{U}^* = \mathbf{P}_\mathbf{B}.$$

Thus, $\mathbf{AB}(\mathbf{AB})^\dagger - (\mathbf{AB})^\dagger\mathbf{AB} = \mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular, i.e., \mathbf{AB} is *co-EP*.

(iii) \Rightarrow (i): Assume that \mathbf{AB} is *co-EP*. Theorem 2.3 of [6] implies that $(\mathbf{AB}) + (\mathbf{AB})^*$ is nonsingular. From (3.1) we have $(\mathbf{AB}) + (\mathbf{AB})^* = \mathbf{U} \begin{bmatrix} \mathbf{XC} + \mathbf{CX}^* & \mathbf{XS} \\ \mathbf{S}^*\mathbf{X}^* & \mathbf{0} \end{bmatrix} \mathbf{U}^*$, where $\mathbf{X} = \mathbf{A}_1\mathbf{CB}_1$. The nonsingularity of $(\mathbf{AB}) + (\mathbf{AB})^*$ yields the nonsingularity

of \mathbf{XS} , which implies that \mathbf{X} is nonsingular. Hence, \mathbf{C} is nonsingular, which implies that $\mathbf{P}_A\mathbf{P}_B - \mathbf{P}_B\mathbf{P}_A$ is nonsingular. \square

A square matrix \mathbf{A} is said to be *group invertible* if there exists a matrix \mathbf{X} such that

$$\mathbf{AXA} = \mathbf{A}, \quad \mathbf{XAX} = \mathbf{X}, \quad \mathbf{XA} = \mathbf{AX}. \quad (3.8)$$

It can be proved (see e.g. [3, Chapter 4]) that for a square matrix \mathbf{A} there is at most one matrix \mathbf{X} satisfying (3.8). Such matrix will be denoted by $\mathbf{A}^\#$. We will use the following result due to Cline (see [10] or [3, Section 4.4]).

THEOREM 3.3. *Let a square matrix \mathbf{A} have the full-rank factorization $\mathbf{A} = \mathbf{FG}$. Then \mathbf{A} has a group inverse if and only if \mathbf{GF} is nonsingular, in which case $\mathbf{A}^\# = \mathbf{F}(\mathbf{GF})^{-2}\mathbf{G}$.*

COROLLARY 3.4. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be EP matrices such that $\mathbf{P}_A\mathbf{P}_B - \mathbf{P}_B\mathbf{P}_A$ is nonsingular. Then \mathbf{AB} is group invertible.*

Proof. By representing \mathbf{A} and \mathbf{B} as in (2.1), (2.2) and recalling that the nonsingularity of $\mathbf{P}_A\mathbf{P}_B - \mathbf{P}_B\mathbf{P}_A$ implies $q = s = p = 0$, in particular, the nonsingularity of $\mathbf{P}_A - \mathbf{P}_B$ and $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$, we obtain from Lemma 3.1 that

$$\mathbf{AB} = \mathbf{U} \begin{bmatrix} \mathbf{XC} & \mathbf{XS} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{X} \\ \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \end{bmatrix} \mathbf{U}^*, \quad (3.9)$$

where $\mathbf{X} = \mathbf{A}_1\mathbf{CB}_1 \in \mathbb{C}_{r,r}$. We have $\text{rk}(\mathbf{AB}) = \text{rk}(\mathbf{XC}) = r$ because \mathbf{X} and \mathbf{C} are nonsingular. We let $\mathbf{F} = \mathbf{U} \begin{bmatrix} \mathbf{X} \\ \mathbf{0} \end{bmatrix} \in \mathbb{C}_{n,r}$ and $\mathbf{G} = \begin{bmatrix} \mathbf{C} & \mathbf{S} \end{bmatrix} \mathbf{U}^* \in \mathbb{C}_{r,n}$. Evidently $\text{rk}(\mathbf{F}) = \text{rk}(\mathbf{X}) = r$ because \mathbf{X} is nonsingular. Furthermore, $r = \text{rk}(\mathbf{I}_r) = \text{rk}(\mathbf{GG}^*) = \text{rk}(\mathbf{G})$. Hence, (3.9) is a full-rank factorization of \mathbf{AB} . From the nonsingularity of \mathbf{CX} and Theorem 3.3, we get that \mathbf{AB} is group invertible. \square

Furthermore, Theorem 3.3 allows us to give a representation of $(\mathbf{AB})^\#$ under the hypotheses of Corollary 3.4.

$$(\mathbf{AB})^\# = \mathbf{U} \begin{bmatrix} \mathbf{X} \\ \mathbf{0} \end{bmatrix} (\mathbf{CX})^{-2} \begin{bmatrix} \mathbf{C} & \mathbf{S} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{C}^{-1}\mathbf{X}^{-1} & \mathbf{C}^{-1}\mathbf{X}^{-1}\mathbf{C}^{-1}\mathbf{S} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*, \quad (3.10)$$

where $\mathbf{X} = \mathbf{A}_1\mathbf{CB}_1$.

EXAMPLE 1. The converse of Corollary 3.4 does not hold in general, as the following example shows. Let $\mathbf{A} = \text{diag}(1, 0)$ and $\mathbf{B} = \text{diag}(0, 1)$. Evidently, \mathbf{A} and \mathbf{B} are two orthogonal projectors and thus, $\mathbf{A} = \mathbf{A}^\dagger = \mathbf{P}_A$ and $\mathbf{B} = \mathbf{B}^\dagger = \mathbf{P}_B$. However, $\mathbf{AB} = \mathbf{0}$ is group invertible and $\mathbf{P}_A\mathbf{P}_B - \mathbf{P}_B\mathbf{P}_A$ is singular.

Evidently, we have $\|\mathbf{AB}\|_2 \leq \|\mathbf{A}\|_2\|\mathbf{B}\|_2$ for any pair of conformable matrices \mathbf{A} and \mathbf{B} . We have a sharper bound provided $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ are EP, $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$, and

$\mathbf{P}_A - \mathbf{P}_B$ is nonsingular. Furthermore, we find some bounds for the norm of several expressions.

COROLLARY 3.5. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be EP matrices and $\theta_1 < \dots < \theta_r$ be the canonical angles between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$.*

(i) *If $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and $\mathbf{P}_A - \mathbf{P}_B$ is nonsingular, then*

$$\|\mathbf{AB}\|_2 \leq \|\mathbf{A}\|_2 \|\mathbf{B}\|_2 \cos \theta_1.$$

(ii) *If $\mathbf{P}_A \mathbf{P}_B - \mathbf{P}_B \mathbf{P}_A$ is nonsingular, then*

(a) $\|(\mathbf{AB})^\dagger\|_2 \leq \|\mathbf{A}^\dagger\|_2 \|\mathbf{B}^\dagger\|_2 / \cos \theta_r.$

(b) $\frac{\sin^2 \theta_1}{\cos \theta_1} \frac{\|\mathbf{B}^\dagger\|_2}{\|\mathbf{A}\|_2} \leq \|(\mathbf{AB})^\dagger - \mathbf{B}^\dagger \mathbf{A}^\dagger\|_2 \leq \|\mathbf{A}^\dagger\|_2 \|\mathbf{B}^\dagger\|_2 \frac{\sin^2 \theta_r}{\cos \theta_r}.$

(c) $\|(\mathbf{AB})^\# \|_2 \leq \|\mathbf{A}^\dagger\|_2 \|\mathbf{B}^\dagger\|_2 / \cos^3 \theta_r.$

Proof. Let us represent \mathbf{A} and \mathbf{B} as in (3.1) and let us define $\mathbf{X} = \mathbf{A}_1 \mathbf{C} \mathbf{B}_1$.

(i): By (3.1), (3.3) and (3.6), we have

$$\|\mathbf{AB}\|_2 = \|\mathbf{X}\|_2 = \|\mathbf{A}_1 \mathbf{C} \mathbf{B}_1\|_2 \leq \|\mathbf{A}_1\|_2 \|\mathbf{C}\|_2 \|\mathbf{B}_1\|_2 = \|\mathbf{A}\|_2 \|\mathbf{B}\|_2 \|\mathbf{C}\|_2.$$

Observe that $\|\mathbf{C}\|_2 = \max\{\cos \theta_i : 1 \leq i \leq r\}$.

(ii): Since $\mathbf{P}_A \mathbf{P}_B - \mathbf{P}_B \mathbf{P}_A$ is nonsingular, \mathbf{C} is nonsingular. From (3.2), (3.3) and (3.7), we have

$$\begin{aligned} \|(\mathbf{AB})^\dagger\|_2 &= \|\mathbf{X}^{-1}\|_2 = \|\mathbf{B}_1^{-1} \mathbf{C}^{-1} \mathbf{A}_1^{-1}\|_2 \\ &\leq \|\mathbf{B}_1^{-1}\|_2 \|\mathbf{C}^{-1}\|_2 \|\mathbf{A}_1^{-1}\|_2 = \|\mathbf{A}^\dagger\|_2 \|\mathbf{B}^\dagger\|_2 \|\mathbf{C}^{-1}\|_2. \end{aligned}$$

To prove (a), observe that $\|\mathbf{C}^{-1}\|_2 = \max\{(\cos \theta_i)^{-1} : 1 \leq i \leq r\}$.

In order to prove (b), define $\mathbf{Y} = \mathbf{B}_1^{-1} \mathbf{C} \mathbf{A}_1^{-1}$. From (3.2) and (3.7) we have

$$\begin{aligned} (\mathbf{AB})^\dagger - \mathbf{B}^\dagger \mathbf{A}^\dagger &= \mathbf{U} \begin{bmatrix} \mathbf{C}(\mathbf{X}^{-1} - \mathbf{Y}) & \mathbf{0} \\ \mathbf{S}^*(\mathbf{X}^{-1} - \mathbf{Y}) & \mathbf{0} \end{bmatrix} \mathbf{U}^* \\ &= \mathbf{U} \begin{bmatrix} \mathbf{C} & -\mathbf{S} \\ \mathbf{S}^* & \mathbf{C} \end{bmatrix} \begin{bmatrix} \mathbf{X}^{-1} - \mathbf{Y} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*. \end{aligned}$$

Thus, $\|(\mathbf{AB})^\dagger - \mathbf{B}^\dagger \mathbf{A}^\dagger\|_2 = \|\mathbf{X}^{-1} - \mathbf{Y}\|_2$. But, $\mathbf{X}^{-1} - \mathbf{Y} = \mathbf{B}_1^{-1}(\mathbf{C}^{-1} - \mathbf{C})\mathbf{A}_1^{-1}$. Therefore,

$$\begin{aligned} \|(\mathbf{AB})^\dagger - \mathbf{B}^\dagger \mathbf{A}^\dagger\|_2 &= \|\mathbf{B}_1^{-1}(\mathbf{C}^{-1} - \mathbf{C})\mathbf{A}_1^{-1}\|_2 \\ &\leq \|\mathbf{B}_1^{-1}\|_2 \|\mathbf{C}^{-1} - \mathbf{C}\|_2 \|\mathbf{A}_1^{-1}\|_2 = \|\mathbf{A}^\dagger\|_2 \|\mathbf{B}^\dagger\|_2 \|\mathbf{C}^{-1} - \mathbf{C}\|_2. \end{aligned}$$

By considering the function $f :]0, \pi/2[\rightarrow \mathbb{R}$ given by $f(\theta) = \frac{1}{\cos \theta} - \cos \theta$, we have $\|\mathbf{C}^{-1} - \mathbf{C}\|_2 = \frac{1}{\cos \theta_r} - \cos \theta_r$. In addition, observe that $\mathbf{C}^{-1} - \mathbf{C} = \mathbf{C}^{-1}(\mathbf{I}_r - \mathbf{C}^2) =$

$\mathbf{C}^{-1}\mathbf{S}^2$. Thus,

$$\begin{aligned}\|\mathbf{B}^\dagger\|_2 &= \|\mathbf{B}_1^{-1}\|_2 = \|\mathbf{B}_1^{-1}(\mathbf{C}^{-1} - \mathbf{C})\mathbf{A}_1^{-1}\mathbf{A}_1\mathbf{C}\mathbf{S}^{-2}\|_2 \\ &\leq \|(\mathbf{A}\mathbf{B})^\dagger - \mathbf{B}^\dagger\mathbf{A}^\dagger\|_2\|\mathbf{A}\|_2\|\mathbf{S}^{-2}\mathbf{C}\|_2.\end{aligned}$$

By studying the function $g(\theta) = \cos\theta/\sin^2\theta$, we have $\|\mathbf{S}^{-2}\mathbf{C}\|_2 = \cos\theta_1/\sin^2\theta_1$. Thus, (b) is proved.

Recall that by Corollary 3.4, the matrix $\mathbf{A}\mathbf{B}$ is group invertible. By (3.10) and if we denote $\mathbf{Z} = \mathbf{C}^{-1}\mathbf{B}_1^{-1}\mathbf{C}^{-1}\mathbf{A}_1^{-1}\mathbf{C}^{-1}$ we have

$$(\mathbf{A}\mathbf{B})^\# = \mathbf{U} \begin{bmatrix} \mathbf{Z}\mathbf{C} & \mathbf{Z}\mathbf{S} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{Z} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S}^* & \mathbf{C} \end{bmatrix} \mathbf{U}^*.$$

Hence, $\|(\mathbf{A}\mathbf{B})^\#\|_2 = \|\mathbf{Z}\|_2 \leq \|\mathbf{C}^{-1}\|_2^3\|\mathbf{B}^\dagger\|_2\|\mathbf{A}^\dagger\|_2$. Thus, (c) is proved. \square

EXAMPLE 2. This example shows that the bounds established in Corollary 3.5 cannot be improved. Let

$$\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Since $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B}) = 1$, according to Definition 1.2 there is only one canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$. Since \mathbf{A} and \mathbf{B} are orthogonal projectors, $\mathbf{A}^\dagger = \mathbf{A}$, $\mathbf{B}^\dagger = \mathbf{B}$, and $\|\mathbf{A}\|_2 = \|\mathbf{B}\|_2 = 1$. In addition, \mathbf{A} and \mathbf{B} are EP matrices, $\mathbf{P}_\mathbf{A} = \mathbf{A}$, and $\mathbf{P}_\mathbf{B} = \mathbf{B}$. Now,

$$\mathbf{A}\mathbf{B} = \mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad \mathbf{B}\mathbf{A} = \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A} = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}.$$

Let us note that \mathbf{A} and \mathbf{B} satisfy any condition of Corollary 3.5.

The singular values of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B}$ are 0 and $\sqrt{2}/2$ (because the singular values of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B}$ are the square root of the eigenvalues of $(\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B})(\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B})^*$). Hence, the unique canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$ is $\pi/4$. Also,

$$\|\mathbf{A}\mathbf{B}\|_2 = \max\{\sigma : \sigma \text{ is a singular value of } \mathbf{A}\mathbf{B}\} = \sqrt{2}/2.$$

Hence, the bound of Corollary 3.5 (i) cannot be improved.

Now, $(\mathbf{A}\mathbf{B})^\dagger = (\mathbf{A}\mathbf{B})^* [(\mathbf{A}\mathbf{B})(\mathbf{A}\mathbf{B})^*]^\dagger = \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$. Hence, $\|(\mathbf{A}\mathbf{B})^\dagger\|_2 = \sqrt{2}$.

Thus, the bound of Corollary 3.5 (ii.a) cannot be improved.

Since $(\mathbf{A}\mathbf{B})^\dagger - \mathbf{B}^\dagger\mathbf{A}^\dagger = \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}$, we have $\|(\mathbf{A}\mathbf{B})^\dagger - \mathbf{B}^\dagger\mathbf{A}^\dagger\|_2 = \sqrt{2}/2$. The bounds of Corollary 3.5 (ii.b) cannot be improved.

Since $2\mathbf{AB}$ is an idempotent, $(2\mathbf{AB})^\# = 2\mathbf{AB}$. Hence, $(\mathbf{AB})^\# = 4\mathbf{AB}$, and therefore, $\|(\mathbf{AB})^\#\|_2 = 4\|\mathbf{AB}\|_2 = 2\sqrt{2}$, which shows that the bound of Corollary 3.5 (ii.c) cannot be improved.

EXAMPLE 3. By means of numerical experiments, we show how good is the bound of Corollary 3.5 (i). In the following m-file (that can be executed in Matlab, Octave, or Freemat), we compute $\|\mathbf{AB}\|_2 / (\|\mathbf{A}\|_2 \|\mathbf{B}\|_2 \cos \theta_1)$ for 29 pairs of 4×4 matrices \mathbf{A} and \mathbf{B} randomly chosen.

```
function example
x = zeros(1,29);
for i=1:29
    % find two EP 4x4 random matrices according Lemma 3.1
    aux1 = rand(2,2); aux2 = rand(2,2);
    A1 = aux1*aux1'+eye(2); B1 = aux2*aux2'+eye(2);
    % XX^* + I is always nonsingular
    A = [A1 zeros(2,2); zeros(2,2) zeros(2,2)];
    th = pi/2*rand(2,1); %two random angles in [0,pi/2]
    C = diag(cos(th)); S = diag(sin(th));
    B = [C*B1*C C*B1*S; S'*B1*C S'*B1*S];
    PA = A*pinv(A); PB = B*pinv(B); sigma = svd(PA*PB);
    % There are two canonical angles
    s1 = sigma(1); s2 = sigma(2);
    % cos(theta1) = s1; cos(theta2) = s2
    x(i) = norm(A*B)/(norm(A)*norm(B)*s1);
end
bar(1:29,x,0.2)
```

One execution of this file produces Figure 3.1. We can see that in general $\|\mathbf{AB}\|_2 \neq \|\mathbf{A}\|_2 \|\mathbf{B}\|_2 \cos \theta_1$.

We can perform analogous numerical experiments to test the remaining bounds. We can insert the following lines in the above code.

```
pab = pinv(A*B); pa = pinv(A); pb = pinv(b);
(norm(pab)*s2)/(norm(pa)*norm(pb))
(norm(pab-pb*pa)*s2)/(norm(pa)*norm(pb)*(1-s2^2))
(norm(pab-pb*pa)*s1*norm(A))/(norm(pb)*(1-s1^2))
giAB = pinv(pab*(A*B)^3*pab);
(norm(giAB)*s2^3)/(norm(pa)*norm(pb))
```

Observe that we have used $(\mathbf{AB})^\# = ((\mathbf{AB})^\dagger(\mathbf{AB})^3(\mathbf{AB})^\dagger)^\dagger$ (see [7, Remark 1]).

From Definition 1.2, it is evident that for a pair of orthogonal projectors \mathbf{P}_1 and




FIG. 3.1.

\mathbf{P}_2 , one has $\|\mathbf{P}_1\mathbf{P}_2\|_2 = \cos \theta_1$, where θ_1 is the least canonical angle between $\mathcal{R}(\mathbf{P}_1)$ and $\mathcal{R}(\mathbf{P}_2)$, which shows again that the bound in Corollary 3.5 (i) cannot be improved. Another related known equality is the following: $\|(\mathbf{P}_1\mathbf{P}_2)^\dagger\| = 1/\cos \theta_r$, where θ_r is the greatest canonical angle between $\mathcal{R}(\mathbf{P}_1)$ and $\mathcal{R}(\mathbf{P}_2)$ (see [11, Lemma 36]).

The right inequality of Corollary 3.5 (ii.b) is related with the reverse order law for the Moore-Penrose inverse. There are known characterizations of the reverse order law (see [1] and [14]). We get that under the hypotheses of Theorem 3.5, if $\theta_r = 0$, then $(\mathbf{AB})^\dagger = \mathbf{B}^\dagger\mathbf{A}^\dagger$.

The nonsingularity of linear combinations of two matrices has been a widely studied topic, see e.g. [2, 5, 12, 15, 18, 19, 20, 22, 23]. Based on Theorem 2.1, we deduce a result concerning this topic.

COROLLARY 3.6. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be EP matrices. If θ is not a canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$; and $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$, then $\mathcal{N}(a\mathbf{A} + b\mathbf{B}) = \mathcal{N}(\mathbf{A}) \cap \mathcal{N}(\mathbf{B})$ for any a, b nonzero complex numbers.*

Proof. Let us represent \mathbf{A} and \mathbf{B} as in Theorem 2.1. Since $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and 0

is not a canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$, then $s = q$ and $p = 0$. We have

$$a\mathbf{A} + b\mathbf{B} = \mathbf{U} \begin{bmatrix} a\mathbf{A}_1 + b\mathbf{CB}_1\mathbf{C} & b\mathbf{CB}_1\mathbf{S} & \mathbf{0} \\ b\mathbf{S}^*\mathbf{B}_1\mathbf{C} & b\mathbf{S}^*\mathbf{B}_1\mathbf{S} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^*.$$

Pick $\mathbf{x} \in \mathcal{N}(a\mathbf{A} + b\mathbf{B})$. By writing $\mathbf{x} = \mathbf{U} \begin{bmatrix} \mathbf{u} \\ \mathbf{v} \\ \mathbf{w} \end{bmatrix}$, where $\mathbf{u}, \mathbf{v} \in \mathbb{C}_{r+s,1}$, we get

$$(a\mathbf{A}_1 + b\mathbf{CB}_1\mathbf{C})\mathbf{u} + b\mathbf{CB}_1\mathbf{S}\mathbf{v} = \mathbf{0} \quad \text{and} \quad b\mathbf{S}^*\mathbf{B}_1\mathbf{C}\mathbf{u} + b\mathbf{S}^*\mathbf{B}_1\mathbf{S}\mathbf{v} = \mathbf{0}. \quad (3.11)$$

Since $b \neq 0$ and \mathbf{S} is nonsingular, the last of the equalities of (3.11) yields

$$\mathbf{B}_1\mathbf{C}\mathbf{u} + \mathbf{B}_1\mathbf{S}\mathbf{v} = \mathbf{0}. \quad (3.12)$$

By substitution of (3.12) into the first of the equalities of (3.11) we get $a\mathbf{A}_1\mathbf{u} = \mathbf{0}$, hence $\mathbf{u} = \mathbf{0}$ in view of $a \neq 0$ and the nonsingularity of \mathbf{A}_1 . Substituting $\mathbf{u} = \mathbf{0}$ into (3.12), we have $\mathbf{v} = \mathbf{0}$ (recall that \mathbf{B}_1 and \mathbf{S} are nonsingular). From $\mathbf{u} = \mathbf{0}$ and $\mathbf{v} = \mathbf{0}$ it is trivial to obtain $\mathbf{A}\mathbf{x} = \mathbf{B}\mathbf{x} = \mathbf{0}$, and thus, $\mathcal{N}(a\mathbf{A} + b\mathbf{B}) \subset \mathcal{N}(\mathbf{A}) \cap \mathcal{N}(\mathbf{B})$. The opposite inclusion is trivial. \square

COROLLARY 3.7. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices such that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular. Then $a\mathbf{A} + b\mathbf{B}$ is nonsingular for all $a, b \in \mathbb{C} \setminus \{0\}$.*

Proof. Since $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular, then $\mathcal{R}(\mathbf{P}_\mathbf{A}) \oplus \mathcal{R}(\mathbf{P}_\mathbf{B}) = \mathbb{C}^n$ (see [17, Theorem 4.1]). From Corollary 3.6 and by using that $\mathcal{R}(\mathbf{P}_\mathbf{A}) = \mathcal{R}(\mathbf{A}) = \mathcal{R}(\mathbf{A}^*)$, $\mathcal{R}(\mathbf{P}_\mathbf{B}) = \mathcal{R}(\mathbf{B}) = \mathcal{R}(\mathbf{B}^*)$ (because \mathbf{A} and \mathbf{B} are EP matrices), we have

$$[\mathcal{N}(a\mathbf{A} + b\mathbf{B})]^\perp = [\mathcal{N}(\mathbf{A}) \cap \mathcal{N}(\mathbf{B})]^\perp = \mathcal{N}(\mathbf{A})^\perp + \mathcal{N}(\mathbf{B})^\perp = \mathcal{R}(\mathbf{A}^*) + \mathcal{R}(\mathbf{B}^*) = \mathbb{C}^n.$$

Thus, $\mathcal{N}(a\mathbf{A} + b\mathbf{B}) = \{\mathbf{0}\}$. \square

In the next results, we give a representation and the norm of the projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{N}(\mathbf{B})$ and the projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{N}(\mathbf{A})$.

COROLLARY 3.8. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be two EP matrices such that $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ is nonsingular. Then $(\mathbf{A}\mathbf{B})(\mathbf{A}\mathbf{B})^\#$ is the oblique projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{N}(\mathbf{B})$ and $\|(\mathbf{A}\mathbf{B})(\mathbf{A}\mathbf{B})^\#\|_2 = 1/\cos\theta_r$, where θ_r is the greatest canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$.*

Proof. Recall that the nonsingularity of $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ implies that $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular and $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$. We represent the matrices \mathbf{A} and \mathbf{B} as in (3.1). Corollary 3.2 yields that matrix \mathbf{C} is nonsingular, and thus the representation (3.1) leads to $\text{rk}(\mathbf{A}\mathbf{B}) = \text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$. From the obvious $\mathcal{R}(\mathbf{A}\mathbf{B}) \subseteq \mathcal{R}(\mathbf{A})$ and $\mathcal{N}(\mathbf{B}) \subseteq \mathcal{N}(\mathbf{A}\mathbf{B})$ we obtain, respectively, $\mathcal{R}(\mathbf{A}\mathbf{B}) = \mathcal{R}(\mathbf{A})$ and $\mathcal{N}(\mathbf{B}) = \mathcal{N}(\mathbf{A}\mathbf{B})$. By recalling that $(\mathbf{A}\mathbf{B})(\mathbf{A}\mathbf{B})^\#$ is the oblique projector onto $\mathcal{N}(\mathbf{A}\mathbf{B})$ along $\mathcal{N}(\mathbf{A}\mathbf{B})$, we finish the proof of the first part of the corollary.

From (3.6) and (3.10), it is easy to get

$$\mathbf{AB}(\mathbf{AB})^\# = \mathbf{U} \begin{bmatrix} \mathbf{I}_r & \mathbf{C}^{-1}\mathbf{S} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* = \mathbf{U} \begin{bmatrix} \mathbf{C}^{-1} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \begin{bmatrix} \mathbf{C} & \mathbf{S} \\ -\mathbf{S} & \mathbf{C} \end{bmatrix} \mathbf{U}^*.$$

Thus, $\|\mathbf{AB}(\mathbf{AB})^\#\|_2 = \|\mathbf{C}^{-1}\|_2 = 1/\cos\theta_r$. \square

COROLLARY 3.9. *Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be EP matrices such that $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{B})$ and $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular. Then the projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{R}(\mathbf{B})$ is $\mathbf{A}(\mathbf{A} + \mathbf{B})^{-1}$. The norm of this projector equals to $1/\sin\theta_1$, where θ_1 is the smallest canonical angle between $\mathcal{R}(\mathbf{A})$ and $\mathcal{R}(\mathbf{B})$.*

Proof. Let us represent \mathbf{A} and \mathbf{B} as in (3.1). Observe that $\mathbf{S} = \mathbf{S}^*$ since \mathbf{S} is real and diagonal. The matrix \mathbf{P} defined as

$$\mathbf{P} = \mathbf{U} \begin{bmatrix} \mathbf{I}_{r+s} & -\mathbf{CS}^{-1} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{U}^* \quad (3.13)$$

obviously satisfies

$$\mathbf{P}^2 = \mathbf{P}, \quad \mathbf{PA} = \mathbf{A}, \quad \mathbf{PB} = \mathbf{0}, \quad \text{rk}(\mathbf{P}) = r + s. \quad (3.14)$$

The second equality of (3.14) yields $\mathcal{R}(\mathbf{A}) \subset \mathcal{R}(\mathbf{P})$. Since $\text{rk}(\mathbf{A}) = \text{rk}(\mathbf{P})$, we deduce $\mathcal{R}(\mathbf{A}) = \mathcal{R}(\mathbf{P})$. The third equality of (3.14) implies $\mathcal{R}(\mathbf{B}) \subset \mathcal{N}(\mathbf{P})$. Since $\dim\mathcal{N}(\mathbf{P}) = n - \text{rk}(\mathbf{P}) = r + s = \text{rk}(\mathbf{B})$ we deduce $\mathcal{R}(\mathbf{B}) = \mathcal{N}(\mathbf{P})$. Thus, the matrix \mathbf{P} defined in (3.13) is the oblique projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{R}(\mathbf{B})$. From (3.14) we get $\mathbf{P}(\mathbf{A} + \mathbf{B}) = \mathbf{A}$, and from Corollary 3.7 we obtain $\mathbf{P} = \mathbf{A}(\mathbf{A} + \mathbf{B})^{-1}$. From (3.13) we have $\|\mathbf{P}\|_2^2 = \|\mathbf{PP}^*\|_2 = \|\mathbf{I}_{r+s} + \mathbf{C}^2\mathbf{S}^{-2}\|_2$, but $\mathbf{I}_{r+s} + \mathbf{C}^2\mathbf{S}^{-2} = \mathbf{S}^{-2}(\mathbf{S}^2 + \mathbf{C}^2) = \mathbf{S}^{-2}$. By using $\|\mathbf{S}^{-2}\|_2 = 1/\sin^2\theta_1$, the proof is finished. Let us notice that by using [17, Theorem 3.1] we can prove also the assertion on the norm. \square

The expressions concerning the norm of the oblique projectors appearing in Corollaries 3.8 and 3.9 are not only conceptually simple, but, as illustrated in Figure 3.2, there is also a particularly nice picture that accompanies it.

Let $\mathbf{A}, \mathbf{B} \in \mathbb{C}_{n,n}$ be EP matrices such that $\mathbf{P}_\mathbf{A}\mathbf{P}_\mathbf{B} - \mathbf{P}_\mathbf{B}\mathbf{P}_\mathbf{A}$ is nonsingular. As we have proved, $\mathbf{P}_\mathbf{A} - \mathbf{P}_\mathbf{B}$ is nonsingular. Let $\mathbf{P} = \mathbf{A}(\mathbf{A} + \mathbf{B})^{-1}$ be the oblique projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{R}(\mathbf{B})$ and $\mathbf{Q} = \mathbf{AB}(\mathbf{AB})^\#$ be the oblique projector onto $\mathcal{R}(\mathbf{A})$ along $\mathcal{N}(\mathbf{B})$. In Figure 3.2, \mathbf{x}_1 and \mathbf{x}_2 are vectors of the unit ball that maximize $\|\mathbf{P}\mathbf{x}\|$ and $\|\mathbf{Q}\mathbf{x}\|$, respectively, when $\|\mathbf{x}\| \leq 1$, and thus, $\|\mathbf{P}\| = \|\mathbf{P}\mathbf{x}_1\|$ and $\|\mathbf{Q}\| = \|\mathbf{Q}\mathbf{x}_2\|$. But, as we can see in the right triangle of Figure 3.2, we have $\sin\theta = 1/\|\mathbf{P}\mathbf{x}_1\|$. In addition, from the left triangle of Figure 3.2, we get $\cos\theta = 1/\|\mathbf{Q}\mathbf{x}_2\|$.




FIG. 3.2.

Acknowledgment. The authors are grateful to the referee for reviewing the manuscript and also for his/her valuable comments. The first author is supported by Project UPV. PAID-06-12 (Ref. SP2012047). The second author is supported by the Guangxi Natural Science Foundation (2013GXNSFAA019008), the Key Project of Education Department of Guangxi (201202ZD031), National Science Foundation of China (11361009), and Science Research Project 2013 of the China-ASEAN Study Center (Guangxi Science Experiment Center) of Guangxi University for Nationalities.

REFERENCES

- [1] E. Arghiriade. Sur les matrices qui sont permutables avec leur inverse généralisée. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.*, 35:244–251, 1963.
- [2] J.K. Baksalary and O.M. Baksalary. Nonsingularity of linear combinations of idempotent matrices. *Linear Algebra Appl.*, 388:25–29, 2004.
- [3] A. Ben-Israel and T.N.E. Greville. *Generalized Inverses: Theory and Applications*, second edition. Springer, New York, 2002.
- [4] J. Benítez. Moore-Penrose inverses and commuting elements of C^* -algebras. *J. Math. Anal. Appl.*, 345:766–770, 2008.
- [5] J. Benítez, X. Liu, and T. Zhu. Nonsingularity and group invertibility of linear combinations of two k -potent matrices. *Linear Multilinear Algebra*, 58:1023–1035, 2010.
- [6] J. Benítez and V. Rakočević. Matrices A such that $AA^\dagger - A^\dagger A$ are nonsingular. *Appl. Math. Comput.*, 217:3493–3503, 2010.
- [7] J. Benítez and X. Liu. On the continuity of the group inverse. *Oper. Matrices*, 6:859–868, 2012.
- [8] Å. Björck. *Numerical Methods for Least Squares Problems*. SIAM, Philadelphia, 1996.
- [9] S.L. Campbell and C.D. Meyer. *Generalized Inverses of Linear Transformations*. SIAM, Philadelphia, 2009.
- [10] R.E. Cline. Inverses of rank invariant powers of a matrix. *SIAM J. Numer. Anal.*, 5:182–197, 1968.

- [11] A. Galántai. Subspaces, angles and pairs of orthogonal projections. *Linear Multilinear Algebra* 56:227–260, 2006.
- [12] H.L. Gau, C.J. Wang, and N.C. Wong. Invertibility and fredholmness of linear combinations of quadratic, k potent and nilpotent operators. *Oper. Matrices*, 2:193–199, 2008.
- [13] G.H. Golub and C.F. Van Loan. *Matrix Computations*, third edition. Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, 1996.
- [14] T.N.E. Greville. Note on the generalized inverse of a matrix product. *J. Soc. Indust. Appl. Math.*, 9:109–115, 1966.
- [15] J. Groß and G. Trenkler. Nonsingularity of the difference of two oblique projectors. *SIAM J. Matrix Anal. Appl.*, 21:390–395, 1999.
- [16] M. Hegland, J. Garcke, and V. Challis. The combination technique and some generalisations. *Linear Algebra Appl.*, 420:249–275, 2007.
- [17] I.C.F. Ipsen and C.D. Meyer. The angle between complementary subspaces. *Amer. Math. Monthly*, 102:904–911, 1995.
- [18] J.J. Koliha and V. Rakočević. Invertibility of the sum of idempotents. *Linear Multilinear Algebra*, 50:285–292, 2002.
- [19] J.J. Koliha and V. Rakočević. Invertibility of the difference of idempotents. *Linear Multilinear Algebra*, 51:97–110, 2003.
- [20] X. Liu, S. Wu, and J. Benítez. On nonsingularity of combinations of two group invertible matrices and two tripotent matrices. *Linear Multilinear Algebra*, 59:1409–1417, 2011.
- [21] R. Penrose. A generalized inverse for matrices. *Proc. Cambridge Phil. Soc.*, 51:406–413, 1955.
- [22] M. Tošić and D.S. Cvetković-Ilić. Invertibility of a linear combination of two matrices and partial orderings. *Appl. Math. Comp.*, 218:4651–4657, 2012.
- [23] M. Tošić, D.S. Cvetković-Ilić, and C. Deng. The Moore-Penrose inverse of a linear combination of commuting generalized and hypergeneralized projectors. *Electron. J. Linear Algebra*, 22:1129–1137, 2011.
- [24] H.K. Wimmer. Canonical angles of unitary spaces and perturbations of direct complements. *Linear Algebra Appl.*, 287:373–379, 1999.