

2013

Minimum number of distinct eigenvalues of graphs

Bahman Ahmadi

Fatemeh Alinaghipour

Michael S. Cavers

Shaun Fallat
shaun.fallat@uregina.ca

Karen Meagher

See next page for additional authors

Follow this and additional works at: <http://repository.uwyo.edu/ela>

Recommended Citation

Ahmadi, Bahman; Alinaghipour, Fatemeh; Cavers, Michael S.; Fallat, Shaun; Meagher, Karen; and Nasser, Shahla. (2013), "Minimum number of distinct eigenvalues of graphs", *Electronic Journal of Linear Algebra*, Volume 26.
DOI: <https://doi.org/10.13001/1081-3810.1679>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact scholcom@uwyo.edu.

Minimum number of distinct eigenvalues of graphs

Authors

Bahman Ahmadi, Fatemeh Alinaghipour, Michael S. Cavers, Shaun Fallat, Karen Meagher, and Shahla Nasser

MINIMUM NUMBER OF DISTINCT EIGENVALUES OF GRAPHS*

BAHMAN AHMADI[†], FATEMEH ALINAGHIPOUR[†], MICHAEL S. CAVERS[‡], SHAUN
FALLAT[§], KAREN MEAGHER[¶], AND SHAHLA NASSERASR^{||}

Abstract. The minimum number of distinct eigenvalues, taken over all real symmetric matrices compatible with a given graph G , is denoted by $q(G)$. Using other parameters related to G , bounds for $q(G)$ are proven and then applied to deduce further properties of $q(G)$. It is shown that there is a great number of graphs G for which $q(G) = 2$. For some families of graphs, such as the join of a graph with itself, complete bipartite graphs, and cycles, this minimum value is obtained. Moreover, examples of graphs G are provided to show that adding and deleting edges or vertices can dramatically change the value of $q(G)$. Finally, the set of graphs G with $q(G)$ near the number of vertices is shown to be a subset of known families of graphs with small maximum multiplicity.

Key words. Symmetric matrix, Eigenvalue, Join of graphs, Diameter, Trees, Bipartite graph, Maximum multiplicity.

AMS subject classifications. 05C50, 15A18.

1. Introduction. Suppose $G = (V, E)$ is a simple graph with vertex set $V = \{1, 2, \dots, n\}$ and edge set E . To a graph G , we associate the collection of real $n \times n$ symmetric matrices defined by

$$S(G) = \{A : A = A^T, \text{ for } i \neq j, a_{ij} \neq 0 \Leftrightarrow \{i, j\} \in E\}.$$

Note that, the main diagonal entries of A in $S(G)$ are free to be chosen.

For a square matrix A , we let $q(A)$ denote the number of distinct eigenvalues of A . For a graph G , we define

$$q(G) = \min\{q(A) : A \in S(G)\}.$$

*Received by the editors on April 2, 2013. Accepted for publication on September 7, 2013.
Handling Editor: Xingzhi Zhan.

[†]Department of Mathematics and Statistics, University of Regina, Regina, Saskatchewan, S4S 0A2, Canada (bahman.ahmadi@gmail.com, fatemeh.naghipour@gmail.com).

[‡]Department of Mathematics and Statistics, University of Calgary, Calgary, AB, T2N 1N4, Canada (michael.cavers@ucalgary.ca).

[§]Department of Mathematics and Statistics, University of Regina, Regina, Saskatchewan, S4S 0A2, Canada (shaun.fallat@uregina.ca). Research supported in part by an NSERC research grant.

[¶]Department of Mathematics and Statistics, University of Regina, Regina, Saskatchewan, S4S 0A2, Canada (karen.meagher@uregina.ca). Research supported in part by an NSERC research grant.

^{||}Department of Mathematics and Statistics, University of Regina, Regina, Saskatchewan, S4S 0A2, Canada (Shahla.Nasserarsr@uregina.ca). Research supported by PIMS and Fallat's NSERC research grant.

It is clear that for any graph G on n vertices, $1 \leq q(G) \leq n$. Furthermore, it is not difficult to show that for a fixed n , there exists a graph G on n vertices with $q(G) = k$, for each $k = 1, 2, \dots, n$, see Corollary 3.6 for further details.

The class of matrices $S(G)$ has been of interest to many researchers recently (see [7, 8] and the references therein), and there has been considerable development of the parameters $M(G)$ (maximum multiplicity or nullity over $S(G)$) and $\text{mr}(G)$ (minimum rank over $S(G)$) and their positive semidefinite counterparts, see, for example, the works [3, 7, 8]. Furthermore, as a consequence interest has grown in connecting these parameters to various combinatorial properties of G . For example, the inverse eigenvalue problem for graphs (see [13]) continues to receive considerable and deserved attention, as it remains one of the most interesting unresolved issues in combinatorial matrix theory.

In the context of the $(0, 1)$ -adjacency matrix, $A(G)$, it is well known that $q(A(G))$ is at least one more than the diameter of G (denoted by $\text{diam}(G)$) (see [4]). This result was generalized to the case of trees, by observing that $q(A) \geq \text{diam}(G) + 1$, whenever $A \in S(G)$ is an entry-wise nonnegative matrix (see [6]). Thus, if G is a tree, it is known that $q(G) \geq \text{diam}(G) + 1$. However, it has been demonstrated that while this inequality is tight for some trees (e.g. path, star), equality need not hold for all trees (see [2], [17], and also [19]).

Our main interest lies in studying the value of $q(G)$ for arbitrary graphs, and as such is a continuation of [12] by de Fonseca. As with many studies of this kind, moving beyond trees leads to a number of new and interesting difficulties, and numerous exciting advances. It is clear that knowledge of $q(G)$ for a graph G will impact current studies of the inverse eigenvalue problem for graphs, and, in particular, the parameters $M(G)$ and $\text{mr}(G)$.

Our work has been organized into a number of components. The next section contains necessary background information and various preliminary-type results including connections between $q(G)$ and existing graph parameters as well as the graphs attaining the extreme values of $q(G)$. The following section provides a simple but surprisingly useful lower bound for $q(G)$. Section 4 is devoted to studying the graphs for which $q(G) = 2$, which is continued into Section 5, whereas the next section considers bipartite graphs and certain graph products. The final two sections focus on the graphs for which $q(G) = |V(G)| - 1$ and some possible further work.

2. Preliminary results. To begin we list some basic results about the minimum number of distinct eigenvalues for graphs. In this work, we use $K_n, K_{m,n}$ and I_n to denote the complete graph on n vertices, complete bipartite graph with parts of sizes m, n , and the identity matrix of order n , respectively. The notations $M_{m,n}, M_n$ are used for the set of real matrices of order $m \times n$ and n , respectively. For $A \in M_n$, the

set of eigenvalues of A is denoted by $\sigma(A)$. For graphs G and H , $G \cup H$ denotes the graph with vertex set $V(G) \cup V(H)$ and edges $E(G) \cup E(H)$, and is called the *union of G and H* .

LEMMA 2.1. *For a graph G , $q(G) = 1$ if and only if G has no edges.*

Proof. If $q(G) = 1$, then there is an $A \in S(G)$ with exactly one eigenvalue, this matrix is a scalar multiple of the identity matrix, thus the graph G is the empty graph. Clearly, if G is the empty graph, then $q(G) = 1$. \square

LEMMA 2.2. *For any $n \geq 2$, we have $q(K_n) = 2$.*

Proof. The adjacency matrix of K_n has two distinct eigenvalues, so $q(K_n) \leq 2$ and by Lemma 2.1, $q(K_n) > 1$, which implies $q(K_n) = 2$. \square

LEMMA 2.3. *If G is a non-empty graph, then for any two distinct real numbers μ_1, μ_2 , there is an $A \in S(G)$ such that $q(A) = q(G)$ and $\mu_1, \mu_2 \in \sigma(A)$.*

Proof. Consider $B \in S(G)$ with $q(B) = q(G)$ and $\lambda_1, \lambda_2 \in \sigma(B)$ with $\lambda_1 \neq \lambda_2$. Then the matrix A defined below satisfies $q(A) = q(B)$ and $\mu_1, \mu_2 \in \sigma(A)$.

$$A = \frac{\mu_1 - \mu_2}{\lambda_1 - \lambda_2} \left(B + \frac{\lambda_1 \mu_2 - \lambda_2 \mu_1}{\mu_1 - \mu_2} I \right). \quad \square$$

COROLLARY 2.4. *If G and H are non-empty graphs with $q(G) = q(H) = 2$, then $q(G \cup H) = 2$. In particular, if G is the union of non-trivial complete graphs, then $q(G) = 2$.*

The parameter q is related to other parameters of graphs, such as the minimum rank of the graph.

PROPOSITION 2.5. *For any graph G , we have $q(G) \leq \text{mr}(G) + 1$.*

Proof. Consider a matrix $A \in S(G)$ with the minimum possible rank, $\text{mr}(G)$. Then, A has $\text{mr}(G)$ nonzero eigenvalues, so the number of distinct eigenvalues of A is less than or equal to $\text{mr}(G) + 1$. \square

Clearly, any known upper bound on the minimum rank of a graph can be used as an upper bound for the value of $q(G)$ for a graph G . For example, a *clique covering* of a graph is a collection of complete subgraphs of the graph such that every edge of the graph is contained in at least one of these subgraphs. Then the *clique covering number* of a graph is the fewest number of cliques in a clique covering. This number is denoted by $\text{cc}(G)$. It is well known that for all graphs G , $\text{mr}(G) \leq \text{cc}(G)$; see [7], and thus, we have the following corollary.

COROLLARY 2.6. *Let G be a graph, then $q(G) \leq \text{cc}(G) + 1$.*

In Corollary 3.6, a family of graphs is given for which this bound holds with equality.

We conclude this section with the exact value of $q(C_n)$ where C_n is a cycle on n vertices. This result can be derived from [9], but in this section, we will prove it using work from [10].

LEMMA 2.7. *Let C_n be the cycle on n vertices. Then*

$$q(C_n) = \left\lceil \frac{n}{2} \right\rceil.$$

Proof. First, suppose $n = 2k + 1$, for some $k \geq 1$. Then, the adjacency matrix of C_n has exactly $\frac{n-1}{2} + 1$ distinct eigenvalues, these eigenvalues are $2 \cos \frac{2\pi j}{n}$, $j = 1, \dots, n$. On the other hand, using [10, Cor. 3.4], any eigenvalue of $A \in S(C_n)$ has multiplicity at most two, so $q(C_n) \geq \frac{n-1}{2} + 1$. Thus, $q(C_n) = \frac{n-1}{2} + 1$.

Next, consider $n = 2k$, for some $k \geq 2$. Again using [10, Cor. 3.4], $q(C_n) \geq k$. Moreover, by [10, Thm. 3.3] for any set of numbers $\lambda_1 = \lambda_2 > \lambda_3 = \lambda_4 > \dots > \lambda_{2k-1} = \lambda_{2k}$, there is an $A \in S(C_n)$ with eigenvalues λ_i , $i = 1, \dots, n$. This implies that if n is even, then $q(C_n) = \frac{n}{2}$. \square

Since $\text{mr}(C_n) = n - 2$ and $q(C_n) \approx n/2$, we know that for some graphs G there can be a large gap between the parameters $\text{mr}(G)$ and $q(G)$.

3. Unique shortest path. There is only one family of graphs for which the eigenvalues for every matrix in $S(G)$ are all distinct, these are paths. This statement is Theorem 3.1 in [12], and also follows from a result by Fiedler [11], which states that for a real symmetric matrix $A \in M_n$ and a diagonal matrix D if $\text{rank}(A + D) \geq n - 1$, then $A \in S(P_n)$. A path on n vertices is denoted by P_n .

PROPOSITION 3.1. *For a graph G , $q(G) = |V(G)|$ if and only if G is a path.*

From this we can also conclude that the parameter q is not monotone on induced subgraphs; as $q(P_n) = n$ while $q(C_n) \approx n/2$. The next result is related to a very simple, but often very effective, lower bound on the minimum number of distinct eigenvalues of a graph that is based on the length of certain induced paths (see also similar notions presented in [6] and even in [12]). Recall that the *length* of a path is simply the number of edges in that path, and that the *distance between two vertices*, (in the same component) is the length of the shortest path between those two vertices.

THEOREM 3.2. *If there are vertices u, v in a connected graph G at distance d and the path of length d from u to v is unique, then $q(G) \geq d + 1$.*

Proof. Assume that $u = v_1, v_2, \dots, v_d, v_{d+1} = v$ is the unique path of length d

from u to v . For any $A = [a_{ij}] \in S(G)$, all of the matrices A, A^2, \dots, A^{d-1} have zero in the position (u, v) , while the entry (u, v) of A^d is equal to $\prod_{i=1}^d a_{v_i v_{i+1}} \neq 0$. Thus, the matrices I, A, A^2, \dots, A^d are linearly independent and the minimal polynomial of A must have degree at least $d + 1$. \square

It is important to note that the induced path from u to v in the proof of Theorem 3.2 is the shortest path from u to v and that it is the only path of this length. The length of such a path is a lower bound on the diameter of the graph and if the path is not unique, then the bound only holds for nonnegative matrices, see also [6].

COROLLARY 3.3. *For any connected graph G , if $A \in S(G)$ is nonnegative, then $q(A) \geq \text{diam}(G) + 1$.*

We note that Theorem 3.2 implies Theorem 3.1 from [12].

THEOREM 3.4. [12, Thm. 3.1] *Suppose G is a connected graph. If P is the longest induced path in G for which no edge of P lies on a cycle, then $q(G) \geq |V(P)|$.*

It is not true that $\text{diam}(G) + 1$ is a lower bound for the minimum number of distinct eigenvalues of an arbitrary graph G , see Corollary 6.9 for a counter-example. However, in the case of trees, since in this case, any shortest path between two vertices is the unique shortest path, we have

COROLLARY 3.5. *For any tree T , $q(T) \geq \text{diam}(T) + 1$.*

There are several other proofs of Corollary 3.5, see [6, 18]. There are also trees with $q(T) > \text{diam}(T) + 1$; see [2]. Further, for any positive integer d , there exists a constant $f(d)$ such that for any tree T with diameter d , there is a matrix $A \in S(T)$ with at most $f(d)$ distinct eigenvalues (this was shown by B. Shader [20] who described $f(d)$ as possibly “super-super-exponential”). It has been shown that $f(d) \geq (9/8)d$ for d large, see [18] and [19].

Using unique shortest paths, it is possible to construct a connected graph on n vertices with $q(G) = k$, for any pair of integers k, n with $1 \leq k \leq n$.

COROLLARY 3.6. *For any pair of integers k, n with $1 \leq k \leq n$, let $G(n, k)$ be the graph on vertices v_1, \dots, v_n , where vertices v_1, \dots, v_{n-k+2} form a clique and vertices $v_{n-k+2}, v_{n-k+3}, \dots, v_{n-1}, v_n$ form a path of length $k - 2$. Then, $q(G(n, k)) = k$.*

Proof. There is a unique shortest path of length $k - 1$ from v_n to any of the vertices v_1, \dots, v_{n-k+1} and there is a clique covering of the graph consisting of $k - 1$ cliques. Thus, by Corollary 2.6 and Theorem 3.2, $q(G(n, k)) = k$. \square

4. Graphs with two distinct eigenvalues. For a graph G , $q(G) = 2$ means that there is a matrix $A \in S(G)$ such that A has exactly two distinct eigenvalues,

and there is no matrix in $S(G)$ with only one eigenvalue. Therefore, the minimal polynomial of A has degree two, thus, A satisfies $A^2 = \alpha A + \beta I$, for some scalars α and β . This implies that A and A^2 have exactly the same zero-nonzero pattern on the off-diagonal entries. Equivalently, for any nonempty graph G , $q(G) = 2$ if and only if $S(G)$ contains a real symmetric orthogonal matrix Q . Using this, we can show the following results with the aid of Theorem 3.2.

LEMMA 4.1. *If $q(G) = 2$, for a connected graph G on $n \geq 3$ vertices, then G has no pendant vertex.*

Proof. Suppose vertex v_1 is pendant and suppose its unique neighbor is v_2 . Since G is connected and has at least 3 vertices there is another vertex v_3 that is adjacent to v_2 . Thus, there is a unique shortest path from v_1 to v_3 of length 2 and the result follows from Theorem 3.2. \square

The previous basic result is contained in the next slight generalization by noting that any edge incident with a pendant vertex is a *cut edge* (that is, its deletion results in a disconnected graph).

LEMMA 4.2. *Suppose G is a connected graph on n vertices with $n \geq 3$. If $q(G) = 2$, then there is no cut edge in the graph G .*

Proof. Assume that vertices v_1 and v_2 form a cut edge. We can assume without loss of generality that there is another vertex v_3 in G that is adjacent to v_2 . Thus, there is a unique shortest path from v_1 to v_3 of length 2 and the result follows from Theorem 3.2. \square

The next result should be compared to Theorem 3.4.

COROLLARY 4.3. *If G is a graph on $n \geq 3$ vertices with $q(G) = 2$, then every edge in G is contained in a cycle.*

Proof. Lemmas 4.1 and 4.2 together imply this result. \square

Consider $\alpha \subseteq \{1, 2, \dots, m\}$ and $\beta \subseteq \{1, 2, \dots, n\}$. For a matrix $A \in M_{m,n}$, $A[\alpha, \beta]$ denotes the submatrix of A lying in rows indexed by α and columns indexed by β . Recall that for any vertex v of a graph G , the *neighborhood set* of v , denoted by $N(v)$, is the set of all vertices in G adjacent to v .

THEOREM 4.4. *For a connected graph G on n vertices, if $q(G) = 2$, then for any independent set of vertices $\{v_1, v_2, \dots, v_k\}$, we have*

$$\left| \bigcup_{i \neq j} (N(v_i) \cap N(v_j)) \right| \geq k.$$

Proof. For the purpose of a contradiction, suppose G is a graph with $q(G) = 2$ and that there exists an independent set S with $|S| = k$ such that

$$\left| \bigcup_{i \neq j} (N(v_i) \cap N(v_j)) \right| < k,$$

and let $X = \bigcup_{i \neq j} (N(v_i) \cap N(v_j))$. Using Lemma 2.3, there exists a symmetric orthogonal matrix $A \in S(G)$. Consider such A and let $B = A[S, \{1, \dots, n\}]$. Observe that B is a $k \times n$ matrix and any column of B not indexed with X contains at most one nonzero entry. Since the rows of B are orthogonal, we deduce that rows of $C = A[S, X]$ must also be orthogonal. However, C is a $k \times |X|$ matrix with $k > |X|$, and orthogonality of these rows is impossible, as they are all nonzero. This completes the proof. \square

The next two statements are immediate, yet interesting, consequences of Theorem 4.4.

COROLLARY 4.5. *Let G be a connected graph on $n \geq 3$ vertices with $q(G) = 2$. Then, any two non-adjacent vertices must have at least two common neighbors.*

COROLLARY 4.6. *Suppose $q(G) = 2$, for a connected graph G on $n \geq 3$ vertices. If the vertex v_1 has degree exactly two with adjacent vertices v_2 and v_3 , then every vertex v that is different from v_2 and v_3 , has exactly the same neighbors as v_1 .*

Along these lines, we also note that if G is a connected graph with $q(G) = 2$, then for any independent set of vertices S , we have $|S| \leq n/2$. Thus, for any graph G with $q(G)$ being two, we have a basic upper bound on the size of any independent set in G .

As a final example, recall that $q(K_n) = 2$ whenever $n \geq 2$. We can build on this result for complete graphs with a single edge deleted.

PROPOSITION 4.7. *Suppose G is obtained from K_n by deleting a single edge e . Then*

$$q(G) = \begin{cases} 1, & \text{if } n = 2; \\ 3, & \text{if } n = 3; \\ 2, & \text{otherwise.} \end{cases}$$

Proof. The cases $n = 2, 3$ follow easily from previous facts. So suppose $n \geq 4$. We will construct a symmetric orthogonal matrix Q in $S(G)$, assuming the edge deleted was $e = \{1, n\}$, without loss of generality. In this case set,

$$u_1 = \frac{1}{\sqrt{n-1}} \begin{bmatrix} 1 \\ \frac{e_{n-2}}{0} \end{bmatrix},$$

where e_{n-2} is the $(n-2)$ -vector of all ones. Then choose u'_2 to be orthogonal to u_1 as follows

$$u'_2 = \begin{bmatrix} 0 \\ \frac{e_{n-3}}{-(n-3)} \\ 1 \end{bmatrix},$$

where e_{n-3} is the $(n-3)$ -vector of all ones. Then set $u_2 = \frac{1}{\|u'_2\|} u'_2$. Finally, set $Q = I - 2(u_1 u_1^T + u_2 u_2^T)$. Then it follows that Q is orthogonal and a basic calculation will show that $Q \in S(G)$. Hence, $q(G) = 2$. \square

5. Join of two graphs. In the previous section, we found several restrictions on a graph G for which $q(G) = 2$. In this section, we show that, despite these restrictions, a surprisingly large number of graphs satisfy this property.

Let G and H be graphs, then the *join of G and H* , denoted by $G \vee H$, is the graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H) \cup \{\{g, h\} \mid g \in V(G), h \in V(H)\}$.

A real matrix R of order n is called an *M -matrix* if it can be written in the form $R = sI - B$ for some $s > 0$ and entry-wise nonnegative matrix B such that its spectral radius satisfies $\rho(B) \leq s$. Recall that the *spectral radius* of a square matrix B is defined to be $\rho(B) = \max\{|\lambda| : \lambda \in \sigma(B)\}$. In the case that $\rho(B) < s$, then R is called a *nonsingular M -matrix*. Recall that for $A \in M_n$, we call $B \in M_n$ a *square root of A* if $B^2 = A$. In [1], it is shown that an M -matrix R has an M -matrix as a square root if and only if R has a certain property (which the authors refer to as *property c*). It is also known that all nonsingular M -matrices have “property c ”.

The following theorem is proved in [1]. If P is a square matrix, $\text{diag}(P)$ means the diagonal entries of P .

THEOREM 5.1. [1, Thm. 4] *Let R be an M -matrix of order n , and let $R = s(I - P)$ be a representation of R for sufficiently large s such that $\text{diag}(P)$ is entry-wise positive and $\rho(P) \leq 1$. Then R has an M -matrix as a square root if and only if R has “property c .” In this case, let Y^* denote the limit of the sequence generated by*

$$Y_{i+1} = \frac{1}{2}(P + Y_i^2), \quad Y_0 = 0.$$

Then $\sqrt{s}(I - Y^*)$ is an M -matrix with “property c ” which is a square root of R .

Using Theorem 5.1, we can prove the following.

THEOREM 5.2. *Let G be a connected graph, then $q(G \vee G) = 2$.*

Proof. Suppose G is a connected graph on n vertices. The goal of this proof is to construct a matrix P such that

$$Q = \begin{bmatrix} \sqrt{P} & \sqrt{I-P} \\ \sqrt{I-P} & -\sqrt{P} \end{bmatrix}$$

is in $S(G \vee G)$. If we can construct such a matrix P then $Q^2 = \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}$ and Q has exactly two eigenvalues. Let $A(G)$ be the adjacency matrix of G and set

$$P = \frac{2n-1}{4n^2} \left(\frac{1}{n}A(G) + I \right)^2.$$

Note that $\text{diag}(P)$ is entry-wise positive. By Gershgorin’s disc theorem (see [4, pg. 89]), every eigenvalue of $\frac{1}{n}A(G) + I$ belongs to the interval $(0, 2)$, and hence, every eigenvalue of P belongs to the interval $(0, \frac{2n-1}{n^2})$. Therefore, $\rho(P) \leq 1$.

Consider the matrix $R = I - P$. Then R is an M -matrix that satisfies the conditions of Theorem 5.1 with $s = 1$. Note that if S is a matrix with eigenvalue λ , then $1 - \lambda$ is an eigenvalue of $I - S$. Thus, the eigenvalues of R are in the interval $(\left(\frac{n-1}{n}\right)^2, 1)$. Hence, R is nonsingular and thus has “property c .” By Theorem 5.1, R has an M -matrix as a square root of the form $I - Y^*$, where Y^* is the limit of the sequence generated by

$$(5.1) \quad Y_{i+1} = \frac{1}{2}(P + Y_i^2), \quad Y_0 = 0.$$

Note that Y^* satisfies

$$(5.2) \quad (I - Y^*)^2 + \frac{2n-1}{4n^2} \left(\frac{1}{n}A(G) + I \right)^2 = I.$$

As G is a connected graph, its adjacency matrix $A(G)$ is an irreducible nonnegative matrix. Thus, $(aA(G) + bI)^n > 0$ for every $a, b > 0$, and hence, $Y^* > 0$ by (5.1).

As P is a real symmetric matrix, the sequence (and consequently the limit) of (5.1) are real symmetric matrices. In particular, Y^* is a real symmetric matrix that may be written as a polynomial in $A(G)$. Therefore, $I - Y^*$ commutes with $A(G)$.

If $\lambda_1, \lambda_2, \dots, \lambda_n$ are the eigenvalues of Y^* , then

$$\text{trace}(Y^*) = \sum_{i=1}^n \lambda_i < 1$$

as each eigenvalue of Y^* belongs to the interval $(0, \frac{1}{n})$. Therefore, $\text{diag}(I - Y^*) > 0$ implying that $I - Y^*$ is an entry-wise nonzero matrix.

Finally consider the block matrix

$$Q = \begin{bmatrix} \frac{\sqrt{2n-1}}{2n} (\frac{1}{n}A(G) + I) & I - Y^* \\ I - Y^* & -\frac{\sqrt{2n-1}}{2n} (\frac{1}{n}A(G) + I) \end{bmatrix}.$$

By (5.2), Q is an orthogonal matrix with two distinct eigenvalues. As $I - Y^*$ is entry-wise nonzero, $Q \in S(G \vee G)$, hence, $q(G \vee G) = 2$. \square

Recall that for any graph $G = (V, E)$, the graph $\overline{G} = (V, \overline{E})$, is called the *complement* of G whenever, $\overline{E} = \{\{i, j\} | \{i, j\} \notin E\}$.

COROLLARY 5.3. *There are graphs G for which the gap between $q(G)$ and $q(\overline{G})$ can grow without bound as a function of the number of vertices of G .*

Proof. Let $G = \overline{P_n} \vee \overline{P_n}$ with $n \geq 4$. Then $q(G) = 2$, while $q(\overline{G}) = q(P_n \cup P_n) = n$. \square

Also, note Theorem 5.2 fails to hold for two different graphs. For example, using Theorem 4.4, we have that $q(P_1 \vee P_4) > 2$. It is still unresolved whether or not the condition that G be connected is required in Theorem 5.2.

6. Bipartite graphs and graph products. Let G be a bipartite graph with parts X and Y such that $0 < |X| = m \leq n = |Y|$. Define $\mathcal{B}(G)$ to be the set of all real $m \times n$ matrices $B = [b_{ij}]$ whose rows and columns are indexed by X and Y , respectively, and for which $b_{ij} \neq 0$ if and only if $\{i, j\} \in E(G)$. We have the following:

THEOREM 6.1. *For any non-empty bipartite graph G , if $B \in \mathcal{B}(G)$, then $q(G) \leq 2q(BB^T) + 1$.*

Proof. Let $B \in \mathcal{B}(G)$ and consider $A \in \mathcal{S}(G)$ with $A = \begin{bmatrix} 0 & B \\ B^T & 0 \end{bmatrix}$. It is well known that BB^T and B^TB have the same nonzero eigenvalues, so the number of distinct nonzero eigenvalues of A^2 is at most $q(B^TB)$. Moreover, the eigenvalues of A are of the form $\pm\sqrt{\lambda}$, where λ is an eigenvalue of A^2 . Thus, A has at most $2q(B^TB) + 1$ distinct eigenvalues. \square

If B is square, then B^TB and BB^T have the same eigenvalues, this implies the following corollary.

COROLLARY 6.2. *For any non-empty bipartite graph G with equal sized parts, $q(G) \leq 2q(BB^T)$.*

LEMMA 6.3. *For any non-empty bipartite graph G , if there is a matrix $B \in \mathcal{B}(G)$ with orthogonal rows and orthogonal columns, then $q(G) = 2$.*

Proof. If $B \in \mathcal{B}(G)$ has orthogonal rows and orthogonal columns, then B is a square matrix. Consider $A \in \mathcal{S}(G)$ with $A = \begin{bmatrix} 0 & B \\ B^T & 0 \end{bmatrix}$. Then, $A^2 = I$, which implies that A has at most two distinct eigenvalues. Thus, by Lemma 2.1, $q(G) = 2$. \square

PROPOSITION 6.4. *Consider a bipartite graph G with parts X and Y . If $q(G) = 2$, then $|X| = |Y|$ and there exists an orthogonal matrix $B \in \mathcal{B}(G)$.*

Proof. Label the vertices of G so that the vertices of X come first. Then, any matrix in $\mathcal{S}(G)$ is of the form $A = \begin{bmatrix} D_1 & B \\ B^T & D_2 \end{bmatrix}$, where $D_1 \in M_{|X|}$ and $D_2 \in M_{|Y|}$ are diagonal matrices. Since $q(G) = 2$, using Lemma 2.3, $A \in \mathcal{S}(G)$ can be chosen with eigenvalues $-1, 1$, therefore $A^2 = I$. On the other hand,

$$A^2 = \begin{bmatrix} D_1^2 + BB^T & D_1B + BD_2 \\ B^TD_1 + D_2B^T & B^TB + D_2^2 \end{bmatrix}.$$

This implies that BB^T and B^TB are diagonal. Therefore, the rows and columns of B are orthogonal, and hence, $|X| = |Y|$. \square

For any $n \geq 1$, there is a real orthogonal $n \times n$ matrix all of whose entries are nonzero. For $n = 1, 2$ this is trivial, and for $n > 2$, the matrix $B = I - \frac{2}{n}J$ is such an orthogonal matrix.

Using the above example and Lemma 6.3, we have the following.

COROLLARY 6.5. *For any m, n with $1 \leq m \leq n$,*

$$q(K_{m,n}) = \begin{cases} 2, & \text{if } m = n; \\ 3, & \text{if } m < n. \end{cases}$$

Proof. If $m = n$, it is enough to normalize the real orthogonal matrix in the example preceding this Corollary and use it in Lemma 6.3. If $m < n$, then according to Proposition 6.4, we have $q(G) \geq 3$. On the other hand, the adjacency matrix of $K_{m,n}$ has 3 distinct eigenvalues. This completes the proof. \square

Next, we consider a group of bipartite graphs for which the lower bound given in Theorem 3.2 is tight. This family is closely related to the ‘‘tadpole graphs’’ discussed in [12] and are of interest since they are *parallel paths* (these graphs are discussed in Section 7). The exact value of the maximum multiplicity of parallel paths is known to be 2 (see [15]).

Define $S_{m,n}$ to be the graph consisting of a 4-cycle on vertices v_1, u_1, v_2, u_2 and edges $u_1v_2, v_2u_2, u_2v_1, v_1u_1$, together with a path P_{m+1} starting at vertex v_1 and a path P_{n+1} starting at v_2 , where P_{m+1} and P_{n+1} are disjoint from each other and they intersect the 4-cycle only on v_1 and v_2 , respectively. Label the vertices on the paths P_{m+1} and P_{n+1} by u_i and v_i , alternating the label u and v so that the graph can be considered as a bipartite graph with parts consisting of vertex sets $\{u_i\}$ and $\{v_i\}$. The graph $S_{m,n}$ has $m + n + 4$ vertices, and the graph $S_{4,4}$ is given in Figure 6.1.




Fig. 6.1: The graph $S_{4,4}$ and $q(S_{4,4}) = 6$.

LEMMA 6.6. *If m and n have the same parity, then*

$$q(S_{m,n}) = \max\{m, n\} + 2.$$

Proof. We assume that m and n are both even, the case when they are both odd is similar. We use the above labelling for $S_{m,n}$, and assume that $m \geq n$. Since there is a unique shortest path with $m + 2$ vertices from the pendant vertex on P_{m+1} to u_1 , by Theorem 3.2 we know that $q(S_{m,n}) \geq m + 2$. Define an $(m + n + 4)/2 \times (m + n + 4)/2$ matrix $B = [b_{ij}] \in \mathcal{B}(S_{m,n})$ with the rows labelled by the vertices u_i and the columns labelled by the vertices v_i . Let $b_{u_2v_1} = -1$ and $b_{u_2v_2} = b_{u_1v_1} = b_{u_1v_2} = 1$. Then, with the proper ordering of the vertices, BB^T has the form $\begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}$ where X is an $\frac{m+2}{2} \times \frac{m+2}{2}$ tridiagonal matrix and Y is an $\frac{n+2}{2} \times \frac{n+2}{2}$ tridiagonal matrix. Using the inverse eigenvalue problem for tridiagonal matrices [5] it is possible to find entries for B such that the eigenvalues for X are distinct and the eigenvalues for Y are a subset of the eigenvalues of X . Thus, $q(BB^T) = \frac{m+2}{2}$ and by Corollary 6.2, $q(S_{m,n}) \leq m + 2$. \square

Since $q(P_n) = n$ and $q(C_n) \approx n/2$, we know that addition of an edge can dramatically decrease the minimum number of distinct eigenvalues. Here we show that the addition of an edge to a graph can also dramatically increase the minimum number of distinct eigenvalues. To see this consider the graph G obtained by adding an edge between vertices u_1 and u_3 in the graph $S_{m,m}$ (see Figure 6.2). We know that $q(S_{m,m}) = m + 2$, but the new graph G has a unique shortest path that contains $2m + 2$ vertices from a pendant vertex to another pendant vertex. Thus, by Theorem 3.2, $q(G) \geq 2m + 2$, and we may conclude that there exist graphs G and an edge e




Fig. 6.2: The graph $S_{4,4}$ with a single edge added. The parameter q for this graph is at least 10.

such that the gap between $q(G)$ and $q(G - e)$ can grow arbitrarily large as a function of the number of vertices.

Similarly, if we consider the graph obtained from $S_{m,m}$ by adding a new vertex w and edges $\{w, u_1\}$ and $\{w, u_3\}$, then this new graph has a unique shortest path between the pendant vertices that contains $2m + 3$ vertices. Hence, there exists a family of graphs G with a vertex v of degree 2 such that the gap between $q(G)$ and $q(G \setminus v)$ can grow arbitrarily large as a function of the number of vertices.

We now switch gears and consider a graph product and a graph operation in an effort to compute q for more families of graphs. The product that we consider is the *Cartesian product*; if G and H are graphs then $G \square H$ is the graph on the vertex set $V(G) \times V(H)$ with $\{g_1, h_1\}$ and $\{g_2, h_2\}$ adjacent if and only if either $g_1 = g_2$ and h_1 and h_2 are adjacent in H or g_1 and g_2 are adjacent in G and $h_1 = h_2$.

THEOREM 6.7. *Let G be a graph on n vertices, then $q(G \square K_2) \leq 2q(G) - 2$.*

Proof. Let $A \in S(G)$ with $q(A) = q(G) = \ell$, and assume $\sigma(A) = \{\lambda_1, \lambda_2, \dots, \lambda_\ell\}$, where $\lambda_1 = 1, \lambda_2 = -1$. Let α, β be nonzero scalars, and consider the matrix

$$B = \begin{bmatrix} \alpha A & \beta I \\ \beta I & -\alpha A \end{bmatrix} \in S(G \square K_2).$$

Then,

$$B^2 = \begin{bmatrix} \alpha^2 A^2 + \beta^2 I & 0 \\ 0 & \alpha^2 A^2 + \beta^2 I \end{bmatrix}$$

and the eigenvalues of B^2 are of the form $\alpha^2 \lambda_i^2 + \beta^2$, for $i = 1, \dots, \ell$. If we choose $\alpha^2 + \beta^2 = 1$, then two eigenvalues of B^2 are equal to 1. Since $\pm \lambda$ is an eigenvalue of B whenever λ^2 is an eigenvalue of B^2 , this implies that $q(B) \leq 2(\ell - 1) = 2q(G) - 2$. \square

The following is implied by Lemma 2.1 and Theorem 6.7.

COROLLARY 6.8. *If $q(G) = 2$, then $q(G \square K_2) = 2$.*

Observe that Corollary 6.8 verifies that the bound in Theorem 6.7 can be tight.

COROLLARY 6.9. *If $n \geq 1$ is an integer, then the hypercube, Q_n , satisfies $q(Q_n) = 2$.*

Proof. Recall that Q_n can be defined recursively as $Q_n = Q_{n-1} \square K_2$, with $Q_1 = K_2$. Since $q(K_2) = 2$, the results follows by application of Corollary 6.8. \square

Note that the diameter of Q_n is n while $q(Q_n)$ is always 2, so for a graph that is not a tree the difference between the diameter and the minimum number of distinct eigenvalues can be arbitrarily large, as a function of the number of vertices.

Next we consider an operation on a graph. Let G be a graph, then the *corona of G* is the graph formed by joining a pendant vertex to each vertex of G .

LEMMA 6.10. *Let G be a graph and let G' be the corona of G , then $q(G') \leq 2q(G)$.*

Proof. Consider the matrix $B = \begin{bmatrix} A & I \\ I & 0 \end{bmatrix} \in S(G')$ when $A \in S(G)$. Assume that λ is an eigenvalue of B with the eigenvector $\begin{bmatrix} x \\ y \end{bmatrix}$. Then, $Ax + y = \lambda x$ and $x = \lambda y$. Hence, $\lambda \neq 0$, and $Ay = \frac{\lambda^2 - 1}{\lambda} y$. Therefore, $\mu = \frac{\lambda^2 - 1}{\lambda}$ is an eigenvalue of A . This implies that for each eigenvalue μ of A there are two real eigenvalues for $\lambda = \frac{\mu \pm \sqrt{\mu^2 + 4}}{2}$, this completes the proof. \square

7. Connected graphs with many distinct eigenvalues. In this section, we address the question of “which connected graphs have the minimum number of distinct eigenvalues near the number of vertices of the graph”. In Proposition 3.1, we observed that $q(G) = |V(G)|$ if and only if G is a path. In this section, we study the connected graphs G with the property that $q(G) = |V(G)| - 1$.

To begin we apply Theorem 3.2 to derive two families of graphs for which q is one less than the number of vertices.

PROPOSITION 7.1. *Let G be the graph with vertices v_1, v_2, \dots, v_n and edge set $E = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-1}, v_n\}, \{v_i, v_{i+2}\}\}$, where i is fixed and satisfies $1 \leq i \leq n - 2$. Then, $q(G) = |V(G)| - 1$.*

PROPOSITION 7.2. *Let G be the graph with vertices v_1, v_2, \dots, v_n and edge set $E = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{n-2}, v_{n-1}\}, \{v_i, v_n\}\}$, where i is fixed and satisfies $2 \leq i \leq n - 2$. Then, $q(G) = |V(G)| - 1$.*

Using Proposition 2.5 we may deduce that any graph G for which $q(G) = |V(G)| -$

1 implies $M(G) = 2$. However, even more can be said about such graphs.

THEOREM 7.3. *If G is a graph that satisfies $q(G) = |V(G)| - 1$, then G has the following properties:*

1. $M(G) = 2$.
2. *If A is in $S(G)$ and A has a multiple eigenvalue, then A has exactly one eigenvalue of multiplicity two, and all remaining eigenvalues are simple.*

The next result verifies that the graphs in Proposition 7.2 are the only trees with $q(G) = |V(G)| - 1$.

LEMMA 7.4. *Suppose T is a tree. If $q(T) = |V(T)| - 1$, then T consists of a path $P_{|V(T)|-1}$, along with a pendant vertex adjacent to a non-pendant vertex in this path.*

Proof. Since $q(T) = |V(T)| - 1$, it follows from Proposition 2.5 that $\text{mr}(T) \geq |V(T)| - 2$. Using Theorem 7.3 (1), $M(T) = 2$, and hence, the vertices of T can be covered by two vertex-disjoint paths (see [14]). Therefore, T consists of two induced paths P_1 and P_2 that cover all of the vertices of T along with exactly one edge connecting P_1 and P_2 . Then T has maximum degree equal to three and contains at most two vertices of degree three. Using Theorem 7.3 (2), if $q(T) = |V(T)| - 1$, then any matrix $A \in S(T)$ realizing an eigenvalue of (maximum) multiplicity two, has all other eigenvalue being simple. In [16], all such trees have been characterized, for all values of $M(T)$. In particular, from Theorem 1 in [16], we may conclude that the subgraph of T induced by the vertices of degree at least three must be empty. Thus, T has exactly one vertex of degree three. Furthermore, deletion of the vertex of degree three yields at most two components that contain more than one vertex (see [16, Thm. 1]), and hence must be of the claimed form. \square

Characterizing general connected graphs G with the property that $q(G) = |V(G)| - 1$ appears to be rather more complicated. By Theorem 7.3, we can restrict attention to certain graphs with $M(G) = 2$. Fortunately, the graphs with $M(G) = 2$ have been characterized in [15] and they include the graphs known as graphs of two parallel paths. A graph G is a *graph of two parallel paths* if there exist two disjoint induced paths (each on at least one vertex) that cover the vertices of G and any edge between these two paths can be drawn so as not to cross other edges (that is, there exists a planar embedding of G with all vertices incident to one face). The graphs $S_{m,n}$ described in Section 6 are examples of graphs of two parallel paths that satisfy $q(S_{m,n}) < |V(S_{m,n})| - 1$. Using [15], our investigation reduces to testing, which graphs G either of two parallel paths or from the exceptional list given in [15], satisfy $q(G) = |V(G)| - 1$.

We first, consider those graphs identified as exceptional type in [15, Fig. B1]. We let C_5 , C'_5 and C''_5 denote the graphs pictured in Figure 7.1, and refer to them as

base exceptional graphs, from which all other exceptional graphs can be formed by attaching paths of various lengths to the five vertices in each of C_5 , C'_5 and C''_5 .




Fig. 7.1: Base exceptional graphs.

LEMMA 7.5. *Each of the graphs C_5 , C'_5 and C''_5 in Figure 7.1 satisfy*

$$q(C_5) = q(C'_5) = q(C''_5) = 3.$$

Proof. We already know that $q(C_5) = 3$. For the remaining equalities it is enough to demonstrate the existence of a matrix with three distinct eigenvalues, since using Theorem 3.2 implies $q(C'_5), q(C''_5) \geq 3$. Consider C''_5 first. Let $A \in S(C''_5)$ be of the form

$$A = \left[\begin{array}{c|c} L & b \\ \hline b^T & a \end{array} \right],$$

where

$$L = \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 2 & -1 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 1 \end{bmatrix}$$

is the Laplacian matrix for a path on four vertices, namely $\{1, 2, 3, 4\}$. We will determine a and b based on some conditions in what follows. It is not difficult to check that the eigenvalues of L are $\{0, 2, 2 \pm \sqrt{2}\}$. The objective here is to choose a and b so that $A \in S(C''_5)$ and the eigenvalues of A are $\{0, 0, 2, 2, \lambda\}$ ($\lambda \neq 0, 2$). This can be accomplished by satisfying the following conditions:

1. $b = Lu = (L - 2I)w$ for some real vectors u, w ;
2. $a = u^T Lu = w^T (L - 2I)w + 2$; and
3. b has no zero entries.

If the eigenvectors of L associated with $2 \pm \sqrt{2}$ are x_2 and x_3 , respectively, then we know that u, w must be in the span of $\{x_2, x_3\}$. Hence, we may choose to write

$$u = \alpha_1 x_2 + \beta_1 x_3, \text{ and } w = \alpha_2 x_2 + \beta_2 x_3,$$

for some scalars $\alpha_1, \beta_1, \alpha_2, \beta_2$. (Note: u lies in the span of $\{e, x_2, x_3\}$, but we do not concern ourselves with e at this point.) In this case, (2) can be re-written as

$$(2 + \sqrt{2})\alpha_1^2 + (2 - \sqrt{2})\beta_1^2 = \sqrt{2}\alpha_2^2 - \sqrt{2}\beta_2^2 + 2,$$

and (1) can be re-written as

$$(2 + \sqrt{2})\alpha_1 x_2 + (2 - \sqrt{2})\beta_1 x_3 = \sqrt{2}\alpha_2 x_2 - \sqrt{2}\beta_2 x_3.$$

Since $\{x_2, x_3\}$ forms a linearly independent set of vectors, we have

$$\alpha_2 = \left(\frac{2 + \sqrt{2}}{\sqrt{2}} \right) \alpha_1, \text{ and } \beta_2 = \left(\frac{\sqrt{2} - 2}{\sqrt{2}} \right) \beta_1.$$

Substituting these values back into (2) gives,

$$(2 + \sqrt{2})\alpha_1^2 = (2 - \sqrt{2})\beta_1^2 - \sqrt{2}.$$

Thus, choosing β_1 large enough will suffice in satisfying all of the conditions (1)-(3) above. For example, if

$$\beta_1 = 2 \text{ and } \alpha_1 = -\sqrt{\frac{(2 - \sqrt{2})\beta_1^2 - \sqrt{2}}{2 + \sqrt{2}}},$$

then A , as constructed above, will have the desired form (that is $A \in S(C_5'')$) and with prescribed eigenvalues $\{0, 0, 2, 2, 18 - 9\sqrt{2}\}$. (The actual entries of A cannot be easily simplified so we have not displayed it here.) Hence, $q(C_5'') = 3$. Similar arguments can be applied to the graph C_5' , to conclude that $q(C_5') = 3$ as well. \square

In fact, using the above techniques, and the results obtained thus far we may deduce the following result.

THEOREM 7.6. *For connected graphs G on at most five vertices only the graphs from Propositions 7.1 and 7.2 satisfy $q(G) = |V(G)| - 1$.*

We have poured considerable effort into extending the above fact to larger orders, but this still has not been resolved. However, we have a strong suspicion that this fact can be extended. For instance, by Lemma 7.4, this is true for trees. Furthermore, using Lemma 7.5 and considering the result in the previous section on coronas, we feel strongly that all of the exceptional graphs G listed in [15, Fig. B1] satisfy $q(G) < |V(G)| - 1$.

8. Possible future directions. There are many open questions concerning the minimum number of distinct eigenvalues of a graph. In this section, we list some of them that we find interesting and provide some possible directions along these lines.

We have seen that adding an edge or a vertex can dramatically change the minimum number of distinct eigenvalues of a graph but we suspect that adding a pendant vertex to a graph could increase the minimum number of distinct eigenvalues by at most one. The next problem that we plan to work on is to determine how adding pendant vertices to a graph affects the minimum number of distinct eigenvalues.

We are also interested in how other graph operations affect the minimum number of distinct eigenvalues. For example, can we determine the minimum number of distinct eigenvalues of a graph that is the vertex sum of two graphs? Or what is the value of $q(G_1 \vee G_2)$ or $q(G \vee G \vee G)$ in general? Similarly, does Theorem 5.2 still hold if G is disconnected? We formulate the following unresolved idea for the join of two distinct graphs. If G_1 and G_2 are connected graphs and $|q(G_1) - q(G_2)|$ is small, then is $q(G_1 \vee G_2) = 2$?

Another unresolved issue deals with strongly-regular graphs. For any nonempty strongly-regular graph G , it is clear that $2 \leq q(G) \leq 3$. Thus, a key question is which strongly-regular graphs satisfy $q(G) = 2$? The complete bipartite graph $K_{n,n}$ and K_n are examples of such graphs. By Corollary 4.5, if G is a strongly-regular graph with parameters (n, k, a, c) (see [4, Chap. 5]), where c is the number of mutual neighbors of any two non-adjacent vertices, and $q(G) = 2$, then $c \geq 2$ (but this is hardly a strong restriction). However, this restriction on c does verify that the minimum number of distinct eigenvalues for the Petersen graph is three. In addition the complete multipartite graphs of the form $G = K_{n_1, n_1, n_2, n_2, \dots, n_k, n_k}$, where n_i are arbitrary positive numbers, have $q(G) = 2$; because G is the join of K_{n_1, n_2, \dots, n_k} with itself; and obviously G is strongly regular. Also, $q(K_{2,2,2}) = 2$ as a 6×6 real symmetric orthogonal matrix can be constructed with 2×2 zero blocks on the diagonal and Hadamard-like 2×2 matrices off the diagonal. Observe that $K_{2,2,2}$ is also strongly regular but is not the join of a graph with itself. At present we are still not sure about $q(K_{2,2,\dots,2})$ or $q(K_{3,3,3})$.

Finally, the last outstanding issue is the characterization of all graphs G for which $q(G) = |V(G)| - 1$. Towards this end, as we eluded to in Section 7, we presented a number of ideas and directions towards a general characterization.

Acknowledgment. We would like to thank Dr. Francesco Barioli and Dr. Robert Bailey for a number of interesting discussions related to this topic, and other connections to certain spectral graph theory problems.

REFERENCES

- [1] G. Alefeld and N. Schneider. On square roots of M -matrices. *Linear Algebra Appl.*, 42:119–132, 1982.
- [2] F. Barioli and S. Fallat. On two conjectures regarding an inverse eigenvalue problem for acyclic symmetric matrices. *Electron. J. Linear Algebra*, 11:41–50, 2004
- [3] M. Booth et al. On the minimum rank among positive semidefinite matrices with a given graph. *SIAM Matrix Anal. Appl.*, 30:731–740, 2008.
- [4] R. Brualdi and H.J. Ryser. *Combinatorial Matrix Theory*. Cambridge University Press, New York, 1991.
- [5] A.L. Duarte. Construction of acyclic matrices from spectral data. *Linear Algebra Appl.*, 113:173–182, 1989.
- [6] A.L. Duarte and C.R. Johnson. On the minimum number of distinct eigenvalues for a symmetric matrix whose graph is a given tree. *Math. Inequal. Appl.*, 5:175–180, 2002.
- [7] S. Fallat and L. Hogben. The minimum rank of symmetric matrices described by a graph: A survey. *Linear Algebra Appl.*, 426:558–582, 2007.
- [8] S. Fallat and L. Hogben. Variants on the minimum rank problem: A survey II. Preprint, arXiv:1102-5142v1, 2011.
- [9] W.E. Ferguson. The construction of Jacobi and periodic Jacobi matrices with prescribed spectra. *Math. Comp.*, 35:1203–1220, 1980.
- [10] R. Fernandes and C.M. da Fonseca. The inverse eigenvalue problem for Hermitian matrices whose graphs are cycles. *Linear Multilinear Algebra*, 57:673–682, 2009.
- [11] M. Fiedler. A characterization of tridiagonal matrices. *Linear Algebra Appl.*, 2:191–197, 1969.
- [12] C.M. da Fonseca. A lower bound for the number of distinct eigenvalues of some real symmetric matrices. *Electron. J. Linear Algebra*, 21:3–11, 2010.
- [13] L. Hogben. Spectral graph theory and the inverse eigenvalue problem of a graph. *Electron. J. Linear Algebra*, 14:12–31, 2005.
- [14] C. Johnson and A.L. Duarte. The maximum multiplicity of an eigenvalue in a matrix whose graph is a tree. *Linear Multilinear Algebra*, 46:139–144, 1999.
- [15] C. Johnson, R. Loewy and P.A. Smith. The graphs for which the maximum multiplicity of an eigenvalue is two. *Linear Multilinear Algebra*, 57:713–736, 2009.
- [16] C. Johnson and C. Saigo. The trees for which maximum multiplicity implies the simplicity of other eigenvalues. *Discrete Math.*, 306:3130–3135, 2006.
- [17] I.-J. Kim and B.L. Shader. Classification of trees each of whose associated acyclic matrices with distinct diagonal entries has distinct eigenvalues. *Bull. Korean Math. Soc.*, 28:95–99, 2008.
- [18] I.-J. Kim and B.L. Shader. Smith normal form and acyclic matrices. *J. Algebraic Combin.*, 29:63–80, 2009.
- [19] N. Reshmi. *Acyclic Matrices with an Extreme Number of Distinct Eigenvalues*. Ph.D. Dissertation, University of Wyoming, 2011.
- [20] B.L. Shader. Personal Communication, 2011.