

2017

## On skew-symmetric matrices related to the vector cross product in $\mathbb{R}^7$

P. D. Beites

*University of Beira Interior*, pbeites@ubi.pt

A. P. Nicolás

*University of Valladolid*, alejandro.pinera@uva.es

José Vitória

*University of Coimbra*, jvitoria@mat.uc.pt

Follow this and additional works at: <http://repository.uwyo.edu/ela>



Part of the [Algebra Commons](#)

---

### Recommended Citation

Beites, P. D.; Nicolás, A. P.; and Vitória, José. (2017), "On skew-symmetric matrices related to the vector cross product in  $\mathbb{R}^7$ ", *Electronic Journal of Linear Algebra*, Volume 32, pp. 138-150.

DOI: <https://doi.org/10.13001/1081-3810.3498>

This Article is brought to you for free and open access by Wyoming Scholars Repository. It has been accepted for inclusion in Electronic Journal of Linear Algebra by an authorized editor of Wyoming Scholars Repository. For more information, please contact [scholcom@uwyo.edu](mailto:scholcom@uwyo.edu).



## ON SKEW-SYMMETRIC MATRICES RELATED TO THE VECTOR CROSS PRODUCT IN $\mathbb{R}^7$ \*

PATRÍCIA D. BEITES<sup>†</sup>, ALEJANDRO P. NICOLÁS<sup>‡</sup>, AND JOSÉ VITÓRIA<sup>§</sup>

**Abstract.** A study of real skew-symmetric matrices of orders 7 and 8, defined through the vector cross product in  $\mathbb{R}^7$ , is presented. More concretely, results on matrix properties, eigenvalues, (generalized) inverses and rotation matrices are established.

**Key words.** Vector cross product, Skew-symmetric matrix, Matrix properties, Eigenvalues, (Generalized) Inverses, Rotation matrices.

**AMS subject classifications.** 15A72, 15B57, 15A18, 15A09, 15B10.

**1. Introduction.** Let  $F$  be a field of characteristic different from 2. An algebra  $A$  over  $F$  is a composition algebra if it is endowed with a nondegenerate quadratic form (the norm)  $n : A \rightarrow F$  which is multiplicative, i.e., for any  $x, y \in A$ ,

$$n(xy) = n(x)n(y).$$

The form  $n$  being nondegenerate means that  $n(x, y) = \frac{1}{2}(n(x+y) - n(x) - n(y))$ , the associated symmetric bilinear form, is nondegenerate.

A classical result known as the generalized Hurwitz Theorem asserts that, over  $F$ , if  $\mathcal{A}$  is a finite dimensional composition algebra with identity, then its dimension is equal to 1, 2, 4 or 8. Furthermore,  $\mathcal{A}$  is isomorphic either to the base field, a separable quadratic extension of the base field, a generalized quaternion algebra or a generalized octonion algebra, [8].

A consequence of the cited theorem is that the values of  $n$  for which the Euclidean spaces  $\mathbb{R}^n$  can be equipped with a binary vector cross product, satisfying the same requirements as the usual one in  $\mathbb{R}^3$ , are restricted to 1 (trivial case), 3 and 7. A complete account on the existence of  $r$ -fold vector cross products for  $d$ -dimensional vector spaces, where they are used to construct exceptional Lie superalgebras, is in [4].

The interest in octonions, seemingly forgotten for some time, resurged in the recent decades, not only for their intrinsic mathematical relevance but also because of their applications, as well as those of the vector cross product in  $\mathbb{R}^7$ . This product was used for the implementation of the seven-dimensional vector analysis method in [15], to estimate the amount of abnormalities in algorithms that provide accurate feedback in rehabilitation.

---

\*Received by the editors on February 18, 2017. Accepted for publication on April 11, 2017. Handling Editor: Michael Tsatsomeros.

<sup>†</sup>Departamento de Matemática and Centro de Matemática e Aplicações (CMA-UBI), Universidade da Beira Interior, Portugal (pbeites@ubi.pt). Supported by Fundação para a Ciência e a Tecnologia (Portugal), project PEst-OE/MAT/UI0212/2013 of CMA-UBI, and the research project MTM2013-45588-C3-1-P (Spain).

<sup>‡</sup>Departamento de Matemática Aplicada and Instituto de Investigación en Matemáticas (IMUVa), Universidad de Valladolid, Spain (alejandro.pinera@uva.es). Supported by the research project MTM2013-45588-C3-1-P (Spain).

<sup>§</sup>Departamento de Matemática, Universidade de Coimbra, Portugal (jvitoria@mat.uc.pt).

Moreover, as mentioned in [9], the octonions play an important role in Physics. Namely, they led up to the theory of fundamental particles known as eightfold way. More recently, in [11], it was shown that if the fundamental particles, the fermions, are assumed to have seven time-spatial dimensions, then the so called hierarchy problem, concerning the unknown reason for the weak force to be stronger than the gravity force, could be solved.

In this work, we extend the results, devoted to the vector cross product in  $\mathbb{R}^3$  and real skew-symmetric matrices of order 4, in [17]. Concretely, we study real skew-symmetric matrices of orders 7 and 8 defined through the vector cross product in  $\mathbb{R}^7$ . These are denoted, for any  $a, b \in \mathbb{R}^7$ , by  $S_a$  and  $M_{a,b}$ , respectively.

The latter matrices, called hypercomplex in [9], can be used to write the coordinate matrix of the left multiplication by an octonion. The particular case  $b = a$  leads to  $M_{a,a}$ , an orthogonal design which, according to [14] and references therein, can be used in the construction of space time block codes for wireless transmissions. Furthermore, if  $b = a = [1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1]^T$ , then  $I_8 + M_{a,a}$  is a Hadamard matrix of skew-symmetric type.

For completeness, in Section 2, we recall some definitions and results related to the binary vector cross product in  $\mathbb{R}^7$ , inverses and skew-symmetric matrices. Throughout the work, for simplicity, we omit the word binary.

In Section 3, we approach the vector cross product in  $\mathbb{R}^7$  from a matrix point of view. For this purpose, we consider the matrices  $S_a$  and establish some related properties.

Section 4 is devoted to the eigenvalues of  $S_a$  and  $M_{a,b}$ . We obtain the characteristic polynomials of these matrices, using adequate Schur complements in the latter case.

In Section 5, we deduce either the inverse or the Moore-Penrose inverse of  $M_{a,b}$ , depending on its determinant. The Moore-Penrose inverse of  $S_a$  is presented in Section 3.

We dedicate Section 6, the last of this work, to the generation of rotation matrices from the Cayley transforms and the exponentials of the skew-symmetric matrices  $S_a$  and  $M_{a,b}$ .

**2. Preliminaries.** Throughout this work,  $\mathbb{R}^{m \times n}$  denotes the set of all  $m \times n$  real matrices. With  $n = 1$ , we identify  $\mathbb{R}^{m \times 1}$  with  $\mathbb{R}^m$ . With  $m = n = 1$ , we identify  $\mathbb{R}^{1 \times 1}$  with  $\mathbb{R}$ .

Consider the usual real vector space  $\mathbb{R}^8$ , with canonical basis  $\{e_0, \dots, e_7\}$ , equipped with the multiplication  $*$  given by  $e_i * e_i = -e_0$  for  $i \in \{1, \dots, 7\}$ , being  $e_0$  the identity, and the below Fano plane, where the cyclic ordering of each three elements lying on the same line is shown by the arrows.




FIGURE 1. Fano plane for  $\mathbb{O}$ .

Then  $\mathbb{O} = (\mathbb{R}^8, *)$  is the real (non-split) octonion algebra. Every element  $\underline{x} \in \mathbb{O}$  may be represented by

$$\underline{x} = x_0 + x, \quad \text{where } x_0 \in \mathbb{R} \text{ and } x = \sum_{i=1}^7 x_i e_i \in \mathbb{R}^7,$$

are, respectively, the *real part* and the *pure part* of the octonion  $\underline{x}$ .

The multiplication  $*$  can be written in terms of the Euclidean inner product and the vector cross product in  $\mathbb{R}^7$ , hereinafter denoted by  $\langle \cdot, \cdot \rangle$  and  $\times$ , respectively. Concretely, as in [9], we have

$$\underline{x} * \underline{y} = x_0 y_0 - \langle x, y \rangle + x_0 y + y_0 x + x \times y.$$

A formula for the *double vector cross product* in  $\mathbb{R}^7$  is

$$x \times (y \times z) = \langle x, z \rangle y - \langle x, y \rangle z + \frac{1}{3} J(x, y, z),$$

[10]. Here  $J$  stands for the *Jacobian*, the alternate application defined by

$$J(x, y, z) = x \times (y \times z) + y \times (z \times x) + z \times (x \times y).$$

For  $(\mathbb{R}^3, \times)$ , a Lie algebra, the well known formula for the double vector cross product in  $\mathbb{R}^3$  arises since, for any  $x, y, z \in \mathbb{R}^3$ ,  $J(x, y, z) = 0$ .

Let  $A \in \mathbb{R}^{m \times n}$ .

A matrix  $A^- \in \mathbb{R}^{n \times m}$  is a *generalized inverse* of  $A$  if  $AA^-A = A$ .

The *Moore-Penrose inverse* of  $A$ , as defined in [1], is the unique matrix  $A^\dagger \in \mathbb{R}^{n \times m}$  satisfying

$$AA^\dagger A = A, \quad A^\dagger AA^\dagger = A^\dagger, \quad (A^\dagger A)^T = A^\dagger A \quad \text{and} \quad (AA^\dagger)^T = AA^\dagger.$$

In particular, if  $u$  is a nonzero vector in  $\mathbb{R}^{m \times 1}$ , then its Moore-Penrose inverse is given by

$$u^\dagger = \frac{u^T}{\|u\|^2},$$

where, hereinafter,  $\|\cdot\|$  denotes the Euclidean norm.

In the remaining part of this section, assume that  $m = n$ .

The matrix  $A$  is a *rotation matrix* if  $A$  is orthogonal ( $A^T A = I$ ) and  $\det A = 1$ .

From now on, assume also that  $A$  is a skew-symmetric matrix. Hence, according to a classical result on skew-symmetric matrices, the eigenvalues of  $A$  are purely imaginary or null.

Due to the skew-symmetry of  $A$ ,  $I_n + A$  is invertible. The *Cayley transform* of  $A$  is the matrix given by  $\mathcal{C}(A) = (I_n + A)^{-1}(I_n - A)$ . It is well known that  $\mathcal{C}(A)$  is a rotation matrix and, as  $I_n - A = 2I_n - (I_n + A)$ ,

$$\mathcal{C}(A) = 2(I_n + A)^{-1} - I_n.$$

This is one of the Cayley formulas in [6], that allow to establish a one-to-one correspondence between the skew-symmetric matrices and the orthogonal matrices that do not have the eigenvalue  $-1$ .

As it is known,  $R = e^A$  is the rotation matrix, called *exponential* of  $A$ , defined by the absolute convergent power series

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!}.$$

Conversely, given a rotation matrix  $R \in \mathbf{SO}(n)$ , there exists a skew-symmetric matrix  $A$  such that  $R = e^A$ , [6]. The combination of these two facts is equivalent to saying that the map  $\exp : \mathfrak{so}(n) \rightarrow \mathbf{SO}(n)$ , from the Lie algebra  $\mathfrak{so}(n)$  of skew-symmetric  $n \times n$  matrices to the Lie group  $\mathbf{SO}(n)$ , is surjective, [2].

**3. Matrix properties of  $S_a$ .** In the present section, following [9] and [10], we consider a matrix representation of the Maltsev algebra  $(\mathbb{R}^7, \times)$  in terms of particular cases of hypercomplex matrices. If  $a \in \mathbb{R}^7$ , then let  $S_a$  be the matrix in  $\mathbb{R}^{7 \times 7}$  defined by

$$S_a x = a \times x$$

for any  $x \in \mathbb{R}^7$ . Hence, for  $a = [a_1 \ a_2 \ a_3 \ a_4 \ a_5 \ a_6 \ a_7]^T$ ,  $S_a$  is the skew-symmetric matrix

$$\begin{bmatrix} 0 & -a_3 & a_2 & -a_5 & a_4 & -a_7 & a_6 \\ a_3 & 0 & -a_1 & -a_6 & a_7 & a_4 & -a_5 \\ -a_2 & a_1 & 0 & a_7 & a_6 & -a_5 & -a_4 \\ a_5 & a_6 & -a_7 & 0 & -a_1 & -a_2 & a_3 \\ -a_4 & -a_7 & -a_6 & a_1 & 0 & a_3 & a_2 \\ a_7 & -a_4 & a_5 & a_2 & -a_3 & 0 & -a_1 \\ -a_6 & a_5 & a_4 & -a_3 & -a_2 & a_1 & 0 \end{bmatrix}.$$

We now establish some properties related to  $S_a$ .

PROPOSITION 3.1. *Let  $a, c \in \mathbb{R}^7$  and  $\alpha, \gamma \in \mathbb{R}$ . Then*

- 1)  $S_{\alpha a + \gamma c} = \alpha S_a + \gamma S_c$ ;
- 2)  $S_a c = -S_c a$ ;
- 3)  $S_a$  is singular;
- 4)  $S_a^2 = aa^T - a^T a I_7$ ;
- 5)  $S_a^3 = -a^T a S_a$ ;
- 6)  $S_a^\dagger = \begin{cases} 0 & \text{if } a = 0 \\ -\frac{1}{a^T a} S_a & \text{if } a \neq 0 \end{cases}$ ;
- 7)  $S_{S_a b} = \frac{3}{2}(ba^T - ab^T) - \frac{1}{2}[S_a, S_b]$ , where  $[\cdot, \cdot]$  denotes the matrix commutator.

*Proof.* Properties 1) and 2) are direct consequences of the bilinearity and of the skew-symmetry of  $\times$ .

As far as 3), on the one hand, if  $a = 0$  then  $S_a = 0$ , being  $S_a$  singular. On the other hand, if  $a \neq 0$  then, from 2), we have  $S_a a = 0$ . If  $S_a$  was invertible then  $a = 0$ , a contradiction.

Regarding 4), for any  $x \in \mathbb{R}^7$ , we have

$$S_a S_a x = a \times (a \times x) = \langle a, x \rangle a - \langle a, a \rangle x = (aa^T)x - (a^T a)x = (aa^T - a^T a I_7)x.$$

Concerning 5), note that  $aa^T S_a = -a(S_a a)^T = 0$  by 2). Hence, by 4),  $S_a^3 = S_a^2 S_a = -a^T a S_a$ .

To obtain 6), since the case  $a = 0$  is trivial, assume that  $a \neq 0$ . By 5),

$$S_a \left( -\frac{1}{a^T a} S_a \right) S_a = -\frac{1}{a^T a} S_a^3 = S_a.$$

Taking into account the skew-symmetry of  $S_a$ ,

$$\left( S_a \frac{-1}{a^T a} S_a \right)^T = -\frac{1}{a^T a} S_a^T S_a^T = S_a \frac{-1}{a^T a} S_a.$$

The remaining equalities of the Moore-Penrose inverse definition can be proved in a similar way.

By 2), for any  $x \in \mathbb{R}^7$ , we get

$$S_{S_a b} x = -S_x S_a b = -x \times (a \times b) = -\langle x, b \rangle a + \langle x, a \rangle b - \frac{1}{3} J(x, a, b).$$

As  $J(x, a, b) = x \times (a \times b) + a \times (b \times x) + b \times (x \times a) = -S_{a \times b} x + [S_a, S_b] x$ , then we obtain

$$S_{S_a b} x = \left( b a^T - a b^T + \frac{1}{3} S_{S_a b} - \frac{1}{3} [S_a, S_b] \right) x,$$

and 7) follows. □

REMARK 3.2. The problem of finding the Moore-Penrose inverse of an order 7 skew-symmetric matrix was proposed by Groß, Troschke and Trenkler in [7]. The ensuing solutions can be found in [16], the proof of 6) in Proposition 3.1 being similar, although independently obtained, to the solution presented therein by Bapat.

**4. Eigenvalues of  $S_a$  and  $M_{a,b}$ .** In this section and in the following ones, we consider real skew-symmetric matrices of order 8 written as bordered matrices in the partitioned form

$$M_{a,b} = \begin{bmatrix} S_a & b \\ -b^T & 0 \end{bmatrix},$$

with  $b = [b_1 \ b_2 \ b_3 \ b_4 \ b_5 \ b_6 \ b_7]^T$ ,  $a = [a_1 \ a_2 \ a_3 \ a_4 \ a_5 \ a_6 \ a_7]^T \in \mathbb{R}^{7 \times 1}$ . These  $8 \times 8$  matrices constitute a generalization of the  $4 \times 4$  matrices in [17].

PROPOSITION 4.1. [13] *Let  $E \in \mathbb{R}^{r \times r}$ ,  $F \in \mathbb{R}^{r \times 1}$  and  $G \in \mathbb{R}^{r \times 1}$ . Then*

$$\det \begin{bmatrix} E & F \\ G^T & 0 \end{bmatrix} = -G^T \text{adj}(E) F.$$

THEOREM 4.2. *The determinant of  $M_{a,b}$  is*

$$\det(M_{a,b}) = (a^T a)^2 (a^T b)^2.$$

*Proof.* By Proposition 4.1,

$$\det(M_{a,b}) = b^T \text{adj}(S_a) b,$$

where  $\text{adj}(S_a)$  is the adjugate of  $S_a$ . Some straightforward calculations lead to  $\text{adj}(S_a) = (a^T a)^2 (a a^T)$ . Hence,  $\det(M_{a,b}) = b^T (a^T a)^2 (a a^T) b = (a^T a)^2 (a^T b)^2$ . □

Before proceeding to the problem of determining the eigenvalues of  $S_a$  and  $M_{a,b}$ , we recall a result related to block determinants.

PROPOSITION 4.3. [13] Let  $E \in \mathbb{R}^{r \times r}$ ,  $F \in \mathbb{R}^{r \times s}$ ,  $G \in \mathbb{R}^{s \times r}$  and  $H \in \mathbb{R}^{s \times s}$ .

$$\det \begin{bmatrix} E & F \\ G & H \end{bmatrix} = \begin{cases} \det(E) \det(H - GE^{-1}F) & \text{when } E^{-1} \text{ exists} \\ \det(H) \det(E - FH^{-1}G) & \text{when } H^{-1} \text{ exists} \end{cases},$$

where  $H - GE^{-1}F$  and  $E - FH^{-1}G$  are the Schur complements of  $E$  and  $H$ , respectively.

THEOREM 4.4. The characteristic polynomial of  $M_{a,b}$  is

$$p_{M_{a,b}}(\lambda) = (\lambda^2 + a^T a)^2 (\lambda^4 + \lambda^2 (a^T a + b^T b) + (a^T b)^2).$$

*Proof.* The characteristic polynomial of  $M_{a,b}$  is given by

$$p_{M_{a,b}}(\lambda) = \det(M_{a,b} - \lambda I_8) = \det \begin{bmatrix} S_a - \lambda I_7 & b \\ -b^T & -\lambda \end{bmatrix}.$$

If  $\lambda = 0$  then  $p_{M_{a,b}}(0) = \det(M_{a,b}) = (a^T a)^2 (a^T b)^2$ . Assume that  $\lambda \neq 0$ . Then  $S_a - \lambda I_7$  is invertible. Since, through straightforward calculations, the adjugate and the determinant of this matrix are, respectively,

$$(\lambda^2 + a^T a)^2 (\lambda(S_a + \lambda I_7) + a a^T) \quad \text{and} \quad -\lambda(\lambda^2 + a^T a)^3,$$

then

$$(S_a - \lambda I_7)^{-1} = -\frac{1}{\lambda^2 + a^T a} \left( S_a + \lambda I_7 + \frac{1}{\lambda} a a^T \right).$$

By Proposition 4.3,

$$\det(M_{a,b} - \lambda I_8) = \det(S_a - \lambda I_7) (-\lambda + b^T (S_a - \lambda I_7)^{-1} b).$$

As  $b^T a a^T b = (a^T b)^2$  and, by 2) of Proposition 3.1,  $b^T S_a b = 0$ , we arrive at  $p_{M_{a,b}}(\lambda) = \det(M_{a,b} - \lambda I_8) = (\lambda^2 + a^T a)^2 (\lambda^4 + \lambda^2 (a^T a + b^T b) + (a^T b)^2)$ .  $\square$

COROLLARY 4.5. The eigenvalues of  $S_a$  are 0 and  $\pm \|a\| i$ .

*Proof.* Since the characteristic polynomial of  $S_a$  is  $-\lambda(\lambda^2 + a^T a)^3$ , it is a consequence of the proof of Theorem 4.4.  $\square$

COROLLARY 4.6. The eigenvalues of  $M_{a,b}$  are the purely imaginary numbers

$$\pm \|a\| i \quad \text{and} \quad \pm \sqrt{\frac{1}{2} (\|a\|^2 + \|b\|^2 \pm \|a - b\| \|a + b\|)} i.$$

*Proof.* From Theorem 4.4, putting  $\lambda^2 = x$  in  $p_{M_{a,b}}(\lambda)$ , we obtain

$$(x + a^T a)^2 (x^2 + (a^T a + b^T b)x + (a^T b)^2) = 0.$$

Thus,

$$x = -a^T a \quad \text{or} \quad x_{1,2} = -\frac{a^T a + b^T b}{2} \pm \frac{\sqrt{(a^T a + b^T b)^2 - 4(a^T b)^2}}{2}.$$

We have  $x_1 + x_2 = -(a^T a + b^T b)$  and  $x_1 x_2 = (a^T b)^2$ . So, invoking Girard-Newton-Viète laws,  $x_1 \leq 0$  and  $x_2 \leq 0$ . Finally, a straightforward computation leads to the result since

$$x = -a^T a \quad \text{or} \quad x_{1,2} = -\frac{a^T a + b^T b}{2} \pm \frac{\sqrt{(a-b)^T(a-b)(a+b)^T(a+b)}}{2}. \quad \square$$

REMARK 4.7. Assume that  $a$  and  $b$  are orthogonal vectors. So,  $\|a\|^2 + \|b\|^2 = \|a+b\|^2$ . By Corollary 4.6, the eigenvalues of  $M_{a,b}$  are  $\pm\|a\|i, 0$  and  $\pm\|a+b\|i$ . Invoking Gerschgorin's Theorem in [13], we obtain

$$\|a+b\| \leq \max\{r_i : i \in \{1, \dots, 8\}\},$$

$$\text{where } r_t = \sum_{\substack{s=1 \\ s \neq t}}^7 |a_s| + |b_t| \text{ for } t \in \{1, \dots, 7\}, \quad r_8 = \sum_{k=1}^7 |b_k|.$$

Taking  $a = [1 \ -1 \ 1 \ -1 \ 0 \ 0 \ 0]^T$  and  $b = [1 \ 1 \ 1 \ 1 \ 1 \ 1 \ 1]^T$ , we see that this upper bound can be sharper than  $\|a\| + \|b\|$ , the one given by the triangle inequality. Concretely, we get  $\max\{3, 4\} < 2 + \sqrt{7}$ .

**5. Inverses of  $M_{a,b}$ .** The Moore-Penrose inverse of  $S_a$  was characterized in Section 3. Depending on the determinant of  $M_{a,b}$ , either the inverse or the Moore-Penrose inverse of  $M_{a,b}$  may be determined. For this purpose, we recall the following result where  $*$ ,  $R(A)$  and  $N(A)$  stand for the conjugate transpose of a matrix, the column space of  $A$  and the nullspace of  $A$ , respectively.

THEOREM 5.1. [12] *Let  $T$  denote the complex bordered matrix  $\begin{bmatrix} A & c \\ d^* & \alpha \end{bmatrix}$ , where  $A$  is  $m \times m$ ,  $c$  and  $d$  are columns, and  $\alpha$  is a scalar. Let  $k = A^\dagger c$ ,  $h^* = d^* A^\dagger$ ,  $u = (I - AA^\dagger)c$ ,  $v = (I - A^\dagger A)d$ ,  $w_1 = 1 + k^* k$ ,  $w_2 = 1 + h^* h$  and  $\beta = \alpha - d^* A^\dagger c$ . Then*

1.  $\text{rank}(T) = \text{rank}(A) + 2$  if and only if  $c \notin R(A)$  and  $d \notin R(A^*)$ ,
2.  $\text{rank}(T) = \text{rank}(A)$  if and only if  $c \in R(A)$ ,  $d \in R(A^*)$  and  $\beta = 0$ .

The Moore-Penrose inverse of  $T$  is as follows.

- 1) When  $\text{rank}(T) = \text{rank}(A) + 2$ ,

$$T^\dagger = \begin{bmatrix} A^\dagger - ku^\dagger - v^{*\dagger}h^* - \beta v^{*\dagger}u^\dagger & v^{*\dagger} \\ u^\dagger & 0 \end{bmatrix}.$$

- 2) When  $\text{rank}(T) = \text{rank}(A)$ ,

$$T^\dagger = \begin{bmatrix} A^\dagger - w_1^{-1}kk^*A^\dagger - w_2^{-1}A^\dagger hh^* & w_2^{-1}A^\dagger h \\ w_1^{-1}k^*A^\dagger & 0 \end{bmatrix} + \frac{k^*A^\dagger h}{w_1 w_2} \begin{bmatrix} k \\ -1 \end{bmatrix} \begin{bmatrix} h^* & -1 \end{bmatrix}.$$

PROPOSITION 5.2. Consider the matrix  $M_{a,b} = \begin{bmatrix} S_a & b \\ -b^T & 0 \end{bmatrix}$ . Following the notation in Theorem 5.1, let:

$$\begin{aligned} k &= S_a^\dagger b, & h &= -(S_a^\dagger)^T b, & w_1 &= 1 + k^T k, & w_2 &= 1 + h^T h, \\ \alpha &= 0, & \beta &= b^T S_a^\dagger b, & u &= (I_7 - S_a S_a^\dagger)b, & v &= -(I_7 - S_a^\dagger S_a)b. \end{aligned}$$



Then  $\beta = 0$  and the subsequent equalities hold:

$$k = h = \begin{cases} 0 & \text{if } a = 0 \\ \frac{1}{a^T a} S_a^T b & \text{if } a \neq 0 \end{cases},$$

$$w_1 = w_2 = \begin{cases} 1 & \text{if } a = 0 \\ 1 + \frac{(a^T a)(b^T b) - (a^T b)^2}{(a^T a)^2} & \text{if } a \neq 0 \end{cases}, \quad u = -v = \begin{cases} b & \text{if } a = 0 \\ \frac{a^T b}{a^T a} a & \text{if } a \neq 0 \end{cases}.$$

*Proof.* The equalities hold trivially when  $a = 0$ . So, assume that  $a \neq 0$ . By the properties of  $S_a$  in Proposition 3.1, we have

$$k = S_a^\dagger b = -\frac{1}{a^T a} S_a b = \frac{1}{a^T a} S_a^T b,$$

$$h = (S_a^\dagger)^T (-b) = -\frac{1}{a^T a} S_a^T (-b) = k,$$

$$w_1 = 1 + k^T k = 1 - \frac{1}{(a^T a)^2} b^T S_a^2 b = 1 + \frac{b^T (a^T a) b - (a^T b)^T (a^T b)}{(a^T a)^2} = 1 + \frac{(a^T a)(b^T b) - (a^T b)^2}{(a^T a)^2},$$

$$w_2 = 1 + h^T h = 1 + k^T k = w_1,$$

$$\beta = b^T S_a^\dagger b = -\frac{1}{a^T a} b^T S_a b = \frac{1}{a^T a} b^T S_b a = -\frac{1}{a^T a} (S_b b)^T a = 0,$$

$$u = (I_7 - S_a S_a^\dagger) b = \left( I_7 + \frac{1}{a^T a} S_a^2 \right) b = \left( I_7 + \frac{1}{a^T a} (a a^T - a^T a I_7) \right) b = \frac{a^T b}{a^T a} a,$$

$$v = -(I_7 - S_a^\dagger S_a) b = -(I_7 - S_a S_a^\dagger) b = -u. \quad \square$$

**THEOREM 5.3.** Consider the matrix

$$M_{a,b} = \begin{bmatrix} S_a & b \\ -b^T & 0 \end{bmatrix}.$$

1) If  $a = 0$  and  $b \neq 0$ , then

$$M_{a,b}^\dagger = -\frac{1}{b^T b} \begin{bmatrix} 0_7 & b \\ -b^T & 0 \end{bmatrix}.$$

2) If  $a = b = 0$ , then  $M_{a,b}^\dagger = 0_8$ .

3) If  $a \neq 0$  and  $a^T b \neq 0$ , then

$$M_{a,b}^{-1} = -\frac{1}{a^T b} \begin{bmatrix} \frac{a^T b}{3a^T a} S_a + \frac{2}{3} S_b - \frac{1}{3a^T a} [S_a, S_a \times b] & a \\ -a^T & 0 \end{bmatrix}.$$

4) If  $a \neq 0$  and  $a^T b = 0$ , then

$$M_{a,b}^\dagger = -\frac{1}{a^T a + b^T b} \begin{bmatrix} (1 + \frac{b^T b}{3a^T a}) S_a - \frac{1}{3a^T a} [S_b, S_a \times b] & b \\ -b^T & 0 \end{bmatrix}$$

and a generalized inverse of  $M_{a,b}$  is

$$M_{a,b}^- = \begin{bmatrix} S_a^\dagger & a \\ -a^T & 0 \end{bmatrix}.$$

*Proof.* Suppose now that  $a = 0$  and  $b \neq 0$ . Then  $\text{rank}(S_a) = 0$  and  $\text{rank}(M_{a,b}) = 2$ . So, the case 1) is a consequence of 1) in Theorem 5.1 and Proposition 5.2.

The case 2) is straightforward.

As far as 3), assume that  $a \neq 0$  and  $a^T b \neq 0$ . Hence, by Proposition 5.2 and Theorem 4.2,  $u \neq 0$  and  $\det(M_{a,b}) \neq 0$ . Consequently,  $b$  does not belong to the column space of  $S_a$  and, so,  $-b$  does not belong to the column space of  $S_a^T$ . By Theorem 5.1, we have  $\text{rank}(M_{a,b}) = \text{rank}(S_a) + 2$ . Thus,  $\text{rank}(S_a) = 6$ . Also by the cited theorem,

$$M_{a,b}^{-1} = \begin{bmatrix} S_a^\dagger - ku^\dagger - (v^T)^\dagger h^T & (v^T)^\dagger \\ u^\dagger & 0 \end{bmatrix}.$$

Invoking Proposition 5.2, we conclude that:

$$\begin{aligned} u^\dagger &= \frac{u^T}{u^T u} = \frac{1}{a^T b} a^T, \\ ku^\dagger &= \frac{1}{a^T a} S_a^T b \frac{1}{a^T b} a^T = -\frac{1}{(a^T a)(a^T b)} S_a b a^T, \\ (v^T)^\dagger &= (-u^T)^\dagger = -(u^\dagger)^T = -\frac{1}{a^T b} a, \\ (v^T)^\dagger h^T &= -\frac{1}{a^T b} a \frac{1}{a^T a} b^T S_a = -\frac{1}{(a^T a)(a^T b)} a b^T S_a. \end{aligned}$$

From these equalities, we arrive at

$$M_{a,b}^{-1} = -\frac{1}{a^T b} \begin{bmatrix} -a^T b S_a^\dagger - \frac{1}{a^T a} (S_a b a^T + a b^T S_a) & a \\ -a^T & 0 \end{bmatrix}.$$

Applying the properties of  $S_a$  in Proposition 3.1, we obtain

$$\begin{aligned} S_a b a^T + a b^T S_a &= S_a b a^T - a b^T S_a^T \\ &= \frac{2}{3} S_{S_a S_a b} + \frac{1}{3} [S_a, S_{S_a b}] \\ &= \frac{2}{3} S_{a a^T b - a^T a b} + \frac{1}{3} [S_a, S_{a \times b}] \\ &= \frac{2}{3} (a^T b S_a - a^T a S_b) + \frac{1}{3} [S_a, S_{a \times b}]. \end{aligned}$$

Therefore,  $(a^T b) S_a^\dagger + \frac{1}{a^T a} (S_a b a^T + a b^T S_a) = -\frac{a^T b}{3 a^T a} S_a - \frac{2}{3} S_b + \frac{1}{3 a^T a} [S_a, S_{a \times b}]$ , and 3) follows.

In order to prove 4), suppose now that  $a \neq 0$  and  $a^T b = 0$ . Thus,  $\det(M_{a,b}) = 0$ . By Proposition 5.2, we have

$$k = h = -\frac{1}{a^T a} S_a b,$$

$$w_1 = w_2 = 1 + \frac{b^T b}{a^T a},$$

and

$$u = v = 0.$$

Moreover,  $b \in N(a^T)$  and  $N(a^T) = R(S_a)$  since  $R(S_a) = (N(S_a^T))^\perp = \langle a \rangle^\perp$ . Consequently,  $\text{rank}(M_{a,b}) = \text{rank}(S_a)$  and, by Theorem 5.1, we get

$$M_{a,b}^\dagger = \begin{bmatrix} S_a^\dagger - w_1^{-1} k k^T S_a^\dagger - w_2^{-1} S_a^\dagger h h^T & w_2^{-1} S_a^\dagger h \\ w_1^{-1} k^T S_a^\dagger & 0 \end{bmatrix} + \frac{k^T S_a^\dagger h}{w_1 w_2} \begin{bmatrix} k \\ -1 \end{bmatrix} \begin{bmatrix} h^T & -1 \end{bmatrix}.$$

Taking into account the properties in Proposition 3.1 and in Proposition 5.2, we have

$$\begin{aligned} \frac{k^T S_a^\dagger h}{w_1 w_2} &= \frac{h^T S_a^\dagger h}{w_1^2} = -\frac{b^T S_a S_a^\dagger S_a b}{(a^T a + b^T b)^2} = -\frac{b^T S_a b}{(a^T a + b^T b)^2} = 0, \\ k^T S_a^\dagger &= h^T S_a^\dagger = -\frac{1}{(a^T a)^2} b^T S_a^2 = \frac{-1}{(a^T a)^2} b^T (a a^T - a^T a I_7) = \frac{1}{a^T a} b^T, \\ S_a^\dagger h &= S_a^\dagger k = (k^T (S_a^\dagger)^T)^T = -(k^T S_a^\dagger)^T = -\frac{1}{a^T a} b, \end{aligned}$$

and

$$-k k^T S_a^\dagger - S_a^\dagger h h^T = \frac{1}{a^T a} (-k b^T + b k^T) = \frac{1}{(a^T a)^2} (S_a b b^T + b b^T S_a).$$

We also obtain

$$\begin{aligned} S_a b b^T + b b^T S_a &= S_a b b^T - b b^T S_a^T \\ &= \frac{2}{3} S_{S_b S_a b} + \frac{1}{3} [S_b, S_{S_a b}] \\ &= \frac{2}{3} S_{-S_b S_b a} + \frac{1}{3} [S_b, S_{a \times b}] \\ &= \frac{2}{3} S_{b^T b a - b b^T a} + \frac{1}{3} [S_b, S_{a \times b}] \\ &= \frac{2}{3} b^T b S_a + \frac{1}{3} [S_b, S_{a \times b}], \end{aligned}$$

$$\begin{aligned} S_a^\dagger + \frac{1}{(a^T a)^2 w_1} (S_a b b^T + b b^T S_a) &= \frac{-1}{a^T a + b^T b} \left( \frac{a^T a + b^T b}{a^T a} S_a - \frac{1}{a^T a} (S_a b b^T + b b^T S_a) \right) \\ &= \frac{-1}{a^T a + b^T b} \left( S_a + \frac{b^T b}{a^T a} S_a - \frac{2b^T b}{3a^T a} S_a - \frac{1}{3a^T a} [S_b, S_{a \times b}] \right) \\ &= -\frac{1}{a^T a + b^T b} \left( \left( 1 + \frac{b^T b}{3a^T a} \right) S_a - \frac{1}{3a^T a} [S_b, S_{a \times b}] \right), \end{aligned}$$

$$w_2^{-1} S_a^\dagger h = -\frac{1}{w_1 a^T a} b = -\frac{1}{a^T a + b^T b} b,$$

$$w_1^{-1} k^T S_a^\dagger = \frac{1}{w_1 a^T a} b^T = \frac{1}{a^T a + b^T b} b^T.$$

Hence, the first part of 4) follows. To finish the proof, observe that

$$\begin{bmatrix} S_a & b \\ -b^T & 0 \end{bmatrix} \begin{bmatrix} S_a^\dagger & a \\ -a^T & 0 \end{bmatrix} \begin{bmatrix} S_a & b \\ -b^T & 0 \end{bmatrix} = \begin{bmatrix} S_a S_a^\dagger S_a - S_a a b^T & S_a S_a^\dagger b \\ -b^T S_a^\dagger S_a & -b^T S_a^\dagger b \end{bmatrix} = M_{a,b}$$

since

$$\begin{aligned} S_a S_a^\dagger S_a - S_a a b^T &= S_a, \\ S_a S_a^\dagger b &= \frac{-1}{a^T a} S_a^2 b = \frac{-1}{a^T a} (a a^T b - a^T a b) = b, \\ -b^T S_a^\dagger S_a &= \frac{1}{a^T a} b^T S_a^2 = \frac{1}{a^T a} (b^T a a^T - b^T a^T a) = -b^T, \end{aligned}$$

and

$$-b^T S_a^\dagger b = \frac{1}{a^T a} b^T S_a b = -\frac{1}{a^T a} b^T S_b a = \frac{1}{a^T a} (S_b b)^T a = 0. \quad \square$$

**6. Rotation matrices from  $S_a$  and  $M_{a,b}$ .** Possible representations for rotation operators are the ones in the form of rotation matrices. In particular, the Cayley transform and the exponential of a skew-symmetric matrix may be considered.

Let us begin with the Cayley transform of  $S_a$  and with the Cayley transform of  $M_{a,b}$ , writing the latter in terms of the former one. With this purpose in mind, we first recall the following result.

PROPOSITION 6.1. [13] *Let  $E \in \mathbb{R}^{r \times r}$ ,  $F \in \mathbb{R}^{r \times s}$ ,  $G \in \mathbb{R}^{s \times r}$ ,  $H \in \mathbb{R}^{s \times s}$  and*

$$N = \begin{bmatrix} E & F \\ G & H \end{bmatrix}.$$

*If  $E$  and  $J = H - GE^{-1}F$ , the Schur complement of  $E$  in  $N$ , are invertible, then*

$$N^{-1} = \begin{bmatrix} E^{-1} + E^{-1}FJ^{-1}GE^{-1} & -E^{-1}FJ^{-1} \\ -J^{-1}GE^{-1} & J^{-1} \end{bmatrix}.$$

THEOREM 6.2. *The Cayley transform of  $M_{a,b}$  is the rotation matrix*

$$\mathcal{C}(M_{a,b}) = \begin{bmatrix} \mathcal{C}(S_a) - \frac{2}{s} S^{-1} b b^T S^{-1} & -\frac{2}{s} S^{-1} b \\ \frac{2}{s} b^T S^{-1} & \frac{2}{s} - 1 \end{bmatrix},$$

*where  $S$  stands for  $S_a + I_7$ ,  $s$  is the Schur complement of  $S$  in  $I_8 + M_{a,b}$  and  $\mathcal{C}(S_a)$  is the Cayley transform of  $S_a$  given by the rotation matrix*

$$\mathcal{C}(S_a) = \frac{1}{1 + a^T a} (-2S_a + 2a a^T + (1 - a^T a)I_7).$$

*Proof.* Let us denote  $S_a + I_7$  by  $S$ . Invoking the proof of Theorem 4.4, we have

$$S^{-1} = -\frac{1}{1 + a^T a} (S_a - I_7 - a a^T).$$

As  $\mathcal{C}(S_a) = 2S^{-1} - I_7$ , then the stated formula for  $\mathcal{C}(S_a)$  follows. Furthermore, the Schur complement  $1 + b^T S^{-1} b$  of  $S$  in  $I_8 + M_{a,b}$  is equal to

$$s = \frac{1 + a^T a + b^T b + (a^T b)^2}{1 + a^T a}.$$

and, so, is invertible. By Proposition 6.1, we obtain

$$(I_8 + M_{a,b})^{-1} = \frac{1}{s} \begin{bmatrix} sS^{-1} - S^{-1}bb^T S^{-1} & -S^{-1}b \\ b^T S^{-1} & 1 \end{bmatrix}.$$

Taking into account that  $\frac{2}{s}(sS^{-1} - S^{-1}bb^T S^{-1}) - I_7 = \mathcal{C}(S_a) - \frac{2}{s}S^{-1}bb^T S^{-1}$  and  $\mathcal{C}(M_{a,b}) = 2(I_8 + M_{a,b})^{-1} - I_8$ , we arrive at the stated matrix for  $\mathcal{C}(M_{a,b})$ .  $\square$

An explicit expression for computing the exponential of an order 3 skew-symmetric matrix  $B$  is given by the Rodrigues' formula, a consequence of  $B^3 = -\alpha^2 B$  for a certain scalar  $\alpha$ . Although this does not hold in general for an order  $n \geq 4$ , hypercomplex matrices are an exception, [9]. Moreover, a generalization of the Rodrigues' formula that allows to compute the exponential of a skew-symmetric matrix of order  $n \geq 3$  was proposed in [5].

**THEOREM 6.3.** [9] *Let  $\underline{a} = a_0 + a \in \mathbb{O}$  with  $\|a\| = \alpha \neq 0$ . Then*

$$e^{tS_a} = I \cos(\alpha t) + S_a \frac{\sin(\alpha t)}{\alpha} + \frac{1 - \cos(\alpha t)}{\alpha^2} aa^T.$$

**THEOREM 6.4.** [5] *Given any non-null skew-symmetric  $n \times n$  matrix  $B$ , where  $n \geq 3$ , if the set of distinct eigenvalues of  $B$  is  $\{i\theta_1, -i\theta_1, \dots, i\theta_p, -i\theta_p\}$ , where  $\theta_j > 0$  and each  $i\theta_j$  (and  $-i\theta_j$ ) has multiplicity  $k_j \geq 1$ , there are  $p$  unique skew-symmetric matrices  $B_1, \dots, B_p$  such that*

$$B = \theta_1 B_1 + \dots + \theta_p B_p, \quad B_i B_j = B_j B_i = 0_n (i \neq j), \quad B_i^3 = -B_i$$

for all  $i, j$  with  $1 \leq i, j \leq p$ , and  $2p \leq n$ . Furthermore,

$$e^B = e^{\theta_1 B_1 + \dots + \theta_p B_p} = I_n + \sum_{i=1}^p (\sin \theta_i B_i + (1 - \cos \theta_i) B_i^2).$$

**THEOREM 6.5.** *Let  $a, b \in \mathbb{R}^7$  such that  $a \neq 0_{7 \times 1}$ . The exponentials of  $S_a$  and of  $M_{a,b}$  are, respectively, the rotation matrices*

$$e^{S_a} = I_7 + \frac{\sin \|a\|}{\|a\|} S_a + \frac{1 - \cos \|a\|}{\|a\|^2} S_a^2$$

and

$$e^{M_{a,b}} = I_8 + \sum_{k=1}^p (\sin \theta_k M_{a,b,k} + (1 - \cos \theta_k) M_{a,b,k}^2),$$

where

$$p = \begin{cases} 2 & \text{if } a^T b = 0 \\ 3 & \text{if } a^T b \neq 0 \end{cases}, \quad \{\theta_j : 1 \leq j \leq p\} = \begin{cases} \{\|a\|, \|a + b\|\} & \text{if } p = 2 \\ \{\|a\|, \sqrt{\frac{1}{2}(\|a\|^2 + \|b\|^2 \pm \|a - b\| \|a + b\|)}\} & \text{if } p = 3 \end{cases}$$

and the  $p$  unique skew-symmetric matrices  $M_{a,b,k}$  can be obtained through the solution of a  $28p \times 28p$  linear equations system deduced from

$$M_{a,b} = \sum_{k=1}^p \theta_k M_{a,b,k}, \quad M_{a,b}^3 = - \sum_{k=1}^p \theta_k^3 M_{a,b,k}, \quad \dots, \quad M_{a,b}^{2p-1} = (-1)^{p-1} \sum_{k=1}^p \theta_k^{2p-1} M_{a,b,k}.$$

*Proof.* Let  $a, b \in \mathbb{R}^7$  such that  $a \neq 0_{7 \times 1}$ .

From 4) in Proposition 3.1, we have  $aa^T = S_a^2 + \|a\|^2 I_7$ . Hence, by Theorem 6.3, we obtain the stated Rodrigues-like formula for the exponential of  $S_a$ .

By Theorem 6.4, we obtain the stated formulas for the exponential of  $M_{a,b}$  and its odd powers, where  $\{\pm\theta_j i : \theta_j > 0, 1 \leq j \leq p\}$  is the set of distinct non-null eigenvalues of  $M_{a,b}$ . From Theorem 4.2, we have  $\det(M_{a,b}) = (a^T a)^2 (a^T b)^2$ . If  $a^T b = 0$  then  $M_{a,b}$  has, at least, an eigenvalue equal to 0 and  $b \neq -a$ . By Corollary 4.6, we obtain  $\theta_1 = \|a\|$  and  $\theta_2 = \|a + b\|$ . Hence,  $p = 2$ . If  $a^T b \neq 0$  then all eigenvalues of  $M_{a,b}$  are different from 0. Thus,  $p = 3$ . Concretely, once again by Corollary 4.6, we get  $\theta_1 = \|a\|$ ,  $\theta_2 = \sqrt{\frac{1}{2}(\|a\|^2 + \|b\|^2 - \|a - b\| \|a + b\|)}$ ,  $\theta_3 = \sqrt{\frac{1}{2}(\|a\|^2 + \|b\|^2 + \|a - b\| \|a + b\|)}$ .  $\square$

The generalization in [5] is theoretically interesting, however, according to [3], its computational cost seems prohibitive unless  $n$  is small. See [3] for details on effective methods for performing the computation of the exponential of a skew-symmetric matrix.

**Acknowledgment.** The authors thank the referee for bringing references [7] and [16] to their attention. A special thanks to M. Garfield for clarifying some English doubts.

#### REFERENCES

- [1] A. Ben-Israel and T. Greville. *Generalized inverses*. Springer-Verlag, New York, 2003.
- [2] T. Bröcker and T. Dieck. *Representations of Compact Lie Groups*. Springer, New York, 1985.
- [3] J. Cardoso and F. Leite. Exponentials of skew-symmetric matrices and logarithms of orthogonal matrices. *Journal of Computational and Applied Mathematics*, 233:2867–2875, 2010.
- [4] A. Elduque. Vector cross products. Talk presented at the Seminario Rubio de Francia of the Universidad de Zaragoza, <http://www.unizar.es/matematicas/algebra/elduque/Talks/crossproducts.pdf>, 2004.
- [5] J. Gallier and D. Xu. Computing exponentials of skew-symmetric matrices and logarithms of orthogonal matrices. *International Journal of Robotics and Automation*, 17:10–20, 2002.
- [6] F. Gantmacher. *The Theory of Matrices*. AMS Chelsea Publishing, Providence, Rhode Island, 2000.
- [7] J. Groß, S.-O. Troschke, and G. Trenkler. Problem 25-1: Moore-Penrose inverse of a skew-symmetric matrix. *IMAGE*, 25:16, 2000.
- [8] N. Jacobson. Composition algebras and their automorphisms. *Rendiconti del Circolo Matematico di Palermo*, 7:55–80, 1958.
- [9] F.S. Leite. The geometry of hypercomplex matrices. *Linear and Multilinear Algebra*, 34:123–132, 1993.
- [10] F.S. Leite and P. Crouch. The triple cross product in  $\mathbb{R}^7$ . *Reports on Mathematical Physics*, 39:407–424, 1997.
- [11] K.L. McDonald. Warping the universal extra dimensions. *Physical Review D*, 80:24–38, 2009.
- [12] C.D. Meyer. The Moore-Penrose inverse of a bordered matrix. *Linear Algebra and its Applications*, 5:375–382, 1972.
- [13] C.D. Meyer. *Matrix Analysis and Applied Linear Algebra*. SIAM, Philadelphia, 2000.
- [14] S. Morier-Genoud and V. Ovsienko. Orthogonal designs and a cubic binary function. *IEEE Transactions on Information Theory*, 59:1583–1589, 2013.
- [15] C. Senanayake and S. Senanayake. A computational method for reliable gait event detection and abnormality detection for feedback in rehabilitation. *Computer Methods in Biomechanics and Biomedical Engineering*, 14:863–874, 2011.
- [16] G.P.H. Styan and H.J. Werner (editors). Problem 25-1 with solutions. *IMAGE*, 26:2–4, 2001.
- [17] G. Trenkler and D. Trenkler. The vector cross product and  $4 \times 4$  skew-symmetric matrices. In: C.R. Rao, H. Toutenburg, and H.C. Shalabh (editors), *Recent Advances in Linear Models and Related Areas*, Springer, Heidelberg, 95–104, 2008.